

**HEREDITY OF THE
BLOOD GROUPS**



KARL LANDSTEINER
Father of Immunohematology
(1868-1943)

HEREDITY OF THE BLOOD GROUPS

By ALEXANDER S. WIENER, A.B., M.D., F.A.C.P., F.C.A.P.

Senior Bacteriologist (Serology) to the Office of the Chief Medical Examiner of New York City, Assistant Professor, Department of Forensic Medicine, New York University Postgraduate Medical School, Attending Immunohematologist, Jewish and Adelphi Hospitals, Brooklyn

and IRVING B. WEXLER, A.B., M.D., F.A.C.P.

Associate Pediatrician, Jewish Hospital of Brooklyn, Associate Immunohematologist, Jewish and Adelphi Hospitals Brooklyn, Assistant Clinical Professor of Pediatrics, State University of New York College of Medicine at New York City

Sole Distributors in
India, Pakistan, Burma & Ceylon,
THE KOTLIER BOOK DEPOT
BOMBAY 12



GRUNE & STRATTON • New York and London

1958

IRONDISPIECF—Karl Landsteiner



PREFACE

I INTRODUCTION	1
Principles of blood group serology	1
History and present status of the blood groups	4
Principles of blood group genetics	12
A-B O groups	12
M N types	13
Rh Hr types	13
Nomenclature	14
II THE A-B O BLOOD GROUPS AND LEWIS TYPES	16
Landsteiner's A B O groups	16
Serology	16
Gene frequency analysis	16
Family studies	18
Subgroups of A	21
Gene frequency analysis	21
Serology	23
Family studies	24
Subgroups A ₁ and A ₂ B	26
Other subgroups	26
Factor C	26
Serologic complexity of the A-B O agglutinogens	28

Library of Congress Catalog Card Number 58-10790

Copyright © 1958
Alexander S Wiener, M D
Brooklyn, N Y

Printed in U S A (B)

Group specific substances in body fluids and secretions	29
Secretors and non secretors	29
Technique	30
Lewis blood group	31
Serology	31
Family studies	32
Theoretico statistical considerations	34
A B O groups in apes and monkeys	38
Anomalies of the A B O blood groups	38
III THE M N-S SYSTEM	41
M N types	41
Serology	41
Heredity	42
Gene frequency analysis	43
Mother child combinations	45
Agglutinin N_2	46
Agglutinogens M and N in monkeys and apes	46
Factors S s and U	48
Factor S	48
Nomenclature	51
Gene frequency analysis	52
Family investigations	53
Factor U	54
Other complexities of the M N-S system	55
IV THE P SYSTEM	57
Serology	57
Genetics	57
Factor T_J^s	58
V THE Rh Hr SYSTEM	60
The Rh_o factor	60
Distribution	60
Family studies	61
Serology	61
Factor rh	62
Linked genes vs multiple alleles	63
Eight Rh types	64
Factor rh''	64

The eight Rh phenotypes	65
Genetic theory	66
Gene frequency analysis	66
Family studies	70
Serology	70
Hr factors	71
Relationship between Hr and Rh factors	71
Nomenclature	74
Family studies	77
Population studies	78
Serology	79
Additional Rh Hr factors	80
Factor rh^w	80
Factor hr	80
Factor hr^v	81
Complexities of the Rh_o factor and associated factors	84
Factor Rh_o	85
Nomenclature	86
Factor Rh_o	87
Theoretical serological considerations	88
Factors Rh^A Rh^B Rh^C	90
Additional complexities of the Rh Hr types	91
C D E notations for the Rh Hr types	92
Historical background	92
Crossing over	95
The C D E notations	100
VI THE KELL BLOOD GROUPS	103
Factor K (kell)	103
Factor k (Cellano)	104
Additional K k factors	105
Serology	107
VII THE LUTHERAN BLOOD GROUPS	108
Family studies	108
Distribution of Iu^A	109
Factor Iu^B	109
VIII THE DUFFY BLOOD GROUPS	110
Terminology	110
Serology	110

Distribution	110
Heredity	111
Factor f	111
IX THE KIDD BLOOD GROUP	112
Nomenclature	112
Distribution	112
Factor j	112
Serology	112
X OTHER BLOOD GROUP SYSTEMS	114
Diego system	114
High frequency blood factors	115
Low frequency blood factors	116
XI MEDICOLEGAL APPLICATIONS	117
Chances of excluding paternity	118
A B O groups	120
M N types	120
Rh Hr types	121
Disputed maternity	123
More complex problems	124
Circumstantial evidence of paternity	125
Case reports	125
Nomenclature	128
Qualifications of experts	129
XII ANTHROPOLOGICAL STUDIES ON THE BLOOD TYPES	132
Classification of human races by blood group studies	132
Effect of selection against heterozygotes	134
Analysis of mixed populations	135
REFERENCES	138
INDEX	147

Preface

DURING the past 15 years, knowledge of the blood groups has progressed with great rapidity, and blood grouping has become a fundamental biological science with applications in clinical and legal medicine, anthropology, human genetics, evolution, tissue transplantation, and veterinary medicine.

This rapid development followed the discovery of the Rh factor by Landsteiner and Wiener and the subsequent demonstration of its clinical importance in transfusion and in isoimmunization by pregnancy. When certain clinical problems could not be solved by the use of the classical agglutination techniques, new methods of blood grouping were devised which disclosed the existence of a second and more important form of antibody, namely, the so-called blocking or "univalent" antibody. These new and more sensitive techniques, the conglutination test, the anti globulin test, and the proteolytic enzyme test became the means for recognizing instances of isoimmunization to other blood factors so that up to present time more than 30 blood factors belonging to 9 different blood group systems have been discovered. The purpose of this monograph is to present in a unified manner the facts known about the heredity of the human blood groups.

The data that the specialist in immunogenetics must master are numerous and exceedingly complex, but fortunately the fundamental concepts are relatively simple. The reader will find that once he has mastered these few fundamental principles, he will have no difficulty in understanding and learning the detailed facts. For example, in all cases heredity is by multiple allelic genes, not only in the ABO system, but also in the Rh Hr and MN-S systems, and even in the case of the complex B blood groups of cattle and other blood group systems in animals. Therefore, emphasis is placed on principles throughout the text, and the analogies in the heredity of the various blood groups systems are pointed out. Blood grouping is primarily a subspecialty of immunology, so that for mastery of the field thorough understanding of the nature of serological reactions in general is essential. Therefore, at the very onset, the important distinction between a blood factor and agglutinin is pointed out, and this concept is applied to explain the serology, genetics, and nomencla-

ture of the A B O, Rh Hr, M N S, and other blood group systems. This basic concept arises from the serological observation that antigens in general have multiple corresponding antibodies so that there is no simple one to one correspondence between antigens and antibodies. The emphasis placed on this principle is unique among monographs on the subject, and, in the opinion of the authors, serves to resolve the contradictions and paradoxes which have inevitably arisen from disregard of the principle.

As Darwin has stated false theories do little harm, while false facts may retard the progress of science. In fact, an incorrect theory may prove helpful so long as it is used only as a blueprint or working hypothesis to give a direction to investigative work. Therefore a debt of gratitude is due to British investigators, notably R. R. Race and R. Sanger, whose observations and ideas have acted as a stimulus to investigators in the field for the past 15 years. However if an investigator refuses to relinquish a theory refuted by readily available facts or if he allows his preconceived ideas to color his judgment in reporting observations, serious errors may result. This has indeed occurred in the case of the Rh Hr types, as exemplified by the erroneous reports of discovery of anti d, reports of non specific rises in maternal Rh antibody titers during pregnancy with Rh negative fetuses etc. The literature of the blood groups has been swamped with reports by poorly trained and inexperienced workers, unfamiliar with Landsteiner's and other basic work in the field. Such workers have the confidence born of ignorance, and are insensitive to the possibility of human error, so they fail to use the blind technique which is so essential for accurate results, as when carrying out the delicate tests for the Lewis blood types. Since the results which are most difficult to explain are those which are not true, an attempt has been made to include in this monograph only those reports which appear to be reliable. For any errors of omission or commission the authors assume full responsibility, and they would appreciate having their attention called to any such errors which may come to the attention of readers.

Readers interested in other aspects of blood grouping are referred to the senior author's 'Blood Groups and Transfusion,' the third and last edition of which was published in 1943. That book covers the earlier literature on all aspects of blood grouping and its applications, but to do justice to the subject would now require a large volume or volumes, in view of the great growth of the field during the past 15

years. For more detailed information regarding the Rh-Hr types and their applications, the reader may consult the "Rh-Hr Types," published in 1934, a collection of the more important contributions of Wiener and his collaborators, together with a review of the literature. A simplified, compact but comprehensive account can be found in Wiener's "Rh-Hr Syllabus," which has been published also in German, Italian, Spanish, and Japanese.

The serological principles underlying blood group reactions were first pointed out by Karl Landsteiner, the father of the field of immunohematology, who received the Nobel Prize in Medicine in 1930 for the discovery of the blood groups. The senior author was fortunate to be closely associated with Karl Landsteiner from 1929 until his death in 1943. During that period, he had the advantage of Landsteiner's advice and help as mentor, collaborator, and personal friend. In appreciation of Karl Landsteiner's pathfinding contributions to the field, and as a token of personal gratitude, the authors dedicate this book to his memory on this, the 90th anniversary of his birth.

March 1958

A. S. WIENER

I. B. WEINER

CHAPTER I

Introduction

More data are available concerning blood group differences than any other heritable characteristic of healthy human beings. The reasons for this are manifold. The techniques of blood grouping have been perfected during the past two decades to a point where the blood groups of any individual can be readily and precisely determined. Moreover, individual blood group differences have no selective value aside from the effect of isosensitization on the fetus *in utero* so that polymorphism is the rule in almost all human populations. The blood groups are determinable at birth and remain unchanged throughout life, unaffected by age, diet, radiation or disease. Because of the implications of the blood groups in clinical medicine for the selection of donors for blood transfusion and for isoimmunization in pregnancy, millions of blood grouping tests are performed every year. Furthermore, because every transfusion and every pregnancy is a possible source of isosensitization, new blood group antibodies are constantly being discovered and the heredity of the corresponding blood factors worked out.¹

Principles of Blood Group Serology

For blood grouping tests, blood is obtained, preferably from a vein and separated into its two gross components which are present in approximately equal proportions, namely the red blood cells and the plasma in which the red blood cells are suspended within the circulation. (Serum is the fluid which is expressed from clotted blood—that is, plasma without the fibrinogen.)

The red cell or erythrocyte has a lipoprotein envelope which from its serological behavior appears to have a regularly patterned mosaic structure. Presumably certain chemical structures named agglutinogens because of the nature of the serological tests used to demonstrate their presence, are repeated about the surface of the erythrocyte. The first discovered of these agglutinogens were the so-called ABO substances which determine the four classic blood groups described by Karl Landsteiner² at the turn of the century. Two other sets of

agglutinogens are the M N substances and the P p substances both of which were discovered by Landsteiner and Levine³ about three decades ago. Aside from the A B O substances, the most important are the Rh Hr substances, discovered by Landsteiner and Wiener⁴ about two decades ago. The discovery of the Rh Hr substances led to new methods of blood typing and as a result during the past two decades the literature has been rich with reports of the discovery of other blood group substances such as K k (Kell), F f (Duffy) and J j (Kidd). Figure 1 is a hypothetical diagram of the mosaic arrangement of the blood group substances as they may be visualized on the surface of the red blood cell.

To determine an individual's blood group, a battery of diagnostic reagents or antiserums anti A, anti B, anti M, anti N, anti Rh₀, anti rh', etc. are required. These reagents contain only antibodies specific for the blood factor in question and no other antibodies. These antiserums can be obtained in a variety of ways. Anti A and anti B occur spontaneously in the serums of healthy adults whose red blood cells lack the corresponding agglutinogen. Since the spontaneously occurring antibodies are generally weak, donors are injected with




FIG 1—Diagrammatic Representation of the Distribution of Blood Group Agglutinogen Loci on the Surface of the Erythrocyte. Legend: ● = A B O Haptens, X = M N Haptens, ⊙ = Rh Hr Haptens, • = Haptens of other specificities human species specific haptens

materials of animal origin which contain A and B like substances and in that way antisera of high titer and potency can be obtained. Anti M and anti N sera are produced by immunizing rabbits. Most of the other antisera are obtained from mothers of erythroblastic babies, or from patients who have had hemolytic transfusion reactions, or from deliberately immunized blood donors.⁵ Recently it has been found⁶ that certain seeds contain proteins which have blood group specificity, namely, anti A, anti A₁, anti B, anti H, and anti N, but such reagents have not yet found wide usage. Most experts in the field prepare their own antisera, but satisfactory reagents are also available commercially. It may be of interest to mention that using all available reagents as many as thirty blood factors of different specificities have been identified to date.

The blood factors detected by the use of the various antisera are not all independent of one another, but fall naturally into sets which are known as blood group systems. Whether or not two blood factors are independent of one another or are members of a common blood group system can be determined from their distribution in the general population, by using a 2×2 contingency table.* For example in the

* In order to determine whether two traits X and Y are independent of one another a table is set up as follows:

Trait	X+	X-	Totals
Y+	a	b	a + b
Y-	c	d	c + d
Totals	a + c	b + d	N

In this table a, b, c and d represent the frequencies of the four possible combinations of the traits X and Y and $N = a + b + c + d$ the total number of individuals examined. If X and Y are independent then the frequency of the Y trait among X+ individuals should be the same as its frequency amongst

X- individuals that is $\frac{a}{a+c} = \frac{b}{b+d}$ so that $ad = bc$ or $ad - bc = 0$.

In cases where $ad - bc$ is not equal to zero whether the difference from zero is statistically significant can be determined by deriving the value of chi square with the aid of the following formula:

$$\chi^2 = \frac{(ad - bc)^2 N}{(a+b)(a+c)(b+d)(c+d)}$$

The number of degrees of freedom is taken equal to one when determining the value of P from the table of chi square.⁷

TABLE I
HISTORY AND PRESENT STATUS OF THE HUMAN BLOOD GROUPS

Blood system	Serology		Genetics		
	Blood factors and blood groups	Discoverers	Year	Nature of contributions	Investigators
A B O and Lewis	Factors A and B	Landsteiner	1900	First family study	Landsteiner and Ottenberg
	Blood groups O A and B	Landsteiner	1901	Theory of independent gene pairs	von Dungern and Hirsfeld
	Blood group AB	DeCastello and Sturli	1902	Theory of multiple alleles	Bernstein
	A ₁ A ₂ subgroups	von Dungern and Hirsfeld	1911	Extension of multiple allele theory to include subgroups A ₁ A ₂	Landsteiner and Levine
	Factor F _A common to human agglutinogen A and sheep blood	Schiff and Adelsberger	1924		Thomsen Friedenreich and Worsaae
	Factors B, B ₁ B ₂ etc. of agglutinogen B	Friedenreich and With	1933		
	Factor C shared by agglutinogens A and B	Hooker and Anderson	1921		
	Factor H of agglutinogens O and A ₂	Landsteiner and With	1926		
	Agglutinogen A ₃	Schiff	1927		
	Agglutinogen A ₄	Fischer and Hahn	1935	Hereditary agglutinogen A ₃	Friedenreich
		Gammelgaard and Marckussen	1940		

M N S	Group specific seed lectins	Renkonen	1948	Secretor and non secretor genes	Schiff and Siskind	1932
	Secretor type blood group substances in body fluids	Boyd and Reguiera Jehrs and Putkonen	1949 1950			
	Substance T in non secretor saliva	Uyeyama	1950			
	Lewis blood types	Mourant Andresen	1946 1948	Relationship between Lewis and secretor types	Grubb	1951
	Blood group A_0 Blood group A_m	Dunsford and Aspinall Wiener	1952 1957			
	Factors M and N and the three M N types	Landsteiner and Levine Crome	1927 1935	Heredity by pair of allelic genes Heredity of agglutino-gen N_2	Landsteiner and Levine Friedenreich	1928 1936
	Rare agglutinogen N_2					
	Factors M M M ₁ etc	Wiener and Landsteiner	1937			
	Factors N, N	Wiener	1937			
	Factor S	Walsh and Montgomery	1947	Relationship of S to M N system demonstrated	Sanger and Race	1947
	Factor s	Levine Kuhmichel Wigod and Koch	1951			
	Factor t	Wiener Unger and Gordon	1953	Heredity as simple Mendelian dominant and probable relation to M N types	Wiener Unger and Cohen	1954
				Relation to M N types proved	Greenwalt et al	1955

TABLE 1—Continued

Blood Group System	Serology			Genetics		
	Blood factors and blood groups	Discoverers	Year	Nature of contribution	Investigators	Year
P	Factor Hu	Landsteiner, Strutton and Chase	1934	Relationship to MN types noticed	Wiener	1930
	Factor He	Hin and Mourant	1931	Relationship to MN types demonstrated	Chalmers, Hin and Mourant	1953
	Factor Me*	Levine, Stock, Kuhnel and Bronikovsky	1950	Relationship of Me* and Gr to the MN types demonstrated	Wallace et al	1950
	Factor Gr	Graydon*	1946			
P	Factor P	Landsteiner and Levine	1927	Heredity is simple Mendelian dominant	Landsteiner and Levine	1930
	Factor Tj*	Levine, Bollitt, Waller and Kuhmichel	1951	Relationship of Tj* to P* system demonstrated	Sanger and Race	1955
Rh Hr	First anti rhesus (Rh ₀) serum prepared, Rh positive and Rh negative types	Landsteiner and Wiener	1937	Heredity of Rh ₀ factor as simple Mendelian dominant	Landsteiner and Wiener	1941
	Rh sensitization as cause of transfusion hemolysis	Wiener and Peters	1939			
	Rh sensitization as cause of erythroblastosis fetalis	Levine, Burnham, Katz and Vogel	1941			

Factor rh	Wiener	1941	Heridity of 4 types determined by factors rh and Rh , by multiple allelic genes	Wiener and Landsteiner	1941-3
Factor hr'	Ievine and Javert	1941	Reciprocal relation of rh and hr'	Ievine, Race and Taylor	1941-3
Factor rh''	Wiener and Soann	1943	Eight Rh types determined by factors Rh , rh and rh'' , and their heredity by multiple allelic genes	Wiener	1943
Intermediate Rh Hr factors especially Mh_0	Wiener	1944	Extension of multiple allele theory to include Rh Hr variants rh'' , hr , etc	Wiener	1944-1957
Factor hr''	Mofrant	1945	Racial distribution of the Rh Hr types	Wiener and Landsteiner	1944-3
Factor rh''	Cullender and Race	1946			
Factor Mh_0	Race, Sanger and Selwyn	1950			
Factor hr	Rosenfield, Vogel, Gabel	1953			
Factor hr''	deVatale, Cahan, Joek, Race and Sanger	1955			
Factors Rh^A , Rh^B , Rh^C , etc	Wiener and Geiger	1956			
	Rosenfield, Hader and Gabel	1956			
	Unger and Wiener	1958			

TABLE I—Continued

Blood group system	Serology			Genetics		
	Blood factors and blood groups	Discoverers	Year	Nature of contribution	Investigators	Year
Kell (K k)	Factor K (or S_1)	Coombs, Mourant and Race	1946	Heredity as simple Mendelian dominant	Wiener and Gordon	1947
	Factor k	Wiener and Gordon Levine, Backer, Wigod and Ponder	1947 1949	Heredity of three K k types by pair of allelic genes	Levine et al	1949
	Factors Kp^a and Kp^b	Allen and Lewis	1957		Cutbush and Molli-son	1950
	Factor F	Cutbush, Molli-son and Parker	1950	Heredity		
Duffy (D d)	Factor f	Ikin, Mourant, Pettenkofer and Blumen-thal	1951			
	Factor I	Allen, Diamond and Nicols	1951	Heredity	Race, Sanger, Allen, Diamond and Nicols	1951
	Factor J	Plaut, Ikin, Mourant, Sanger and Race	1953			
	Factor Iu^a Subtypes	Callender and Race Mannering and Pickles	1946 1948	Heredity as simple Mendelian dominant	Callender and Race	1946
Lutheran (Lu)						

Diagnosis (Data)	Factor Data	Levine Koch McGee and Hill	1934	Heredity Racial distribution	Levine et al. Farnese and Arends	1934 1936
Unclassified	Factors of very high frequency in population					
	Yel	Simmons and Miller	1932			
	I	Wiener Unger and Cohen	1936			
	Y ₁ ^a	Fulton et al.	1937			
	Factors of very low frequency in population					
	Levas	Callender and Race	1946			
	Jellins	Callender	1947			
	Becker	Ellis and Prokop	1931			
	Yen	van Lijchem and van der Hart	1932			
	Re ^a	Davies and Stern	1933			
		Strausner and Spurr				
	C ₁	Wiener and Brancato	1933			

^a Recent work of Simmons et al. has demonstrated that factor Y₁ of Hart et al. (1934) is probably identical with factor C₁.

case of the four blood group O A B and AB the relationship $\bar{O} \times \bar{AB} = \bar{A} \times B$ does not hold which proves that factors A and B are not independent but belong to the same blood group system. The thirty-odd known blood factors have been shown to belong to some ten blood group systems corresponding to the different chemical substance or agglutino-gen that go to make up the mosaic of the surface of the red blood cell namely the A B O system the M N-S system the P p system the Rh Hr system etc (cf table 1). A situation similar to that occurring in man has been found to exist in regard to the blood groups of cattle fowl dogs mice rats and other animals.

The tests themselves are simple although considerable experience is required to obtain reliable and reproducible result since the reactions are delicate and have numerous pitfalls so that in unskilled hands they may be misinterpreted. Different methods are used depending on the reagent but the general principle is to bring the red blood cells into contact with the appropriate antiserum and allow them to react. If the cells are agglutinated or clumped into large masses generally visible to the naked eye this is a positive reaction whereas if the cells remain evenly suspended as seen under the microscope the reaction is negative (cf fig. 2).

It is necessary to emphasize the difference between a blood factor and an agglutino-gen.⁸ The agglutino-gen is the substance itself while the blood factors are the serological properties by which agglutino-gens are recognized. Thus when a blood is said to have the blood factor A this by definition means merely that the blood cells are clumped by anti A serum. It does not necessarily mean that the blood cells have the agglutino-gen A of human group A blood. As a matter of fact there are many agglutino-gens other than that of the human blood group A that react with particular anti A serum notably an agglutino-gen of sheep cells. Similarly some of the blood factors B which characterize the agglutino-gen B of human blood are shared by agglutino-gens of the red cells of rabbits and other rodents.

The nature of the combination between antibody and agglutino-gen can perhaps be best visualized by comparing it to the fitting of a lock by a key.⁹ The antibody is a modified serum gamma globulin which is indistinguishable by ordinary chemical means from the normal gamma globulin. It is believed that a small portion of the surface of the antibody molecule has been altered so as to make it an electro-



FIG. 2.—Microscopic Appearance of Blood Grouping Tests Magnified 150
 A Negative Reaction (No agglutination) B Positive Reaction (Showing intense agglutination of the red blood cells)

chemical counterpart of a part of the surface of the agglutinin molecule. In conformity with this concept it follows that not all antibodies capable of combining with an agglutinin would necessarily have the identical configuration any more than that all keys capable of opening a particular lock are identical. In fact there is evidence that antisera in general actually contain a spectrum of antibodies of related specificity as can be demonstrated by absorption experiments. For example, anti-M sera prepared by immunizing rabbits by injection of human type M cells often cross-react with blood cells of anthropoid apes and lower primates.⁹ By absorption experiments such serum can be fractionated into five or more components anti-M, M₁, M₂, etc., each specific for a corresponding M factor. Thus the human agglutinin M is characterized not by a single blood factor but by a multiplicity of blood factors. In fact the number of blood factors which characterize an agglutinin is theoretically unlimited, the only practical limitation being the resourcefulness and energy of the investigator in searching for and finding new antibodies. Despite the fact that each agglutinin has multiple blood factors, in many cases a single reagent may suffice for identification. For example, the

agglutinogens A and B can be identified by the corresponding serums anti A and anti B, respectively. The complex agglutinogens of the Rh Hr systems on the other hand, can be identified only by the use of a battery of anti-serums.

Table 1 summarizes the history and present status of the human blood groups.

Principles of Blood Group Genetics

A-B-O Groups. The hereditary nature of the blood groups was recognized shortly after Landsteiner's discovery. The first family studies were done by Epstein and Ottenberg¹⁰ in 1908, while in 1910 von Dungern and Hirschfeld¹¹ suggested that the four A B O groups are inherited by means of two independent pairs of allelic genes, Aa and Bb . In 1924, Felix Bernstein,¹² a mathematician, showed that this theory is incorrect, since it would require that in the general population the following relationship among the frequencies of the blood groups should hold, $\bar{O} \times \overline{AB} = \bar{A} \times B$. As has already been shown, this relationship does not hold, which not only proves that agglutinogens A and B belong to the same blood group system, but also that their heredity is not dependent on genes located in different chromosomes. Other workers suggested that the gene pairs Aa and Bb might be linked within the same chromosome pair,¹³ but as Bernstein pointed out, such a situation would yield the same result at equilibrium as independent assortment, although equilibrium is attained more slowly in the case of linkage.

According to Bernstein's theory, there are three allelic genes which have been variously designated as A , B , and O , or I^A , I^B , and I^O , respectively. (The base letter I was selected because the A B O blood grouping tests depend upon isoagglutination.) Accordingly, corresponding to the four blood groups there are six genotypes as follows: group O, genotype $I^O I^O$; group A, genotypes $I^A I^A$ and $I^A I^O$; group B, genotypes $I^B I^B$ and $I^B I^O$; and group AB, genotype $I^A I^B$. Despite Bernstein's mathematical demonstration of inheritance of the A B O blood groups by multiple allelic genes, the theory of linked genes enjoyed considerable vogue, until when no crossing over could be demonstrated this theory was abandoned. In 1927, Furuhata¹⁴ in Japan proposed a theory of complete linkage, postulating the existence of three completely linked gene pairs ab , Ab , and aB . Obviously, this theory is identical with the theory of multiple alleles, because all that

is necessary is to make the substitutions $I^A = Ab$, $I^B = aB$ and $I^O = ab$, which shows that the difference between the two concepts is merely a semantic one. Moreover, Furuhashi's postulation fails to account for the absence of a linked gene pair AB . Therefore, Furuhashi's theory is no longer mentioned in modern blood group literature. This is of interest because the concept of completely linked genes has recently been resurrected to explain the heredity of the Rh Hr blood types and the M N S types.

Landsteiner and Levine¹⁵ showed that the subgroups of group A and group AB are hereditary, and Thomsen, Friedenreich and Worsaele¹⁶ modified Bernstein's theory to include the subgroups by postulating four allelic genes instead of three, namely, I^A , I^{A_2} , I^B and I^O . To account for the rare agglutinin A_2 still another allele I^{A_2} had to be invoked and further extensions had to be made to take care of more recent discoveries.

M-N types. At first the heredity of the M N types appeared to be simpler than the heredity of the A B O blood groups. To account for the existence of only three M N types, Landsteiner and Levine¹⁷ postulated heredity by a pair of allelic genes designated M and N, or L^M and L^N , respectively. Thus, there are three genotypes corresponding to the three phenotypes as follows: type M, genotype $L^M L^M$; type N, genotype $L^N L^N$; and type MN, genotype $L^M L^N$. Sanger and Race,¹⁸ showed that the blood factor S discovered by Walsh and Montgomery¹⁹ is related to the M-N system, and postulated heredity by four completely linked gene pairs as follows, MS , M_s , NS , N_s . Wiener prefers to consider this another example of multiple allelism, and has designated the four allelic genes L^S , L , I^S , and I , respectively.²⁰ The blood factor U discovered by Wiener, Unger and Gordon,²¹ was shown to be related to the M-N system by Greenwalt et al.,²² since those rare individuals whose blood cells lack the factor U also lack the two factors S and s. Therefore, to allow for this rare type of blood it is necessary to extend the allelic series to include at least 6 allelic genes, L^S , L , L^s , I^S , I , and I^s , respectively. To allow for still other blood factors such as Henshaw and Hunter, corresponding extensions must be made in the series of allelic genes, as shown by Shapiro.²³

Rh-Hr Types. The most complicated example of multiple allelism in man is provided by the Rh Hr system. When only the original Rh₀ factor was known, Landsteiner and Wiener²⁴ showed that its inheritance could be explained

pair of allelic genes Rh and rh . With the discovery of the two related factors rh and rh Wiener²⁵ found it necessary to postulate a minimum of 6 allelic genes r r r R^0 R^1 and R^2 respectively. With the aid of family studies and the use of anti hr serum British investigators²⁷ succeeded in demonstrating the rare gene R^2 to which was later added the still rarer gene r^0 by Wiener.²⁸

Wiener²⁹ showed that each of the Rh factors has variants the most important of which are the variants of Rh_0 and that these variations are hereditary necessitating a further extension of the allelic series. Moreover discovery of still other Rh Hr factors rh^* hr hr hr^v Rh^A Rh^B etc. disclosed an ever increasing complexity of the Rh Hr system so that at the present time a minimum of 30 alleles must be postulated. Only in cattle has a situation been found more complex than that of the Rh Hr system. Here Stormont, Owen and Irwin³⁰ have described a blood group system with more than 100 alleles.

Probably the other blood group systems in man similarly depend upon multiple allelic genes for their hereditary transmission. For example the P blood types which were originally considered to be inherited by a pair of alleles P and p are now known to be more complex since Sanger³¹ showed that the rare Tj^+ negative type described by Levine³² depends on a corresponding rare allelic gene belonging to the P system. Moreover the recent observations of Allen³³ show that the K k system is also inherited by a complex series of multiple alleles. The same probably applies to the other blood group systems as will be shown when these are discussed individually in detail.

None of the blood group genes is known to be sex linked and the genes of each blood group system appears to be located on different pairs of chromosomes as is shown by their independent heredity. Thus each blood group system can be used as a marker for studies in human linkage. Other applications of the hereditary nature of the blood group systems are as a test for the zygosity of twins³⁴ and in medicolegal problems of disputed parentage.

Nomenclature

When devising symbols for the blood groups one must take into account the distinction between agglutinogens and blood factors and between genotypes and phenotypes. Therefore the convention has

been adopted of using regular type for symbols for agglutinogens and phenotypes, bold face type for symbols representing blood factors and their corresponding antibodies, while italics are used for the symbols representing genes and genotypes. For example, blood group A is characterized by the presence in the red cells of agglutinin A as evidenced by clumping of the red cells by anti A but not by anti B serum, which shows the presence in the cells of blood factor A, but not blood factor B. Moreover, agglutinin A is inherited by means of a corresponding allelic gene I^A so that corresponding to group A there are two theoretically possible genotypes, $I^A I^A$ and $I^A I^O$. The full implications of this terminology will become clearer as each individual blood group system is discussed.

CHAPTER II

The A-B-O Blood Groups and Lewis Types

Landsteiner's A-B-O Groups

Serology. According to Landsteiner's original theory, the four blood groups are determined by two agglutinogens A and B on the red blood cells and two corresponding naturally occurring iso agglutinins in the serum, anti A (alpha) and anti B (beta) respectively. Thus, the complete serological formula of the four blood groups is as follows $O\alpha\beta$, $A\beta$, $B\alpha$, and $AB\phi$. According to Landsteiner's law the alpha and beta agglutinins are regularly present in the serum when the corresponding agglutinogen is absent from the red cells except during the neonatal period, when the antibody producing mechanism is immature.

Gene Frequency Analysis. Bernstein's theory of the heredity of the A-B-O blood groups has already been described. The accuracy of the theory has been confirmed by family studies, and by statistical studies on the distribution of the blood groups in the general population. Bernstein's method of gene frequency analysis is as follows. Let p , q , and r represent the frequency of the three allelic genes I^A , I^B and I^O , respectively, so that $p + q + r = 1$, or 100 per cent. Then the frequencies of the four phenotypes can be expressed as follows.

Phenotypes	Genotypes	Frequencies
O	$I^O I^O$	r^2
A	$I^A I^A$ and $I^A I^O$	$p^2 + 2pr$
B	$I^B I^B$ and $I^B I^O$	$q^2 + 2qr$
AB	$I^A I^B$	$2pq$

Thus, if \bar{O} , \bar{A} , \bar{B} , and \bar{AB} represent the frequencies of the four blood groups in the general population, then $r^2 = \bar{O}$

$$\text{and } r = \sqrt{\bar{O}} \quad (1)$$

$$\text{Moreover, } p^2 + 2pr = \bar{A}$$

$$\text{so that } p^2 + 2pr + r^2 = (p + r)^2 = \bar{O} + \bar{A}$$

$$\text{Therefore, } p + r = \sqrt{\bar{O} + \bar{A}}$$

$$\text{so that } p = \sqrt{\bar{O} + \bar{A}} - \sqrt{\bar{O}} \quad (2)$$

$$\text{Similarly, } q = \sqrt{\bar{O} + \bar{B}} - \sqrt{\bar{O}} \quad (3)$$

$$\text{Since, } p + q + r = 1$$

$$\text{then } \sqrt{\bar{O} + \bar{A}} + \sqrt{\bar{O} + \bar{B}} - \sqrt{\bar{O}} = 1 \quad (4)$$

By substituting the frequencies of the blood groups A, B, and O of any population in formula (4) above, the validity of the theory of multiple alleles can be tested. Application of this test to large series of populations has served to substantiate Bernstein's theory. The test is only valid with populations at genetic equilibrium, that is, when there is random mating or panmixia. It can readily be shown that after a single generation of random matings the population will reach such equilibrium.²⁴

Actually, the formulae used by Bernstein himself in making this test are slightly different from the ones given above. Since $p + q + r = 1$, then the following set of formulae may be derived from formulae (1), (2), and (3) above,

$$r = \sqrt{\bar{O}} \quad (5)$$

$$p = 1 - \sqrt{\bar{O} + \bar{B}} \quad (6)$$

$$q = 1 - \sqrt{\bar{O} + \bar{A}} \quad (7)$$

In making the statistical test, Bernstein derived the values of p , q , and r from the formulae (5) to (7), and then calculated the deviation D of the sum from 100 per cent.

Thus,

$$D = 1 - (p + q + r) \quad (8)$$

To test whether the value of the deviation D is significant for a particular population, the standard error of D must be known. This Bernstein showed to be as follows

$$\sigma_D = \sqrt{\frac{pq}{2N(1-p)(1-q)}} \quad (9)$$

where N represents the number of individuals in the sample of the population being examined, and p and q are the calculated frequencies of genes I^A and I^B , respectively

For the purpose of testing the validity of the theory of multiple alleles, the original report of L and H Hirszfeld,³⁵ who were the first to demonstrate the differences in the distribution of the A-B O blood groups in different ethnic groups, is ideal. The data of these authors were collected during the first world war in the Balkans, six years before Bernstein proposed his genetic theory. The observations of L and H Hirszfeld and the results of Bernstein's statistical test of the theory of multiple alleles are given in table 2. It can readily be seen that results conform satisfactorily with the expectations under the theory.

In general, the calculated values of p , q and r would not be expected to add up to 100 per cent exactly. The values of these genes frequencies, as calculated by either set of formulae, while consistent, do not have the maximum efficiency in the statistical sense. The most efficient set of estimates can be obtained by the maximum likelihood method, or more simply by the formulae suggested by Bernstein,³⁶ as follows

$$p' = p \left(1 + \frac{D}{2} \right) \quad (10)$$

$$q' = q \left(1 + \frac{D}{2} \right) \quad (11)$$

$$r' = \left(r + \frac{D}{2} \right) \left(1 + \frac{D}{2} \right) \quad (12)$$

where p , q , and r , are the first estimates of the gene frequencies obtained with formulae (5) to (7), while p' , q' and r' are the improved estimates. These formulae are of particular value to anthropologists who are interested in population studies on the A-B O blood groups. The sum of the values, p' , q' , and r' will not equal 100 per cent exactly, but $1 - \frac{D^2}{4}$. Since D generally will be small, the difference

$\frac{D^2}{4}$ will be insignificant.

Family studies. For a satisfactory test of the theory of multiple alleles by family studies, the fact that group A and group B indi-

TABLE 2
STATISTICAL TEST OF BERNSTEIN'S THEORY
(Made with data accumulated by L. and H. Hirschfeld)

Race	Number of People	Frequencies of Groups				ϕ	q	r	$p + q + r$	Dev	P E	Dev P E
		O	A	B	AB							
English	500	0 464	0 434	0 072	0 031	0 268	0 052	0 681	1 001	0 001	0 0037	0 3
French	500	0 432	0 426	0 112	0 030	0 262	0 074	0 657	0 993	0 007	0 0037	1 9
Italians	500	0 472	0 380	0 110	0 038	0 237	0 077	0 687	1 001	0 001	0 0034	0 3
Serbians	500	0 380	0 418	0 156	0 046	0 268	0 107	0 516	0 991	0 009	0 0015	2 0
Greeks	500	0 382	0 416	0 162	0 040	0 262	0 107	0 618	0 987	0 013	0 0044	3 0
Bulgarians	500	0 390	0 406	0 142	0 062	0 271	0 108	0 624	1 003	0 003	0 0045	0 7
Arabs	500	0 432	0 324	0 190	0 050	0 209	0 129	0 660	0 998	0 002	0 0042	0 5
Turks (Macedonia)	500	0 368	0 380	0 186	0 066	0 256	0 136	0 607	0 999	0 001	0 0049	0 2
Russians	1000	0 107	0 312	0 218	0 063	0 210	0 152	0 638	1 000	0	0 0030	0
Spanish Jews	500	0 388	0 330	0 232	0 050	0 213	0 153	0 627	0 989	0 011	0 0017	2 3
Madagascan	400	0 458	0 262	0 237	0 045	0 168	0 154	0 675	0 997	0 003	0 0046	0 7
Senegal Negroes	500	0 432	0 224	0 292	0 050	0 149	0 189	0 657	0 995	0 005	0 0043	1 2
Annamese	500	0 420	0 224	0 284	0 072	0 161	0 108	0 648	1 008	0 008	0 0015	1 8
Hindus	1000	0 313	0 190	0 412	0 085	0 119	0 291	0 560	1 000	0	0 0040	0

After Wiener *Blood Groups and Transfusion*, 3rd ed., C. C. Thomas, Springfield Ill (1943)

individuals may be homozygous or heterozygous must be taken into account. If the distribution of the blood groups in the general population is known the proportion of homozygous and heterozygous individuals can be estimated from the frequency of the genotypes as follows

$$\begin{aligned}\text{Genotype } I^A I^A &= p^2 \\ \text{and genotype } I^A I^O &= 2pr\end{aligned}$$

Therefore

$$\text{the proportion of homozygous individuals in group A} = \frac{p^2}{p^2 + 2pr} \quad (13)$$

Similarly

$$\text{the proportion of homozygous individuals in group B} = \frac{q^2}{q^2 + 2qr} \quad (14)$$

When making these calculations it is preferable to use the improved estimates of the gene frequencies given by formulae (10) to (12)

It is frequently possible to distinguish homozygous and heterozygous individuals by studying the blood groups of parents and children. For example, if a group A individual has a group O parent or child he must of necessity be of genotype $I^A I^O$. When neither of the parents and none of the children are group O this leaves open the question of zygosity except in certain special cases such as when both parents are group AB.

Sometimes it is possible to distinguish heterozygous from homozygous individuals by serological means since individuals homozygous for a particular blood factor generally have red blood cells which are clumped more strongly by the corresponding antisera. This is known as the gene dose effect. The gene dose effect however is not sufficiently pronounced to be dependable in the case of the A B O groups and in addition the existence of the subgroups of group A interferes with the interpretation of the reactions. The possibility that an anti O reagent might be prepared which could detect the presence of a product of gene I^O has frequently been discussed in the literature. Such antibodies have indeed been found in the serum of certain sensitized group AB persons³⁷⁻³⁸ but they are extremely rare and those encountered to date have been of too low titer and avidity to yield reliable results.

If the gene frequencies p , q and r for the general population are known the percentages of children of the various types for every possible mating can be calculated. Schiff³⁹ has derived general formulae

lae for expressing these frequencies in terms of p , q and r , as shown in table 3. From the qualitative point of view, it is evident that no child will have blood containing the agglutinin A or B unless that agglutinin is present in the blood of one or both parents. In addition a parent of group AB cannot have a group O child, and a group O parent cannot have a child of group AB. Studies on thousands of families by many independent workers have yielded data supporting the expectations under the theory of multiple alleles, when allowance is made for technical errors and a certain incidence of illegitimacy.

The study of mother-child combinations can also yield useful data for testing the genetic theory. There are 16 different theoretically possible mother-child combinations, but of these two are excluded under the theory of multiple alleles, namely, mother AB and child O, and group O mother with an AB child. In such studies the problem of illegitimacy does not enter, and it is significant that the results show no deviation from the genetic theory. Formulae for the 14 possible mother-child combinations in terms of p , q and r are readily derived as shown in table 4.

Subgroups of A

Gene Frequency Analysis. As already mentioned the two most common subgroups of group A and group AB are readily included in the scheme by postulating two allelic genes, I^{A_1} and I^{A_2} , in place of I^A , where I^{A_1} is dominant to I^{A_2} . Corresponding to the 6 possible phenotypes there are 10 possible genotypes as follows: group O genotype $I^O I^O$, subgroup A_1 genotypes $I^{A_1} I^{A_1}$, $I^{A_1} I^{A_2}$, and $I^{A_1} I^O$, subgroup A_2 genotypes $I^{A_2} I^{A_2}$ and $I^{A_2} I^O$, group B genotypes $I^B I^B$ and $I^B I^O$, subgroup $A_1 B$ genotype $I^{A_1} I^B$, and subgroup $A_2 B$ genotype $I^{A_2} I^B$. If the frequencies of the 4 alleles I^{A_1} , I^{A_2} , I^B , and I^O are represented by the symbols p_1 , p_2 , q and r , respectively, then the frequencies of the genes can be calculated from the frequencies of the 6 phenotypes in the population, with the aid of the following formulae, which are derived like the formulae for p , q and r of equations (1) to (3)

$$p_1 = \sqrt{\bar{O} + \bar{A}_1 + \bar{A}_2} - \sqrt{\bar{O} + \bar{A}_2} \quad (15)$$

$$p_2 = \sqrt{\bar{O} + \bar{A}_2} - \sqrt{\bar{O}} \quad (16)$$

$$q = \sqrt{\bar{O} + \bar{B}} - \sqrt{\bar{O}} \quad (17)$$

$$r = \sqrt{\bar{O}} \quad (18)$$

TABLE 3
FREQUENCIES OF TYPES OF MATINGS AND THEIR OFFSPRING

Type of Mating	Frequency of Mating	Offspring			
		O	A	B	AB
O × O	r^4	r^4	none	none	none
O × A	$2pr^2(p+2r)$	$2pr^3$	$2pr^2(p+r)$	none	none
O × B	$2qr^2(q+2r)$	$2qr^3$	none	$2qr^2(q+r)$	none
O × AB	$4pqr^2$	none	$2pqr^2$	$2pqr^2$	none
A × A	$p^4(p+2r)^2$	p^3r^2	$p^3(p+r)(p+3r)$	none	none
A × B	$2pq(p+2r)(q+2r)$	$2pqr^3$	$2pqr(p+r)$	$2pqr(q+r)$	$2pq(p+r)(q+r)$
B × B	$q^4(q+2r)^2$	q^3r^2	none	$q^2(q+r)(q+3r)$	none
A × AB	$4p^2q(p+2r)$	none	$2p^2q(p+2r)$	$2p^2qr$	$2p^2q(p+r)$
B × AB	$4pq^2(q+2r)$	none	$2pq^2r$	$2pq^2(q+2r)$	$2pq^2(q+r)$
AB × AB	$4p^2q^2$	none	p^2q^2	p^2q^2	$2p^2q^2$
Totals	1 00	r^2	p^3+2pr	q^2+2qr	$2pq$

Modified after Schiff *Technik der Blutgruppenuntersuchung*, p 68, Julius Springer, Berlin (1932)

TABLE 4
FREQUENCIES OF MOTHER CHILD COMBINATIONS IN TERMS OF p q AND r
(After Schiff)*

Group of Mother	Children's Groups			
	O	A	B	AB
O†	r^2	pr^2	qr^2	0
A	pr^2	$p(p^2 + 3pr + r^2)$	pqr	$pq(p + r)$
B	qr^2	pqr	$q(q^2 + 3qr + r^2)$	$pq(q + r)$
AB	0	$pq(p + r)$	$pq(q + r)$	$pq(p + q)$

* Schiff *Ann. Hoch.* 5 303 (1927)

† As can be seen, if families with group O mothers are considered separately, the percentages of children of groups O, A, and B are equal to the gene frequencies r , p , and q respectively.

The values for p , q , and r , obtained by formulae (15) to (18) while consistent, are not efficient, and generally their sum is not exactly equal to 100 per cent. A corrected set of gene frequencies which closely approximates the maximum likelihood estimates, can readily be obtained by starting with the values of p' , q' , and r' given by formulae (10) to (12), and then taking

$$p_1' = \frac{p' p_1}{p_1 + p} \quad (19)$$

$$p_2' = \frac{p' p_2}{p_1 + p} \quad (20)$$

Serology. As has already been mentioned, the agglutinin A occurs in two principal forms, designated as A_1 and A_2 respectively. Anti-A serum, which is generally obtained from group B individuals, contains a mixture of antibodies of closely related specificities. If such a serum is absorbed with red cells of subgroup A_2 , a reagent can be obtained which clumps A_1 cells but not A_2 cells. This reagent, known as anti- A_1 serum, can be used to subdivide group A individuals into subgroups A_1 and A_2 , and group AB individuals into subgroups A_1B and A_2B respectively.

There is evidence that the difference between A_1 and A_2 is qualitative, yet the serological reactions disclose certain striking quantitative effects. Absorption of anti- A serum (from group B individuals), for example, by a small amount of A_1 cells has almost the same effect

is absorption by a larger amount of A_2 cells. In fact when the anti A_1 reagent is prepared if an excess of A_2 cells is used for the absorption the reactivity for A_1 cells may be destroyed. Thus the difference between agglutinogens A_1 and A_2 is less sharp than between agglutinogens A and B. Another difficulty is that the presence of agglutinin B reduces the activity of the agglutinin A and vice versa if both agglutinogens are present in the same red cell. This is particularly noticeable in the case of blood of subgroup A_2B where the reactivity of the A agglutinin is so diminished that its presence may be overlooked unless potent anti A reagents are used. Accordingly, a common error is to classify individuals of subgroup A_2B incorrectly as group B.

Another difficulty in technique is created by the fact that in newborn babies the red cells are less agglutinable by anti A and anti B serum and as a result red cells from newborn infants of subgroup A_1 may fail to react with anti A_1 serum and be classified incorrectly as subgroup A_2 . As may be anticipated the difficulty in determining the subgroups is considerably increased when the baby belongs to group AB. Not infrequently therefore it is necessary to wait until the infant is 6 to 12 months old before the subgroups can be reliably determined. Still another difficulty results from the existence of blood cells which give reactions intermediate between A_1 and A_2 ; this intermediate subgroup occurs most frequently among Negroes. While anti A_1 lectins (reagents prepared from seeds) may give more striking reactions than anti A_1 reagents prepared from human group B serum, their use does not obviate the difficulties encountered in subgrouping group A and group AB blood especially of newborn infants and Negroes.

A useful reagent to have when carrying out subgrouping tests is the so-called anti O (A_2) or anti II serum. This reagent reacts regularly with red cells of group O and of subgroup A_2 and more weakly with cells of the other groups. Jandsterner and Levine⁶⁰ first suggested the use of such reagents to confirm the results of the subgrouping tests.

Family Studies. The theory of inheritance of the subgroups of A by means of the corresponding allelic genes I^{A_1} and I^{A_2} has been verified by studies in families by many independent investigators. The studies carried out by Wiener and his collaborators between the years 1930 and 1941 are summarized in table 5. According to the theory the following rules of heredity must hold

TABLE 5

SUMMARY OF PUBLISHED STUDIES ON THE HEREDITY OF THE SUBGROUPS OF GROUP A AND GROUP B

(After Wiener, *Blood Groups and Transfusion* 3rd edition 1943)

Parental Combinations	Number of Families	Number of Children in each Group						Totals
		O	A ₁	A ₂	B	A B	A ₂ B	
A ₂ × O	105	121	(4)	163	0	0	0	291
A ₂ × A ₂	13	13	(1)	26	0	0	0	40
A ₂ × B	43	21	0	34	29	0	32	116
A ₁ B × O	16	0	0	37	27	0	0	64
A ₂ B × A ₂	3	0	0	3	3	0	1	7
A ₂ B × B	10	(1)	0	10	15	0	6	32
A ₁ × O	387	404	605	69	0	(1)	0	1169
A ₁ × A ₁	168	79	391	16	0	0	(1)	490
A ₁ × A ₂	79	52	127	51	0	0	0	233
A ₁ × B	120	57	90	8	51	84	6	299
A ₁ × A ₁ B	16	0	14	9	10	8	2	43
A B × O	38	0	55	0	53	0	0	108
A ₁ B × A ₁	35	0	65	0	19	40	5	129
A ₁ B × A ₂	8	0	18	0	5	(1)	10	34
A ₁ B × B	25	0	15	0	32	16	0	63
A ₁ B × A ₁ B	2	0	6	0	2	8	0	16
Totals	1,068	751	1,481	429	249	158	63	3,134

Note The rare parental combination A₁B × A B and A₂B × A B were not encountered in these studies

1) Agglutinin A₁ cannot occur in a child's blood unless it is present in one or both parents

2) The combination of an A₁B parent with an A₁ child, or an A₂ parent with an A₁B child cannot occur

3) In the mating A₁ × O, if there are any children of group O there cannot be any children of group A₂, and, conversely, if there are any children of subgroup A₂ there cannot be any children of group O (There are other matings in which similar rules can be enunciated, depending upon the fact that an individual of subgroup A₁ must belong to one of the three genotypes, I^{A1}I^{A1}, I^{A1}I^{A2}, or I^{A1}I^O)

4) In the matings A₁B × A₁B and A₁B × B, children of subgroup A₂B cannot occur

Among the 3,134 children of table 5, there were a number of ap

parent exceptions to the genetic theory. These which are indicated by parentheses are present in a larger number than may reasonably be attributed to illegitimacy alone. Probably an important cause for these discrepancies is errors in classifying individuals due to the difficulties of subgrouping already pointed out.

Subgroups A_3 and A_3B The subgroup A_3 is a rare variant of group A which is characterized by the very weak agglutinability of the cells even with potent anti A serum.⁴¹ Such blood is readily recognized when tested with anti A serum because the clumping though distinct is incomplete and the microscopic pattern is that of clumps of red cells on a background of many free cells.⁴² Because of the low frequency of this subgroup the number of heredity studies have been limited but the data obtained support the theory⁴³ that the agglutinogen A_3 is determined by a corresponding gene I^{A_3} recessive to I^{A_1} and I^{A_2} but dominant over I^O .

Other Subgroups Even rarer than bloods of subgroup A_3 are those variants of agglutinogen A which are so weakly reactive that clumping of the red cells occurs only with anti A serums of extremely high titer and avidity whether these serums are derived from group B and group O human beings or produced by immunizing rabbits or other animals. Such bloods have been assigned the designations subgroup A_4 , A_5 , etc. and should not be confused with blood of group A_0 described on page 27. One may anticipate that blood of the subgroups A_3B or A_4B will in most cases be mistaken for group B blood while in newborn babies blood of subgroups A_3 and A_4 will be practically indistinguishable from group O blood. Tests for the blood group substances in saliva (cf. page 30) will be of little or no help in resolving such problems because of the corresponding weak inhibition reactions which will be obtained. Fortunately, in view of the extreme rarity of the agglutinogens A_3 and A_4 this problem hardly ever arises in clinical work, and has not yet arisen in any medical case.

Factor C

When group O individuals are injected with group A red cells, not infrequently the titer of the serum is increased not only for group A cells but also for group B cells. Similarly, injection of group B red cells into group O individuals may raise the serum's titer for group A as well as for group B cells. Conversely, if group O serum is absorbed

with group A cells its reactivity for group B cells may be markedly diminished. Similarly, when group O serum is absorbed with group B cells, a diminution in the reactivity against group A cells may occur. This indicates the presence in group O serum of an antibody which reacts with both group A and group B cells. This antibody has been designated as anti C and the corresponding blood factor, which is common to the two agglutinogens A and B, is known as blood factor C.⁴⁴ Thus, in place of the simple scheme of the four blood groups the more complex one given in table 6 is necessary to account for these observations. It may be remarked that reagents of specificity anti C can also be obtained from animal serums and from seed extracts.

It is clear from table 6 that the factor C is another serological attribute of the agglutinogens A and B. Tests for factor C are not needed to identify the agglutinogens A and B, and need not be taken into account in heredity studies or discussions of genetic theory. The existence of factor C is mentioned here because of its theoretical significance in relation to the difference between a blood factor and an agglutinin, and because of its practical importance in the pathogenesis of ABO hemolytic disease.

A subgroup of group A has been described⁴⁵ which has been variously designated as A_x or A₀. The symbol A₀ was selected by Levine⁴⁶

TABLE 6
SEROLOGIC REACTIONS OF THE FOUR A B O BLOOD GROUPS

Group	Red Cells		Agglutinins in Serum†
	Agglutinogens	Blood Factors	
O	—*	—*	Anti A anti B and anti C
A	A	A and C	Anti B
B	B	B and C	Anti A
AB	A and B	A B and C	—

* For simplicity, agglutinin O and blood factor H are omitted from this table.

† The antibodies of each of the designated specificities are not homogeneous but consist of a spectrum of antibodies. Thus in addition to anti A which acts almost equally on agglutinogens A₁ and A₂ group O and group B serums contain anti A₁ which acts much more strongly on A₁ than on A₂. Similarly in addition to anti C group O serum may contain anti C_A reacting more strongly with agglutinin A than agglutinin B and anti C_B reacting more strongly with agglutinin B than agglutinin A.⁴⁴ Similarly, anti B actually comprises a spectrum of antibodies anti B anti B etc.

because while such cells generally give little or no reaction with potent anti A serum they are clumped by the great majority of group O sera. A more appropriate symbol for this blood group may be C because the most prominent feature of the red cells is the presence of blood factor C. Moreover the sera of such individuals usually agglutinate group A as well as group B cells but the reactions with group A cells are generally of low titer.⁴⁷ While there is some serological evidence that the red cells may contain a very weakly reacting A but no B factor, the almost regular reactions with group O serum suggest that the most important attribute of these cells may be the presence of the C factor.⁴⁸ The report that group A saliva inhibits the agglutination of Λ_0 cells by group O serum but not group B saliva suggests that the factor in such cells should perhaps be designated C_A . Heredity studies have been hampered by the very low frequency of this blood type. The irregular expression of this blood type Λ_0 in the few families studied suggests the operation of a modifier rather than a new allelic gene.

Serological Complexity of the A-B-O Agglutinogens

Immunization of rabbits with human group A blood may stimulate the production of antibodies which react not only with group A cells but which also lyse sheep cells. Similarly the injection of sheep red cells into rabbits may stimulate the production of high titer anti A agglutinins. This demonstrates the existence of a blood factor common to sheep red cells and human group A blood.⁴⁹ Because of the similarity of behavior to *Forsman* antiserum which also hemolyze sheep cells the factor has been designated Γ_A . When human anti A serum is absorbed with sheep red cells in order to remove the I_A antibody a fraction of antibody often remains which agglutinates human group A cells but does not react with sheep cells. This shows that there exist at least two blood factors peculiar to group A cells which are not shared with sheep red cells.

Similarly the anti B agglutinins of human group A serums are not homogeneous as can be shown by their cross reactions with blood cells from lower animals. Thus almost all human anti B serums strongly agglutinate blood from rabbits. Absorption of such serums by rabbit red cells generally leaves behind a fraction which clumps human B cells but not rabbit cells though some human anti B serums exist which can be completely absorbed by rabbit cells as well as by

TABLE 7

SOME OF THE SEROLOGICAL PROPERTIES OF THE A B O AGGLUTINOGENS

Genes	Corresponding Agglutinogens	Blood Factors
I^O	O	H
I^{A_1}	A	A A F _A C
I^{A_2}	A ₂	A H F _A C
I^B	B	B B B C

hum in group B cells.⁵⁰ If we designate the human anti B agglutinins which react exclusively with human B cells as anti B and the agglutinins which react with both rabbit and human B cells as anti B then it is clear that human group B cells have at least two factors B and B while rabbit red cells have factor B but not B. Similarly comparative studies carried out with blood of other mammal⁵¹ such as opossums and guinea pigs disclose additional complexities so that in general human group A serum appears to contain a spectrum of anti B agglutinins while agglutino-gen B has corresponding multiple blood factors B B B etc.

The agglutinogens A and B determined by the corresponding genes I^A and I^B regularly possess all of the appropriate blood factors as described above, which confirms the concept that the factors are merely serological attributes of their single respective agglutino-gen (cf. table 7).

Group-Specific Substances in Body Fluids and Secretions

Chemical studies have disclosed that the A B O blood group substances occur in the body in two forms, one soluble in organic solvents and the other as water soluble mucopolysaccharides. The former are found in the red cells of all individuals of the appropriate blood group and also in the body cell. The water soluble form does not occur in all individuals. Among Caucasoids approximately 75 per cent of all individuals (secreters) have the water soluble form in their bodies whereas 25 per cent (non-secreters) lack it or have only insignificant amounts.⁵²⁻⁵⁴

The water soluble group specific mucopolysaccharides are heat stable so that they can be extracted from tissues with boiling water. They occur in highest concentration in secretory organs such as the

salivary glands, gastric and duodenal mucosa, the pancreas and the seminal vesicles. In the living subject the secretor/non secretor status can be diagnosed most simply by testing the saliva where the group specific polysaccharides occur in high concentration in the case of secretors.

Technique The saliva specimens are collected in wide mouth dry tubes. Only small amounts are necessary for the test. Immediately following the collection the tubes are placed in boiling water for ten minutes to destroy the enzymes which saliva contains, since these may destroy the blood group substances. Coagulated mucus is removed by centrifugation and the supernatant opalescent fluid is used for the test. In infants and small children the saliva may be collected on cotton swabs which are expressed into narrow tubes.⁵⁴

The principle of the test is to determine the ability of the saliva to inhibit group specific agglutination reactions. Thus group A secretor saliva prevents anti A serum from clumping group A cells, but does not prevent anti B serum from clumping B cells and similarly for group B and group AB saliva from secretors (cf table 8). On the other hand group O saliva and non secretor saliva will not inhibit anti A and anti B serums.

Schiff has shown that saliva from secretors of all groups inhibit the agglutination of group O cells by so called anti O serums obtained from animals. Such reagents which are now commonly designated as anti H are more conveniently obtained from seed extracts such as of *Ulex europaeus* and these reagents are very useful for determining the secretor status of group O individuals as well as individuals of the other three blood groups (cf table 8). By family studies Schiff

TABLE 8
INHIBITION TESTS FOR THE PRESENCE OF A B O GROUPS IN SALIVA*

Saliva of	Anti A Serum + Group A Cells + Saliva	Anti B Serum + Group B Cells + Saliva	Anti H Serum + Group O Cells + Saliva
Secretor group O	+	+	-
Secretor group A	-	+	-
Secretor group B	+	-	-
Secretor group AB	-	-	-
Non secretor all groups	+	+	+

* + = agglutination - = no agglutination

and Sasaki⁴⁴ were able to show that the secretor status of an individual is genetically determined, and depends on a pair of allelic genes *Se* and *se*. Thus secretors may be homozygous genotype *SeSe* or heterozygous, genotype *Se se*, while non secretors are always homozygous, genotype *se se*. In family studies the genes *Se* and *se* segregate independently of the genes *I^A*, *I^B*, and *I^O*, so that they may be assumed to be located on a different pair of chromosomes. The secretor type thus provided the first example in blood group genetics of gene interaction.

Aside from the inhibition technique the blood group substances can be demonstrated in the saliva by precipitation techniques. Thus if a potent anti A serum prepared by immunizing rabbits is layered with group A or group AB secretor saliva a white ring will appear at the interface but no such ring appears with non secretor saliva or saliva of group O or group B. This technique is not suitable for routine use because of the difficulty of preparing the reagents. Moreover so far attempts to produce comparable anti B reagents have been unsuccessful. Of interest in this connection is the observation of Ujeyama⁴⁵ who found that about 16 per cent of normal fowl serums contain precipitins for non secretor saliva but not for secretor saliva. Ujeyama designated the antibody in question as anti T and asserted that it detected a substance T, present in saliva of non secretors but not in secretors. This finding conforms with the observation that saliva from non secretors is not devoid of mucopolysaccharides and suggests that the non secretor status is due to some molecular alteration in the blood group mucopolysaccharide which eliminates A B O activity and substitutes T reactivity. Under this concept H and T may be considered to be contrasting products of the genes *Se* and *se* respectively.

Lewis Blood Groups

In 1946 Mourant⁴⁷ encountered a serum in a blood donor, which agglutinated the blood cells of 21 of 96 random Englishmen of group O. The reactions of this serum was stronger at low temperature than at body temperature the antibodies having properties of type specific cold agglutinins as described by Landsteiner and Levine. The blood factor detected by this serum was named Lewis after one of the donors whose serum contained the antibody. In 1948, Andresen⁴⁸ reported that in the course of a year he and Freiesleben had en-

countered 8 examples of an anti-serum which agglutinated the red cells of 21 per cent of adult Danes. The blood factor detected by the sera Andre en designated as L which is fortunate since it proved to be identical with the Lewis factor of Mourant. Shortly thereafter Andre en⁵⁹ reported the existence of a second type of serum which gave reactions almost antithetical to his original anti L serum and which he now called anti L₁ to distinguish it from the second serum which he called anti L. In this review however the symbols Le^a and Le^b will be used for the two Lewis factors in conformity with the recommendation of British workers.

The facts concerning the Lewis blood factors are not entirely clear. Perhaps the main reason for this is that up to now it has been impossible to prepare reagents of satisfactory avidity and titer. The reagents in general have the properties as mentioned above of cold agglutinins of low titer and generally have associated with them non specific cold agglutinins that are difficult to remove even by absorption. Nevertheless certain interesting facts have emerged.

In 1948 Grubb⁶⁰ reported that practically all persons who have the Le^a factor on their red cells are non secretors of the A B O blood group substances while those lacking the Le^a factor almost always are secretors of the A B O blood substances. Tests on the saliva of Le^a positive individuals using the inhibition technique have shown that the saliva can neutralize anti Le^a sera suggesting that the Lewis substance takes the place of the A B O substance in appropriate instances. In this connection it is interesting to note that chemical studies by Morgan et al.⁶¹ prove that the Lewis substance is a mucopolysaccharide chemically indistinguishable by present methods from the A B O substances.

Family studies on heredity of the Le^a factor show that the transmission of Le^a parallels that of the secretor/non secretor trait. Thus two Le(a+) parents who would both be non secretors of the A B O blood group substances will have children all of whom are Le(a+) and non secretors of the A B O substances. This expectation has been fulfilled by family studies carried out by Race et al.⁶² and by Andre en et al.,⁶³ and others who in 17 such families found 44 children all of whom were Le(a+). On the other hand when both parents are Le(a-) children who are Le(a+) as well as Le(a-) occur in a ratio consistent with the hypothesis that factor Le^a, in contrast to other blood factors behaves in inheritance like a Mendelian recessive.

Studies on the heredity of the Le^a factor are complicated by the variable reactivity of the red cells at different ages. Andre⁶¹ early observed that as many as 80 per cent of infants less than 3 months old had $Le(a+)$ red cells. Among older babies the percentage of positive reactions was lower but not until the end of the first year of life did the frequency drop to the adult 20 per cent level. On the other hand Unger found that cord blood specimens are all $Le(a-)$ and according to Rosenfield and Ohno the baby's cells become agglutinable only after the first few weeks of life. Grubb and Morgan observed moreover, that the more $Le(a+)$ cells are washed with saline solution the less agglutinable by anti Le^a serum do they become. However Race and Sanger were unable to wash the Le^a antigen off the red cells and they suggest that such observations may depend on the antiserum used.

Sneath and Sneath⁶⁴ reported that when $Le(a+)$ persons were transfused with $Le(a-)$ blood, the donor's red cells could not be demonstrated in the recipient's circulation by differential agglutination tests employing anti Le^a serum although when antisera which detected other agglutinogens were used the donor's cells could readily be demonstrated. This discrepancy in the results of the differential agglutination tests could be explained by postulating that the donor's $Le(a-)$ cells adsorbed the Le^a substance present in the serum of the recipient and thus became $Le(a+)$. This suggests that in contrast to other blood group substances the Lewis substance is primarily produced in the tissues and body fluid, and secondarily adsorbed onto the red cells. Additional evidence supporting this idea was obtained by studies on a rare instance of mosaicism in a pair of fraternal twins.⁶⁵ The blood vessels of the c twins contained a mixture of two kinds of red blood cells having the genetic blood groups of the two twins respectively. The red cells of the twins reacted in differential agglutination tests like blood obtained from individuals who have had blood transfusions namely like a mixture of two types of blood. However, the twins failed to exhibit mosaicism of their Lewis blood systems. The c twins had the same Lewis substance and had all $Le(a-)$ blood group

systems

When red cells are tested with both sera anti Le^a and anti Le^b , conflicting results are obtained. One cause of difficulty is that anti Le^b

serums give many more positive reactions with cells of group O and group A₂ than with cells of group A₁ and B. When tests are confined to blood cells of group O and A₂, Le^a and I e¹ prove to be almost but not completely antithetical. For example, among 238 group O individuals, Andresen found 46 Le(a+b-), 178 Ie(a-b+), 14 Ie(a-b-) and no Le(a+b+). Other investigators have confirmed these findings, and although a number of genetic theories have been proposed, none satisfactorily explains these findings.

In conclusion, the following hypothesis is offered in an attempt to account for some of the puzzling observations regarding the relationship between the Lewis blood types and the A B O blood group substances and secretor types. It is suggested firstly that the substance I e^a is identical with the substance T found in non secretor saliva by Uyejima, and that substance Le^b is the same as or related to the substance H of secretor saliva,* while both substances are believed to be modifications of a single basic mucopolysaccharide. If this is so, then individuals who are genetically non secretors (genotype *se se*) would be expected to have the substance T (or I e^a) in their secretions and body fluids while secretors (genotype *Se Se* and *Se se*) will have substance H (or I e^b) although heterozygotes may have small amounts of I e^a (or T) in addition. The group substances of secretors, in addition to factor H will have factors A or B or both or neither, depending on the A B O blood group. The genetically determined group specific substance (mucopolysaccharide) present in the body fluid is absorbed onto the red cells which thus may acquire the I e^a (or T), or Le^b (or H) factor according to the secretor type in addition to the group properties originally present. According to this hypothesis one might anticipate that the red cells of group O individuals who are secretors should react more strongly with anti H serums than the red cells of non secretors. Indeed, some preliminary tests using extracts of *Ulex europaeus* as anti H reagents appear to support this expectation.

Theoretico-Statistical Considerations. As has already been pointed out, the presence of Le^a factor on the red cells is demonstrable only in homozygous individuals, genotype *Le^aLe^a*, bearing in mind

* According to W. T. J. Morgan (personal communication) however some preparations of H substance have Le^b activity, while other preparations do not.

TABLE 9
THE LEWIS TYPES*

Red Cell Phenotypes		Corresponding Genotypes		Four Salivary Types	
Designation	Observed frequencies (per cent)	Designations†	Theoretical frequencies	Designation	Estimated frequencies (per cent)
Le(a-b-)	6.01	$lele$ $Le^a le$ $Le^a Le^a$	c^2	I_o I_a I_b	0.4
Ie(a+b-)	22.38		$2ac$		23.0
Ie(i-b+)	71.61		a^2		27.4
		$Le^b le$ $Le^b Le^b$ $Le^a Le^b$	$2bc$	I_b L_{ab}	41.2
			b^2		
			$2ab$		

*Modified after Wiener [*Lab. Digest* (St. Louis) Vol. 18 No. 5 Nov. 1954]

† Gene Le^b is believed to be identical with gene Se ; similarly, genes Le^a and le may both correspond to gene se .

that gene Le^a and gene se for the ABH non secretor trait may be identical.⁶⁸ This may represent a gene dosage effect since all the available anti Le^a serums have been of very low titer and avidity. (Similarly weak anti N serums fail to agglutinate type MN cells while clumping type N cells distinctly.) In secretors of the ABH substance the group substances have proved to be present in saliva, gastric juice and other secretions in considerably higher concentrations than on the red cells. If the same is true of the Le^a substance, then positive reactions by the inhibition test on saliva would be expected in heterozygous as well as homozygous Le^a positive individuals. Thus bearing in mind that the Le^b and H factors are closely related and that genes Le^b and Se may be identical the distribution of the Lewis types as determined by tests on saliva would be expected to be different from that obtained by testing red cells.

Let I_o , I_a , I_b and L_{ab} represent the four Lewis types as determined by testing saliva, where I_o represents absence of both substances, I_a represents presence of Le^a and absence of Le^b , etc. Also let a , b and c represent the frequencies of genes Le^a , Le^b , and le , respectively, where Ic^2 determines the presence of Ic^a substance, Ic^b determines the presence of substance Le^b , and le determines neither. Then table 9 can be drawn up.⁶⁷

Table 9 also gives the distribution of the three Lewis red cell phenotypes in England as reported by Race and Singer⁶⁶ in their

book. From the frequencies of these three phenotypes the frequencies of the three Lewis genes can be estimated as follows:

$$a = \sqrt{\text{Le}(a+ b-)} = \sqrt{0.2238} = 47.3 \text{ per cent}$$

$$c = \sqrt{\text{Le}(a- b-) + \text{Le}(a+ b-)} = \sqrt{\text{Le}(a+ b-)} = 6.0 \text{ per cent,}$$

$$\text{And } b = 1 - a - c = 46.7 \text{ per cent}$$

From the estimated gene frequencies the expected distribution of the four salivary Lewis types can be calculated as follows:

$$L_0 = c^2 = (0.06)^2 = 0.36 \text{ per cent}$$

$$L_a = a^2 + 2ac = (0.473)^2 + 2(0.473)(0.06) = 28.05 \text{ per cent}$$

$$L_b = b^2 + 2bc = (0.467)^2 + 2(0.467)(0.06) = 27.4 \text{ per cent}$$

$$L_{ab} = 2ab = 2(0.473)(0.467) = 44.2 \text{ per cent}$$

Unfortunately very little data are available with which to test these calculations. Grubb and Morgan⁶⁶ have reported observations on the correlation between the Le^a phenotype of the red cells and the ABH secretory type in a series of 122 individuals as follows:

1 erythrocytes $\text{Le}(a+)$ ABH non secretor	5
1 erythrocytes $\text{Le}(a-)$ ABH secretor	163
1 erythrocyte $\text{Le}(a+)$ ABH secretor	0
1 erythrocytes $\text{Le}(a-)$ ABH non secretor	2

If we assume that the two non secretors who were of red cell phenotype $\text{Le}(a-)$ had no Lewis substance in their saliva this would make the frequency of the L_0 type only 2 among 222 individuals or 0.9 per cent which does not deviate significantly from the expected frequency of 0.4 per cent.

Quite different results have been reported by Ceppellini⁶⁷ who tested the saliva of 518 individuals for the ABH and Le^a substances. He designated ABH secretors as S and non-secretors as ns, similarly Le^a secretors were designated as I and non secretors as ni. Ceppellini⁶⁷ thus identified 4 salivary types with the following distribution among 518 individuals:

ns I	11 or 2.13 per cent
ns I	83 or 16.02 per cent
S ni	50 or 9.25 per cent
S I	374 or 72.25 per cent

When these findings were arranged in a 2×2 contingency table Ceppellini found $\chi^2 = 0.036$, which in his opinion strongly indicated that the secretions of the Le^a and ABH substances were genetically independent of one another. If, as seems likely, the four types $ns\ nl$, $ns\ L$, $S\ nl$, and $S\ L$, correspond to the salivary types L_a , L_b , L_{ab} , and L_{ab} , respectively, this would mean that the heredity of Le^a and Le^b depends on independent pairs of genes rather than corresponding allelic genes. Thus, the conflict between Ceppellini's findings and views and the hypothesis favored by the authors is reminiscent of the polemic between Hirsfeld and Bernstein regarding the heredity of the four A B O groups. A serious objection to Ceppellini's hypothesis* is that it does not provide a satisfactory explanation for the reciprocal relationship between the Ie^a blood type and the ABH secretor type*. On the other hand, since the frequency 72.25 per cent which he observed for type $S\ L$ or L_{ab} significantly exceeds 50 per cent, this would argue against the concept of Ie^a and Ie^b as allelic genes. However, Ceppellini's observations conflict with those of Grubb and Morgan and the explanation possibly lies in the difficulty of testing for Le^a substance in saliva. As already pointed out, the anti Le^a reagents available are of low titer and avidity, so that inhibition tests with such reagents do not give clear cut results. In fact, examination of sample protocols given by Ceppellini shows no sharp division between types L and nl so that not infrequently the diagnosis seems arbitrary. Thus, the introduction of a slight bias when reading the reactions would strongly influence the results obtained. Obviously, until better reagents become available, judgment must be suspended.

A third hypothesis has been suggested by Grubb and Morgan. They suggest that there are three pairs of allodromorphic genes, Sc and sc , Le^a and Ie^a , Ie^b and Ie^d , occupying contiguous loci. This hypothesis differs from the one proposed here in the same way as Fisher's theory of triply linked genes differs from Wiener's theory of multiple alleles for the Rh Hr blood types. In the absence of evidence of crossing over the theory of separate gene loci degenerates into a theory of complete

linkage, and since completely linked genes operate as a unit, the difference from the theory of multiple alleles becomes solely a matter of semantics

A-B-O Blood Groups in Apes and Monkeys

In 1925, Landsteiner and Müller⁶⁹ showed that blood of chimpanzees, orangutans and gibbons have isoagglutinogens and isoagglutinins indistinguishable from those which characterize the human blood groups. To date more than 100 chimpanzees have been tested, of whom approximately 90 per cent proved to belong to group A and the remainder to group O. Group A chimpanzee blood resembles human blood of subgroup A₁ more closely than subgroup A₂.⁷⁰ Many fewer of the other anthropoid apes have been tested, among 18 orangutans, 7 proved to be group A, 8 group B, and 3 group AB. Among 7 gibbons, Landsteiner found 1 group A, 5 group B, and 1 group AB.

Wiener tested the blood of two dead gorillas and found anti A in the serums, but tests on the cells failed to reveal the presence of the expected agglutininogen B, in conflict with Landsteiner's law. Examination of the salivary glands, however, showed them to contain blood group substance B. In studies on various species of monkeys, Wiener, Candela and Goss⁷¹⁻⁷² found that their red cells were not agglutinated by anti-A or anti-B serums, but the serum of individual monkeys differed in that some contained anti A alone, some anti B alone, others showed both anti A and anti B, while the remainder showed neither. Examination of the saliva and salivary glands explained these observations since the expected blood group substances were found to be invariably present. Thus, in gorillas and monkeys Landsteiner's law does hold, but the reciprocal relation is between the isoagglutinins in the serum and the blood group substances in glandular tissue and secretions rather than on the red cells themselves.

Anomalies of the A-B-O Blood Groups

Recently, Wiener and Gordon⁷³ encountered two individuals having anti B and not anti A agglutinins in their serums, but whose red cells failed to react with anti A serum. Examination of the saliva showed the presence of the expected blood group substance A. Thus, these two rare individuals gave blood group reactions resembling those of lower monkeys, and for this reason this anomalous blood type has

been designated A_m . Additional examples of this blood type have been observed by Weiner et al,⁷⁴ who postulated the existence of a pair of allelic genes Y and y independent of the A B O genes, where y is extremely rare. The extremely rare homozygotes of genotype yy are postulated to be capable of producing blood group substances in the secretions but not on the red blood cells. The counterpart of A_m , namely, B , has recently been encountered in a father and 2 of his 4 children by Yokoyama et al.⁷⁵

In 1951, Bhende et al.⁷⁶ described a very rare type of blood which has been called 'Bombay'. The red cells of individuals of this type were not clumped by anti A, anti B, or anti H and the serum contained anti H as well as anti A and anti B. The saliva of these persons was devoid of A B O group specific substance. Levine et al.⁷⁷ have reported a family in which three members had the 'Bombay' blood type. From the blood groups of the parents and children they were led to postulate the existence of a rare modifying gene which in homozygous individuals, genotype xx , prevents the blood group genes from producing A B O specific substances both on the red cells and in the saliva. The red cells of all these individuals are $Ie(1+)$, which is further evidence of the reciprocal relationship between Le and H . Levine has assigned to blood of the Bombay type the symbol, O_b , to indicate that the red cells react like group O cells, but lack the blood factor H .

The recent developments demonstrate that the heredity of the A B O blood groups is quite complex, and the suggestion made by Haldane that the blood group substances are the direct products of the blood group genes does not seem tenable in the light of these observations. Apparently, all or almost all humans are capable of producing a basic blood group mucopolysaccharide, and the capacity to produce this substance presumably depends upon the action of two or more pairs of intermediate genes. The blood group genes I^A , I^B and I^O apparently convey the capacity to produce certain enzymes which act on the basic mucopolysaccharide and give it its blood group specificity. Still another pair of genes Se and se determine the capacity to produce the polysaccharide in glandular tissue as well as in the erythrocytes and body cells, and confer Le^b (or H) and Le^a (or T) specificity on these substances. Individuals of the very rare genotype yy apparently can produce group specific substances in the secretions

but not on the red blood cells, while the mucopolysaccharides of individuals of the very rare genotype xx ("Bombay" type) completely lack ABH group specificity both in secretions and on red cells. The presence of genotype xx , while it prevents the expression of the blood group, does not interfere with the hereditary transmission of the A-B O genes, so that the blood group will show up as expected in other individuals of the family who are not homozygous for gene x .

CHAPTER III

The M-N-S System

M-N Types

The three M-N types provide one of the simplest examples of Mendelian heredity in man

Serology To determine the M-N types of an individual anti-M and anti-N reagents from rabbit immune anti-serums are still used today, as in the original investigations of Landsteiner and Levine^{79, 80} Occasionally, human serums have been encountered containing anti-M and anti-N agglutinins but these are so rare that their use is not practicable for routine blood typing Similarly, adequate quantities of seeds yielding anti-N lectins have not been generally available for this purpose so far while anti-M has not been found in seeds

The anti-M serums produced by immunizing rabbits with human type M blood contain in addition to anti-M also anti-human species-specific agglutinins The serum must therefore be processed by absorbing it with type N cells which removes the species-specific antibodies but leaves behind the anti-M agglutinin Anti-N serum is processed similarly by absorbing it with type M cells While there is little difficulty in preparing potent anti-M reagents in this way it has not been possible to obtain high titer anti-N reagents The reason is that the type M blood cells used to absorb the species antibody can also absorb some of the desired anti-N agglutinin, in fact if an excess of cells is used the reagent may be made devoid of activity or weakened so that only type N, but not type MN cells are agglutinated (Also anti-N agglutinins in human serum can be absorbed by type M cells*) The practical importance of this phenomenon is that there is danger of obtaining false negative reactions if the anti-N reagents have been overabsorbed On the other hand, if the serum has been underabsorbed in an attempt to avoid the removal of the type-specific agglutinin, reagents may result which give false positive reactions This difficulty has been responsible for serious errors in the application of the M-N types to anthropology and legal medicine, when the

* This is another of the numerous examples of the cross reactivity of blood group antibodies as discussed on page 11

tests were undertaken by workers who did not first qualify themselves by becoming experienced in serologic techniques

Anti M and anti N reagents yield only three pairs of reactions with human red cells namely $M+N-$, $M+N+$ and $M-N+$ corresponding to the three types designated M MN and N respectively. Bloods lacking both agglutinogens have not been encountered in tests on many millions of blood specimens

Heredity According to the theory of Landsteiner and Levine²⁰ the agglutinogens M and N are transmitted by two corresponding allelic genes L^M and L^N . This theory allows for only three possible genotypes corresponding to the three known phenotypes and at the same time precludes the existence of type M-N-. There are six matings theoretically possible and the distribution of the types to be expected among the children of each mating is shown in table 10. While most studies conform with the predictions some workers have reported an excess of type MN children in those families where both parents are type MN and this supposed excess of type MN children has been cited as an example of selection in favor of the heterozygote. In table 11 is shown the summary of the investigations carried out by Wiener and his collaborators during the years 1929-52 on 1,800 families with 3,379 children. Among the 800 children from the mating $MN \times MN$ 405 belonged to group MN, a finding consistent with simple random redistribution of the genes L^M and L^N without the need to invoke other mechanisms. The excess of type MN children reported by other workers was therefore probably due to the use of incompletely absorbed antisera, as already explained.

From the point of view of medicolegal application the heredity

TABLE 10
HEREDITY OF THE M-N TYPES

Matings		Children (per cent)		
Phenotypes	Genotypes	M ($L^M L^M$)	MN ($L^M L^N$)	N ($L^N L^N$)
M \times M	$L^M L^M \times L^M L^M$	100	0	0
M \times N	$L^M L^M \times L^N L^N$	0	100	0
N \times N	$L^N L^N \times L^N L^N$	0	0	100
M \times MN	$L^M L^M \times L^M L^N$	50	50	0
N \times MN	$L^N L^N \times L^M L^N$	0	50	50
MN \times MN	$L^M L^N \times L^M L^N$	25	50	25

TABLE 11

HEREDITY STUDIES ON THE M N TYPES (1929-1952)*

(After Wiener A S DiDiego N and Sokol S *Acta Genet Med et Gemellol* 2:391 1953)

Parental Combination	Number of Families		Children Belonging to Type			Total
			M	N	MN	
M × M	153	Number Per cent	326 92.7	0 0	(1)† 0.3	327
M × N	173	Number Per cent	(1)† 0.3	0 0	3.6 10.7	377
N × N	57	Number Per cent	0 0	106 100.0	0 0	106
MN × M	463	Number Per cent	433 51.3 ± 1.1†	(1)† 0.1	4.3 48.6 ± 1.1	973
MN × N	351	Number Per cent	(3)† 0.4	382 47.3 ± 1.2	411 51.7 ± 1.1	796
MN × MN	377	Number Per cent	139 24.7 ± 0.8	196 24.5 ± 1.0	405 50.8 ± 1.1	800
Totals	1580		1028	685	1666	3379

* Since that time many hundred additional families have been studied

† The e apparent exceptions to the laws of inheritance are probably due to illegitimacy

‡ Figure following ± sign represents probable error

of the M N types may be summarized by the following two laws:

- 1) The agglutminogen M or N cannot appear in the blood of a child unless it is present in the blood of one or both parents.
- 2) A type M parent cannot have a type N child, and a type N parent cannot have a type M child.

The few exceptions to these rules in table 10 may reasonably be attributed to illegitimacy, and in support of this interpretation it may be pointed out that while there are families in which the blood types of the supposed fathers conflict with the second law, there are none in which this is true of the mother.

Gene Frequency Analysis. The frequency of the allelic genes L^M and L^N in the general population can be readily estimated from the distribution of the three phenotypes.²¹ Since there is only one pheno-

type corresponding to each genotype the gene frequencies can be determined by direct count. If m represents the frequency of gene I^M and n represents the frequency of gene I^N

$$\text{Then } m = \bar{M} + \frac{\overline{MN}}{2} \quad (21)$$

$$\text{And } n = \bar{N} + \frac{\overline{MN}}{2} \quad (22)$$

If V represents the size of the sample then $2V$ is the number of genes counted

$$\text{Therefore } \sigma_m = \sigma_n = \sqrt{\frac{mn}{2V}} \quad (23)$$

From the gene frequencies the distribution of the three MN types can be recalculated as follows

$$M_0 = m^2 \quad (24)$$

$$N_0 = n^2 \quad (25)$$

$$\overline{MN}_0 = 2mn \quad (26)$$

As a rule the recalculated frequencies will not be identical with the observed frequencies. Assuming random intermarriage which theoretically should bring the distribution of the MN types to equilibrium in a single generation the genetic theory can be tested by calculating the chi square value as follows⁸¹

$$\chi^2 = \frac{(a - a_0)^2}{a_0} + \frac{(b - b_0)^2}{b_0} + \frac{(c - c_0)^2}{c_0}$$

Here a , b and c represent the observed frequencies of the types M , MN and N respectively in the population and a_0 , b_0 and c_0 the theoretically expected values according to formulae (24) to (26). This then reduces to the following formula

$$\chi^2 = \frac{(b^2 - 4ac)^2 V}{(2a + b)^2 (b + 2c)^2} \quad (27)$$

where V is the number of individuals examined. This test has been applied in a large number of population studies and the results lend further support to the genetic theory.

The frequencies of the genes L^M and L^N can also be calculated by the square root method as follows

$$m' = \sqrt{\bar{M}} \quad (28)$$

$$n' = \sqrt{\bar{N}} \quad (29)$$

These formulae for the gene frequencies, though consistent are less efficient than formulae (21) and (22), the latter being identical with the maximum likelihood estimates

The formulae (27) and (28) are the basis for a different test of the genetic theory. Since theoretically, $m' + n' = 1$ then $\sqrt{\bar{M}} + \sqrt{\bar{N}}$ should equal unity or 100 percent. In general, the sum $\sqrt{\bar{M}} + \sqrt{\bar{N}}$ will not be exactly equal to unity, and the size of the deviation can be used as a test of the genetic theory. If the deviation

$$D = 1 - (\sqrt{\bar{M}} + \sqrt{\bar{N}}) \quad (30)$$

Then, as Wiener⁸¹ has shown,

$$\sigma_D = \frac{1}{2\sqrt{V}} \quad (31)$$

where V represents the number of persons tested. Here again the observations confirm the genetic theory.

If we represent the frequency of type MN by H ,

$$\text{Then, } \frac{dH}{dm} = \frac{d}{dm} m(1-m) = 1 - 2m$$

To find the maximum value of H , set $\frac{dH}{dm} = 0$, so that $m = \frac{1}{2}$

Therefore, the maximum frequency that type MN can have under this genetic theory is 50 per cent.

This important result applies to any heterozygote in a population at equilibrium, so that, for example, the frequency of group AB can not exceed 50 per cent. As will be pointed out later, this law was applied by Wiener to elucidate the heredity of the Rh Hr blood types (cf. page 63).

Mother-Child Combinations. As already pointed out, the study of mother-child combinations supports the theory because of the absence of the combinations M mother/A child or A mother/M child. Further evidence supporting the theory has been obtained

TABLE 12
FREQUENCIES OF MOTHER CHILD COMBINATIONS

Mothers of Type	Children of Type		
	M	MN	N
M	11^2	11^2n	0
MN	1^2n	$11n$	$11n^2$
N	0	mn^2	n^3

by gene frequency analysis by applying the formulae of table 12 to test the results of studies on the MN types in mothers and children.

Agglutininogen N. As in the ABO system a number of rare anomalies of the MN types have been encountered. The most important is the so called agglutininogen N_2 first described by Crome⁸² and which according to Friedenreich⁸³ is due to a special corresponding rare allelic gene L^{N_2} . This type of blood is characterized by its very weak reactivity with anti-N serums so that with many anti-N reagents clumping may fail to occur or may be so weak as to be overlooked. This may lead to serious errors, as in medicolegal cases. For example if the putative father belongs to type MN_2 (genotype $L^NL^{N_2}$) and the child to type N (genotype $L^NL^{N_2}$) the former might erroneously be reported as type M with resulting false exclusion of paternity.

Agglutinogens M and N in Monkeys and Apes. In tests on human blood no differences are demonstrable among different anti-M reagents or different anti-N reagents. For example all anti-M serums, regardless of source and method of preparation regularly agglutinate red cells of type M and of type MN but not of type N. However when the cross reactions of these reagents are studied using red cells of anthropoid apes and monkeys it is found that not all anti-M serums give the same reactions nor do all anti-N serums.^{84, 85}

In table 13 for example are shown the differences in behavior of six selected anti-M serums prepared from immune rabbit serums in tests on red cells from anthropoid apes and monkeys. In the table, the anti-serums have been arranged so that those with more cross reactions precede those with fewer cross reactions, and the blood specimens have been arranged according to the position in the zoological scale of the primates from which they were obtained. Although the anti-M serums with the highest titers give the most cross reactions, the

differences were not merely quantitative as was shown by absorption experiments. For example, reagent M1 had a titer of 64 units for human red cells and 16 units for rhesus red cells. Absorption by human cells completely removed all the antibodies while absorption by rhesus cells produced a reagent with a titer of 16 units for human M cells but not reacting with rhesus cells. On the other hand, reagent M5 which likewise had a titer of 64 units for human M cells and about 24

TABLE 13

M FACTORS IN BLOOD OF APES AND MONKEYS

(After Wiener A S *Blood Groups and Transfusion* 3rd ed 1943)

Source of Blood Specimens	Anti M Testing Fluids					
	M5	M1	M21	M35	M2	M92
Human M	+++	+++	++±	++±	++±	++±
Human N	0	0	0	0	0	0
Chimpanzee	+++	+++	+++	++	+++	±
Old World Monkeys (Cercopitheidae)						
Sphinx Baboon	+++	++	++±	0		0
Drill Baboon	+++	+++	++±	(±)	(+)	(+±)
Chacma Baboon	+++	+++	++±	0	tr	0
M rhesus	+++	+++	++±	(+±)	±	0
Java Macaque	+++	+++	+±	0	0	0
Sooty Mangabey	+++	+++	++±	tr	±	±
Green Monkey	+++	+++	0	0	0	0
New World Monkeys (Platyrrhini)						
White Spider Monkey	++±	0	0	0	0	0
Black Spider Monkey	±	0	0	tr		0
Woolly Monkey	0	0	(±)	0	0	0
Brown Ringtail (Capuchin Monkey)	0	0	0		0	0
Moss Monkey	0	0	f tr	±		0
Lemur	0	0	0	0	0	0
Average titer of testing fluids	64	64	32	24	16	16

Of several species two individuals were tested of brown ringtails 4 of *Macacus rhesus* 45

Reactions placed in parentheses were found not to be removed by absorbing the serum with type M blood cells

units for rhesus red cells was completely absorbed by either human M cells or rhesus red cells.

Besides providing a striking example of biochemical evolution table 13 is another example of the mosaic structure of red cell agglutinogens. To explain the reactions it is necessary to recognize that not all anti M agglutinins are identical in specificity and therefore by definition there are more than one kind of M factor. If we designate as anti M_1 the agglutinin which reacts with human red cells but not with blood from chimpanzees or monkeys and as anti M_{11} the agglutinin which detects a factor common to human M blood and chimpanzee blood but absent from the blood of lower primates, etc. it follows that the human M agglutimogen has all of the blood factors $M_1 M_2 M_{11}$, etc. chimpanzee blood factors $M_1 M_{11}$, etc. One could anticipate that the injection of rhesus monkey blood into rabbits would yield a reagent specific for human M cells and not reacting with type N cells. Indeed Landsteiner and Wiener⁸⁴ found it possible to prepare satisfactory anti M reagents in this way and it is of historical interest that these experiments led to the discovery of the Rh (rhesus) factor.

By similar experiments it has been possible to demonstrate that the agglutimogen N has a mosaic structure.⁸⁴ Due to difficulty in obtaining potent reagents, as already explained, only two such factors have been demonstrated so far, namely, N_1 peculiar to human N cells alone and N_2 , a factor shared by human N cells and chimpanzee red cells.

Obviously, when a human being inherits an M or N agglutimogen this carries with it all of the serological factors that characterize the agglutimogen, as shown below:

Gene	Corresponding agglutimogen	Blood factors
I^M	M	M, M_1, M_2, M_{11} , etc.
I^N	N	N, N_1, N_2 , etc.

Factors S, s and U

Factor S. In 1947, Walsh and Montgomery⁸⁶ studied the serum of a patient known to be sensitized to the Rh_0 factor and found the serum to contain an additional blood group antibody which gave reactions different from any previously described. A sample of this serum was studied by Singer and Race⁸⁷ who were able to show that the blood factor detected by it was related to the agglutimogen M.

TABLE 14
DISTRIBUTION OF THE M N S TYPES AMONG 394 BLOOD DONORS

Phenotype		Number	Per Cent
M	S+	88	22.3
	S-	37	9.4
N	S+	27	7.0
	S-	49	12.4
MN	S+	106	26.9
	S-	87	22.0
Total	S+	221	56.2
	S-	173	43.8

and N. This was done by making use of the 2×2 contingency table as already explained on page 3.

When persons of type M, MN, and N were subdivided according to their reactions with the anti serum of Walsh and Montgomery, named anti S by Sanger and Race, 6 types of blood could be distinguished. For example, in 394 tests done on professional donors in New York City, the distribution shown in table 14 was found.⁵⁵

On inspection it can be seen that the incidence of the S factor is not the same in the three M N types, since more than two thirds of type M individuals are S positive while only about one third of type N individuals are S positive. That this difference is not merely accidental but statistically significant can be shown by a 2×2 contingency table as follows:

	M + MN	N	
S+	194	27	221
S-	121	49	173
Totals	318	76	394

Applying the formula on page 3, $\chi^2 = 16$ for one degree of freedom and the likelihood of this occurring by chance alone is much less than 0.001.

To account for the findings, Sanger and Race mention two genetic hypotheses. According to one, the six possible M N-S groups are at

TABLE 15
THE M N S PHENOTYPES AND GENOTYPES

Phenotypes	Reactions with Serums			Corresponding Genotypes	
	Anti M	Anti N	Anti S	Linked genes	Multiple alleles
MS	+	-	+	MS/MS & MS/Ms	L ^S L ^S & I ^S I
Ms	+	-	-	Ms/Ms	LL
NS	-	+	+	NS/NS & NS/\s	I ^S I ^S & I ^S I
\s	-	+	-	\s/\s	ll
MN S	+	+	+	MS/\s Ms/\s	L ^S I ^S I ^S I & I I ^S
MN s	+	+	-	Ms/\s	Ll

tributed to 4 allelic genes *MS*, *Ms*, *NS* and *\s* where *S* is considered to be a mutation in the *M* and *N* genes which renders the red cells agglutinable by anti *S* serum. According to the second hypothesis *S* is considered to be a separate gene with an allele *s* and closely or possibly completely linked with the *M N* genes. The six possible genotypes, their reactions and corresponding genotypes according to each of these hypotheses are shown in table 15. Though Race and Singer prefer the linked gene theory, the present authors regard the concept of multiple alleles to be more consistent with the known genetic and serological data.⁵⁹

When factor *S* is considered alone, two types are distinguishable, namely *S* + and *S* - the former having the frequency of 56.1 per cent and the latter 43.9 per cent in tests done on 394 blood donors in New York City. Assuming heredity by a pair of genes *S* and *s* so that corresponding to phenotype *S* there are two genotypes *SS* and *Ss* and corresponding to phenotype *s* there is only one genotype *ss*, the frequencies of genes *S* and *s* can be estimated as follows:

$$s = \sqrt{0.439} = 0.663 \text{ or } 66.3 \text{ per cent}$$

so that *S* = 0.337 or 33.7 per cent

Therefore, the estimated frequencies of the three genotypes are *SS* = 0.114, *Ss* = 0.447, and *ss* = 0.439.

At this stage of knowledge it could be surmised that there might exist a factor, *s*, corresponding to the gene *s*, and demonstrable in the blood cells of individuals of genotypes *Ss* and *ss*. In fact, Levine et al.⁶⁰ did find a serum in the mother of an erythroblastic infant which

gave reactions corresponding to those expected for factor *s*. Subsequently additional examples of this anti serum were encountered by other workers. However while a fair number of satisfactory anti *S* serums have been found to date serum of the specificity of anti *s* is in very short supply and more or less of a serological curiosity. Therefore the number of studies carried out on factor *s* have been quite small.

Nomenclature The vast majority of studies on the M N system have been confined to the blood factors M and N alone since these have been known for the longest time and the reagents for their recognition can be procured at will by immunizing rabbits. As mentioned previously three types are distinguished with the aid of the anti-serums. When tests for factor *S* are also carried out the number of phenotypes in the system is increased to six. When in addition anti *s* serum is used nine phenotypes can be recognized. Since an increasing number of studies will be made in the future using the various serums at different stages it is desirable to have an unambiguous nomenclature for each of the three stages which at the same time will indicate clearly not only the phenotypes but also the theoretically possible genotypes. The nomenclature used by Wiener² is shown in table 16.

TABLE 16
NOMENCLATURE OF THE M N S s TYPE²
(After Wiener)

Three M N Phenotypes			Six M N S Phenotypes		Nine M N S-s Phenotypes		Corresponding Genotypes
Designation	Reaction with serum		Designation	Reaction with serum	Designation	Reaction with serum	
	Anti M	Anti N					
M	+	-	MS	+	MSS	-	L ^S L ^S
			Ms	-	MSs	+	L ^S L
					Mss	+	LL
N	-	+	NS	+	NSS	-	I ^S I ^S
			Ns	-	NSs	+	I ^S I
					Nss	+	II
MN	+	+	MN S	+	MN SS	-	L ^S I ^S
					MN Ss	+	L ^S I and LI ^S
					MN ss	+	LI

Gene Frequency Analysis.⁹¹⁻⁹² If the distribution of the M N S phenotypes in the general population is known, the hypothetical gene frequencies can be readily calculated

Stage 1 At this stage only tests for factors M and N have been done, and three phenotypes distinguished. In this case, the gene frequencies, m and n can be readily calculated, as previously explained, by direct count

Stage 2 Here tests have been made for factor S as well as for M and N. The following relations hold by identity

$$L^s + L = m \quad (32)$$

$$l^s + l = n \quad (33)$$

where m and n are the frequencies of gene L^M and gene L^N respectively [cf. formulae (21) and (22)]

$$\text{Moreover, } L^s + l^s = S \quad (34)$$

$$\text{and } L + l = s \quad (35)$$

Since, corresponding to the phenotype N s there is only the single genotype ll , then, assuming panmixia, it follows that

$$l = \sqrt{\bar{N}s} \quad (36)$$

where $\bar{N}s$ represents the frequency of type N s in the general population

$$\text{Similarly, } L = \sqrt{\bar{M}s} \quad (37)$$

These preliminary estimates must now be adjusted to make them satisfy the identity $L + l = s$. This can be done by solving the simultaneous equations,

$$L - l = \sqrt{\bar{M}s} - \sqrt{\bar{N}s} \quad (38)$$

$$\text{and } L + l = s \quad (39)$$

$$\text{where } s = \sqrt{\text{type } s} = \sqrt{\bar{M}s + \bar{N}s + \bar{MN}s} \quad (40)$$

From these values of L and l , the values of L^s and l^s can be obtained with the aid of identities (30) and (31)

Stage 3 Here tests have been carried out with all four anti-serums, so that 9 phenotypes are distinguishable (cf. table 16). The calculations

are the same as at stage 2 except that now the frequency of gene s can be determined by direct count as follows

$$s = \overline{ss} + \frac{\overline{Ss}}{2} \quad (41)$$

$$= (\overline{Mss} + \overline{Nss} + \overline{MNs}) + \frac{\overline{MS} + \overline{NSs} + \overline{MNS}}{2}$$

After the gene frequencies have been estimated from the distribution of the phenotypes in the population the expected frequencies of the phenotypes can be recalculated with the aid of table 15 or table 16. Then, the chi square test can be applied as a test of the genetic theory. This has been done by Fisher,²² using maximum likelihood estimates of the gene frequencies, which differ only slightly from the estimates obtained with the simpler method (cf. Wiener²³) described here. For example, in one such test, a chi square value of 1.74 was obtained with 2 degrees of freedom, indicating a satisfactory agreement with the genetic theory.

Family Investigations When the factor S is considered by itself, family investigations support the hypothesis of heredity as a simple Mendelian dominant. For example, Rice and Sanger²⁴ have compiled 412 families with 941 children, and in 97 families where both parents were type s , all 221 children were likewise type s .

In family studies, when the bloods of all the individuals are tested for factors M , N and S simultaneously, the relationship between them becomes apparent immediately. In family 1 of table 17, the parents are type MN and type M respectively, and as expected about half of the children are type MN and half are type M . Also the parents are type S and type s respectively, and the former is evidently heterozygous, since three of the children are type S and four are type s . However, while the type M parent lacked the factor S the situation is reversed in the children, that is the type M children have the factor S . The reason for this becomes apparent when the phenotype of each individual is converted into the appropriate genotype in conformity with the genetic theory. The same method of analysis is readily applicable to other families in the table, and in published studies on the subject.

Significantly when in a family factor S is associated with blood factor M in a parent, it is similarly associated in his children and in other members of his family. The same holds for factor S when it is

TABLE 17

ILLUSTRATIVE FAMILIES DEMONSTRATING THE RELATIONSHIP BETWEEN
FACTOR S AND THE THREE M N TYPES(After Race and Sanger *Blood Groups in Man* 2nd ed p 68)

Family Number	Mat ing	Children						
		1	2	3	4	5	6	7
1	MN S × M s	M S	MN s	MN s	MN s	M S	MN s	M S
	L ^s l × LL	L ^s L	ll	ll	ll	L ^s L	ll	L ^s L
2	N s × MNS	MN s	N S	MN s				
	ll × Ll ^s	ll	l ^s l	ll				
3	MN S × N s	MN s	MN s	N S				
	ll ^s × ll	ll	ll	L ^s l				
4	MN S × MN s	MN S	N s	MN s				
	L ^s l × Ll	L ^s l	ll	ll				

associated with the factor N. To date, no evidence of crossing over between M/N and S/s has been found in family studies. This, in our opinion, indicates that factors S and M are not properties of different substances, but of one and the same agglutininogen, and thus supports the theory of multiple alleles in preference to that of linked genes.

Factor U. Wiener, Unger and Gordon,⁹⁵⁻⁹⁶ in 1954, encountered a new antibody in the serum of a woman who had a fatal hemolytic transfusion reaction. The factor it recognized was designated as U because of its almost universal distribution, as follows:

	Positive	Negative
Negroids	977	12
Caucasoids	1,100	0

The investigators also studied a family in which the father and two children lacked the factor while the mother and one child possessed the factor. This and other results supported the hypothesis that the blood factor is transmitted as a simple Mendelian dominant. It was noticed that the twelve individuals who lacked the factor U were all of type N or type MN, suggesting some relationship to the M/N types. Subsequently, in studies carried out with a second anti-U serum, Greenwalt et al.⁹⁷ confirmed these observations, and noted that all individuals lacking factor U also lacked both factors S and s. In this

TABLE 18
THE M-N-S-U SERIES OF ALLELIC GENES

Gene	Approximate Frequency (Per Cent) Among		Corresponding Agglutino- gen	Blood Factors Present*
	Caucasoids	Negroids†		
<i>L^S</i>	21.7	8.7	M _s	M, S, and U (and <i>ℑ</i>)‡
<i>L</i>	28.3	40.2	M _s	M, s, and U (and <i>ℑ</i>)‡
<i>I^u</i>	Extremely rare		M _u	M (and <i>ℑ</i>)‡
<i>I^s</i>	8.1	7.4	N _S	N, S, and U
<i>I</i>	38.9	43.7	N _s	N, s, and U
<i>i^u</i>	Extremely rare		N _u	N

* For simplicity in this chart the factors M, M_s, M_u, N, N_s, etc. are not listed (cf. page 46)

† Based on the data of Miller, Rosenfield, and Vogel, (*Amer. J. Phys. Anthropol.*, 9: 115, 1951)

‡ Agglutino-gen M reacts weakly with anti-N serum. This by definition, means that the agglutino-gen M has an N-like factor which may be designated *ℑ*

way, the series of alleles in the M-N system was increased to six as shown in table 18.

Other Complexities of the M-N-S System

In 1934, Landsteiner, Strutton and Chase⁹⁸ immunized rabbits with blood of a Negroid (Mr. Hunter) and obtained an anti-serum which agglutinated the blood of 7.3 per cent of Negroids and of 0.5 per cent of Whites. Wiener⁹⁹ also tested this serum and noticed that all blood giving distinct positive reactions were type N or type MN, suggesting a relationship to the M-N system.

Chalmers, Ikin and Mourant,⁹⁹ in an attempt to repeat these studies, obtained an anti-serum, prepared by injecting the blood of a Nigerian (Mr. Hen-law) into a rabbit. In tests with this anti-serum (anti He), 38 among 1,390 West Africans, or 2.7 per cent, gave positive reactions, while no positive reactions were obtained with the blood cells of 1,500 Europeans. Comparative tests with the original anti-Hu serum, some of which was still available, showed anti-Hu and anti-He to be different. With fresh blood from the original donor (Mr. Hunter) rabbits were immunized, and with the newly prepared anti-serum Chalmers et al., and Shapiro¹⁰⁰ carried these studies further. Family studies showed that factors He and Hu are related to the properties

M and N in a manner similar to that of factors S, s, and U. For example, when Hu or He were associated with M in the parents there was a similar association in the children. This again indicates that Hu and He do not represent attributes of separate substances, but together with M, N, S, s, and U, as the case may be, are factors of one and the same agglutinin. Because of these observations, the series of allelic genes for this system was extended to 24 by Shapiro, in order to account for the results of his studies in which the simultaneous heredity of M, N, S, s, U, Hu, and He was examined.

Additional complexities were added by the reports of two closely related blood factors designated Gr and Mi*, which have been shown to be similarly associated with the M-N S system.^{101 102 103} The most interesting of the recent developments is Allen's finding¹⁰⁴ of a potent serum for an extremely rare blood factor which he showed to be related to the M-N system. He reported a pedigree in which this blood factor occurred in 7 members of the family. In this family he encountered two apparent contradictions to the heredity of the three M-N types, namely, two brothers, both type N and each with a type M child. The paradox could be explained by postulating a new extremely rare allele which produced an agglutinin having the newly discovered blood factor but lacking both the blood factors M and N. If the allele is designated L^p , then the type N father's genotype could be LL^p while the type M child's genotype would be Ll^p .

CHAPTER IV

The P System

In 1927, when Landsteiner and Levine prepared the first anti M and anti N rabbit immune serums, they also obtained an antiserum which detected still another property of human blood that was designated by them as P. Subsequently they found the same antibody anti P, in the serums of occasional normal human beings and in animals (horse pig rabbits),¹⁰⁵ and it has since been found that the same antibody can occasionally result from immunization in response to blood transfusion.¹⁰⁶ In 1936 Furuhashi and Imamura¹⁰⁷ described a blood factor which they designated as Q, detected with the aid of absorbed pig serum. Japanese investigators have carried out many studies on factor Q but since comparative tests have shown P and Q to be the same, the original symbol P will be used exclusively in this discussion. It is of interest that among Japanese the frequency of the P (or Q) factor is only approximately 30 per cent in contrast to the frequency of 79 per cent among Caucasoids and 98 per cent in Negroids.¹⁰⁸

Family studies have shown the factor P to be inherited as a simple Mendelian dominant (cf. table 19).

Almost all anti P reagents available up to now have had the prop-

erty of sorption tend to weaken the anti P agglutinins as well. In addition, positively reacting cells from different individuals do not react with equal intensity, and all gradations from strong agglutination to weak clumping may be observed. Thus testing for the P factor presents difficulties similar to those of subgrouping blood of groups A and AB.

In a review on the origin of naturally occurring hemagglutinins Wiener¹⁰⁹ wrote: "Since anti P cold agglutinins occur not infrequently in normal human serum, the present author hazards the prediction that future studies on bacteria may reveal them to contain antigens related to agglutininogen P in specificity." In fact among 132 cases of *typhoid fever* of *adone* about 2% might be expected to be P negative. Cameron and Staveley¹¹⁰ found 2 with strong

anti P agglutinins. Of 25 cyst fluids, 19 inhibited anti P serum. In only 6 of the 25, where the scolices were absent or degenerated, did inhibition fail to occur. This confirms Wiener's theory of the heterogenetic origin of cold anti P agglutinins, the antigenic stimulus apparently being provided by the P like substance present in echinococcus cyst fluids containing scolices.

From the incidence of the P factor in the population, the gene frequencies are readily calculated. Thus, among Caucasoids, $p = \sqrt{(P-)} = \sqrt{0.21} = 0.46$ or 46 per cent, and $P = 1 - p = 0.54$ or 54 per cent. Similarly, among Negroids, $p = \sqrt{0.02} = 0.14$ or 14 per cent, and $P = 0.86$ or 86 per cent; while, among Chinese, $p = \sqrt{(0.70)} = 0.84$ or 84 per cent, and $P = 0.16$ or 16 per cent.

Factor Tj^a In 1951, Levine et al.¹⁰⁹ encountered in the serum of a woman an antibody which agglutinated the red cells of 3,000 consecutive random group O individuals, but did not react with the red blood cells of the donor herself or those of one of her three siblings. The factor detected by this serum was named the Jay or Tj^a factor after the donor, Mrs Jay. Since 1951, in eight additional investigations carried out in South Africa, Australia, Japan, the United States, and other parts of the world, anti-serums of identical specificity

TABLE 19
HEREDITY OF THE P AGGLUTINOGEN
(After Dahr and Zehner)

Mating	Number of Families	Number of Children		
		P+	P-	Totals
P+ × P+	240	677	79	756
P+ × P-	134	286	179	465
P- × P-	34	(4)	91	95
Totals	417	967	352	1319

TABLE 20
THE P SERIES OF ALLELIC GENES

Genes	Corresponding Agglutinogens	Blood Factors
P ¹	P ₁	P, Tj ^a
P ²	P ₂	Tj ^a
p	p	None demonstrated to date

were encountered. In a number of these studies the T_J^a negative type was found in more than one member of the family. It is of interest that many of the T_J^a negative individuals had the T_J^a antibody in their sera.

If one assumes that the T_J type is determined by a pair of allelic genes T_J^a and T_J^b then $T_J(a-)$ individuals would all be of genotype $T_J^b T_J^b$. The frequency of gene $T_J^a = \sqrt{1_J(a-)}$ or less than $\sqrt{0.00033} = 0.018$ or 1.8 per cent while the frequency of gene T_J^b is greater than 0.982 or 98.2 per cent.

Sanger¹¹⁰ tested 15 $T_J(a-)$ individuals from 9 families and found that they all were also P negative which suggested a relationship between the T_J^a factor and the P system. In fact she found that when anti T_J^a serum was absorbed with P negative blood until it no longer reacted with the absorbing blood an antibody fraction remained in the serum that gave reactions corresponding to anti P. To account for these findings Sanger has postulated the existence of three allelic genes P^1 , P and p at the P locus. The P^1 gene is postulated to determine the agglutinin found in the type previously classified as P positive. P^1 determines the agglutinin of P negative blood while p determines the agglutinin of $T_J(a-)$ blood. These concepts are summarized in table 20.

CHAPTER V

The Rh-Hr System

The Rh-Hr system is the most complicated and second to the ABO system is the most important of the human blood systems by virtue of its clinical implications. The past fifteen years have witnessed a phenomenal advance in the knowledge of blood group serology and genetics which was initiated by the discovery of the Rh-Hr types.

The Rh_0 Factor

Of the Rh-Hr factors the factor Rh_0 holds a position of special prominence. It is of both theoretical and historical interest that its discovery came about in an indirect manner with the use of an anti-serum prepared in rabbits against the us monkey blood.¹ The Rh_0 factor is the most common source of clinical symptoms because as demonstrated by experimental immunization studies carried out in human volunteer it is the most antigenic of the Rh-Hr factors.¹² It appears to represent a special structure within the Rh-Hr agglutinin since red cells can be coated with Rh_0 blocking antibody without interfering with the reactions of the red cells with other antibodies such as anti rh , anti rh' and anti hr . To indicate the special position of the Rh_0 factor therefore the symbol of this factor has been assigned a capital R while all of the other Rh factors have a small r . The subscript of the symbol Rh_0 also sets it apart from the other Rh-Hr factors.

In the first study on the heredity of the Rh-Hr types tests could be made only for the factor Rh_0 . In this investigation carried out by Landsteiner and Wiener¹³ in 1941 the anti-serums prepared in guinea pigs against the us monkey blood as well as human anti Rh_0 serum were used.

Among 448 persons tested with these serums the following distribution was found:

	Number	Per Cent
Rh positive	39	84.6
Rh negative	69	15.4

Under the hypothesis that the Rh_0 factor is inherited as a simple Mendelian dominant, the gene frequencies are readily estimated as follows

$$rh = \sqrt{Rh \text{ negative}} = \sqrt{0.154} = 39.2 \text{ per cent}$$

$$Rh = 1 - rh = 60.8 \text{ per cent}$$

The expected frequencies of the three theoretically possible genotypes can then be readily calculated

$$Rh \text{ positive} \begin{cases} \text{genotype } RhRh = (0.608)^2 = 37.0 \text{ per cent} \\ \text{genotype } Rhrh = 2(0.608 \times 0.392) = 47.6 \text{ per cent} \end{cases}$$

$$Rh \text{ negative, genotype } rh rh = (0.392)^2 = 15.4 \text{ per cent}$$

Family Studies. Landsteiner and Wiener tested 60 families with 237 children for the Rh_0 factor and in table 21 their findings are summarized. When the distribution of the Rh factor is compared with that expected under the genetic theory the findings prove to be in satisfactory agreement.

Landsteiner and Wiener also showed that the Rh_0 factor is not sex-linked and excluded any relation with the ABO and MN systems.

Serology. While in the original investigations anti-rhesus immune serum was used it was subsequently found that antiserums of greater avidity and higher titer are more readily obtained and in larger amounts from mothers of erythroblastic babies as well as from deliberately immunized donors. Therefore anti-rhesus serums are no longer used and remain only of theoretical and historical im-

TABLE 21
HEREDITARY TRANSMISSION OF THE Rh FACTOR
(From Landsteiner and Wiener 1941)

Mating	Number of Families	Number of Children		Totals
		Rh+	Rh-	
Rh+ × Rh+	42	151	7	158
Rh+ × Rh-	12	37	11	48
Rh- × Rh-	6	0	31	31
Totals	60	188	49	237

portance. It is interesting that the human anti-Rh₀ serum, even those of maximum titer and avidity, give little or no reaction with red cells of rhesus monkeys. This seemingly paradoxical observation is another example of so-called non reciprocal reactions, and is inexplicable under the hypothesis of a strict one to one correspondence between antigen and antibody.

It was early observed,¹¹⁴ in many Rh negative patients with clinical manifestations of iso-sensitization, that Rh antibodies could not be demonstrated by the classic agglutination method in saline media. This led Wiener in 1941 to postulate that Rh antibodies in particular (and all antibodies in general), exist in two molecular forms, one of which was capable of combining with Rh positive cells in saline media but without clumping the cells. In 1944, when suitable human anti Rh agglutinating serum was more easily available, Wiener¹¹⁵ devised the blocking test to demonstrate the second form of antibody. Later on, Race¹¹⁶ and Diamond¹¹⁷ independently encountered the Rh₀ blocking antibody when they tried unsuccessfully to produce a polyvalent anti-Rh₀' serum by pooling anti Rh₀ and anti rh' serums. The antibody which clumps cells in saline media is variously known as "bivalent", agglutinating, or "complete" antibody, while the antibody detected by the blocking test has been designated "univalent", blocking, conglutinating, or "incomplete" antibody.

The original blocking test detected univalent Rh antibodies only when present in high titer, and for this test a satisfactory agglutinating anti Rh serum is also needed. Fortunately, other more sensitive and more practicable tests for univalent antibodies have been devised, notably, the conglutination¹¹⁸ and albumin tests,¹¹⁹ the anti globulin test,¹²⁰ and the use of test cells treated with proteolytic enzymes.¹²¹ Since univalent antibodies are encountered far more frequently than bivalent antibodies, these new tests resulting from the discovery of the Rh factor, opened up the entire field of blood grouping, and may be credited with the phenomenal advances of the past 15 years.

Factor rh'

In 1941, Wiener¹¹⁴ encountered a patient who had a hemolytic transfusion reaction and whose serum contained an antibody which reacted with 70 per cent of Caucasoid individuals, in contrast with the original anti Rh serum which agglutinated the red cells of 85

per cent of such persons (cf table 22) By the use of a 2×2 contingency table, it was possible to show that the factor demonstrated by this serum was independent of the A B O, M N and P systems but related to the Rh_0 factor The problem then arose as to the nature of the relationship between the new blood factor, which is to-day known as rh' , and the original Rh_0 factor

It is instructive as well as of historical interest to give the reasoning as originally presented in the 1943 edition of Wiener's book *Blood Groups and Transfusion*, making necessary changes to correspond to modern terminology in order to avoid misunderstanding The quotation modified in this way reads as follows *

"The reactions could be explained most simply by postulating the existence of two agglutinogens Rh_0 and rh' corresponding to the agglutinins anti Rh_0 and anti rh' The four sorts of blood would then have the compositions Rh_0rh' , Rh_0 , rh' and Rh negative respectively This assumption would imply the existence, however of two corresponding genes R^0 and r' which would have to be either independent, linked or allelic The first two possibilities are excluded since the product of the frequencies $Rh_0rh' \times$ Rh negative is much greater than $Rh_0 \times rh'$ Moreover the existence of allelic genes R^0 and r' in individuals whose blood reacts with both anti Rh_0 and anti rh' would necessitate that this class not exceed 50 per cent † while the actual frequency is 70 per cent Accordingly the observations are best explained by assuming the existence of 3 qualitatively different Rh agglutinogens instead of only 2, one type reacting with anti Rh_0 serum but not anti rh' , a second reacting with anti rh' but not with anti Rh_0 and a third reacting with both sorts of anti Rh serums "

At this stage, the genetics of the Rh Hr system could be summarized as follows Four allelic genes are postulated Gene r gives rise to an agglutinogen lacking both factors Rh_0 and rh' , gene r' corresponds to an agglutinogen having only factor rh' , similarly gene R^0 controls an agglutinogen having only factor Rh_0 and finally gene R^1 deter-

* In the original quotation the anti Rh_0 serum was designated anti Rh_1 while anti rh' was designated as anti Rh_2

† On page 45 when discussing this question it was also pointed out that the frequencies of type MN and of group AB similarly could not exceed 50 per cent in a population at equilibrium

TABLE 22

RELATIONSHIP AMONG THE REACTIONS OF DIFFERENT ANTI Rh SERUMS

Anti serum	Originally Described by	Present Designation	Approximate Per Cent Positive	Reactions of Various Bloods among White Individuals			
				About 0°C	About 15°C	About 25°C	About 35°C
Type 1 (standard anti Rh)	Landsteiner and Wiener	Anti Rh ₀	80	Pos	Pos	Neg	Neg
Type 2	Wiener and Landsteiner	Anti rh	72	Pos	Neg	Pos	Neg
Type 3	Levine, Burnham, Katzin and Vogel	Anti Rh ₀	87	Pos	Pos	Pos	Neg

TABLE 23

THE FOUR PHENOTYPES DETERMINED BY FACTORS Rh₀ AND rh AND THEIR TEN THEORETICALLY POSSIBLE GENOTYPES

Phenotype	Reaction with Serum		Possible Genotypes
	Anti Rh ₀	Anti rh	
rh	—	—	rr
rh	—	+	r r and r r
Rh ₀	+	—	R ⁰ r and R ⁰ R ⁰
Rh ₁	+	+	R ¹ r, R ¹ R ⁰ , P ⁰ r, R ¹ R ¹ and I ¹ r

mines an agglutinin characterized by both factors Rh₀ and rh'. Then, corresponding to the four phenotypes there could theoretically be 10 possible genotypes as shown in table 23.

Wiener and Landsteiner¹²² studied the simultaneous heredity of factors Rh₀ and rh' in 47 families with 133 children. In this small series, unfortunately, there appeared no instance of the rarest of the four blood types, namely, type rh'. However, the results, as far as they went, were in full accord with the genetic theory.

Eight Rh Types

Factor rh'' Early in 1943, Wiener¹²³ described still another anti serum which agglutinated the red cells of only 30 per cent of the white population. Statistical analysis proved that this factor was re-

lated to the Rh Hr system, and to indicate that it is now known as **rh''**. Considering only the reactions of anti **rh'** and anti **rh''** serums, four types of blood are possible which could be designated as **rh**, **rh'**, **rh''**, and **rh'rh''**, respectively.* The distribution of these four types among 280 individuals proved to be as follows: type **rh**, 19.4 per cent, type **rh'**, 50.3 per cent, type **rh''**, 12.8 per cent, and type **rh'rh''**, 17.5 per cent. Since the product $rh' \times rh''$ is considerably greater than $rh \times rh'rh''$ it is clear that **rh'** and **rh''** are not independent. A tentative genetic theory was proposed by Wiener,¹² that the factors **rh'** and **rh''** are transmitted by triple allelic genes in a manner comparable to the heredity of the four A B O blood groups. Based on this hypothesis, the gene frequencies were readily estimated as follows:†

$$r = \sqrt{rh} = \sqrt{0.194} = 44.5 \text{ per cent}$$

$$r' = \sqrt{rh + rh'} - \sqrt{rh} = \sqrt{0.697} - \sqrt{0.194} = 39 \text{ per cent}$$

$$r'' = \sqrt{rh + rh''} - \sqrt{rh} = \sqrt{0.322} - \sqrt{0.194} = 12 \text{ per cent}$$

So that, $r + r' + r'' = 95.5 \text{ per cent}$

Thus, the agreement with the genetic theory was not entirely satisfactory, since the sum of the estimated gene frequencies fell short of 100 per cent, and the theoretical frequency of type **rh'rh''**, which should equal $2 \times r' \times r''$ or about 9.5 per cent, was considerably less than the observed frequency. These discrepancies were correctly attributed to difficulties in technique, due to the poor quality of the anti-serums available at that time. In addition, no allowance was made for additional genes giving rise to agglutinogens with both factors **rh'** and **rh''**. Such genes are now known to exist but their incidence in the population is so low that failure to take them into account must have been only a minor contributing factor to the deviation found in this early study.

The Eight Rh Phenotypes. When tests are made with all three anti-serums simultaneously, eight types of blood can be distinguished

* In the original paper, the four types were designated as Rh negative, Rh₁, Rh₂, and Rh₁Rh₂, respectively.

† The symbols for the genes as used here of necessity have a different meaning from the same symbols when used in connection with the calculations that include factor Rh₃, as will be seen shortly.

TABLE 21
 SCHEME OF THE EIGHT Rh TYPES
 (Wiener & S. Dale County M. A. B. R. 1919)

Rh Negative Types				Rh Positive Types			
Designation of types	Reaction with serum			Designation of types	Reaction with serum		
	Anti rh	Anti rh*	Anti Rh ₀		Anti rh	Anti rh*	Anti Rh ₀
rh	—	—	—	Rh	—	—	+
rh	+	—	—	Rh	+	—	+
rh	—	+	—	Rh ₂	—	+	+
rh rh	+	+	—	Rh ₁ Rh ₂	+	+	+

The types are named after the Rh blood factors (and agglutinogens) present in the red cells. Rh being short for Rh rh' (or Rh₀) and Rh₂ short for Rh rh'' (or Rh₁). Similarly, it is often convenient to use rh in place of rh rh' and Rh₂ in place of Rh Rh' (or Rh₀ Rh').

as shown in table 21. These eight phenotypes are easy to remember if they are arranged in the form of a double scheme of four types each, analogous to the four A B O blood groups.

The distribution of the types is different in diverse ethnic groups, and the results of few representative studies are shown in table 25. As can be seen from the table, Caucasoids have the highest frequency of type rh, while this type is virtually absent in Mongoloids. The most striking feature of negroid populations is the very high frequency of type Rh₀.

Genetic Theory. To account for the existence of the eight phenotypes, Wiener¹²⁴ found it necessary to postulate the existence of at least six allelic genes, as shown in table 26. Under this concept there are 21 theoretically possible genotypes corresponding to the eight possible phenotypes as shown in table 27.

Gene Frequency Analysis. The theory of multiple alleles to account for the eight Rh types has been tested by gene frequency analysis and by family studies. With the aid of table 27, the following formulae for the frequencies of the genes in terms of the frequencies of Rh phenotypes can readily be derived:

$$r = \sqrt{rh} \quad (31)$$

$$r' = \sqrt{rh' + rh} - \sqrt{rh} \quad (32)$$

$$r'' = \sqrt{rh'' + rh} - \sqrt{rh} \quad (33)$$

TABLE 25
REPRESENTATIVE DATA ON THE RACIAL DISTRIBUTION OF THE
EIGHT Rh TYPES

Ethnic Group	Approximate Frequency (Per Cent) of the Blood Types							
	rh	rh	rh*	rh rh*	Rh ₁	Rh ₂	Rh ₃	Rh Rh ₃
Caucasoids (N Y C) ¹	13.5	1.0	0.5	0.2	2.5	53.0	15.0	14.5
Negroids								
N Y C ¹	7.5	1.5	0	0	45.0	25.0	15.5	5.5
Africa ²	3.75	0.75	0	0	70.0	15.0	9.0	1.5
Puerto Ricans ³	10.1	1.7	0.5	0	15.1	39.1	19.6	14.0
Chinese ⁴	1.5	0	0	0	0.9	60.6	3.0	34.4
Japanese ⁵	0.6	0	0	0	0	51.7	8.3	39.4
Filipinos ⁶	0	0	0	0	0	87.0	2.0	11.0
Mexican Indians ⁷	0	0	0	0	1.1	48.1	9.5	41.2

(Modified after Wiener & S., *Am J Clin Path* 16: 477, 1946)

¹ Wiener et al., Unger et al.,
Levine

² Wiener et al., Levine

³ Hubinont et al.

⁴ Torregrosa

⁵ Wiener et al.

⁶ Waller and Levine, Miller and
Taguchi

⁷ Simmons and Crayden

⁸ Wiener et al.

$$R^0 = \sqrt{Rh_0 + rh} - \sqrt{rh} \quad (34)$$

$$R^1 = \sqrt{Rh_1 + rh' + Rh_0 + rh} - \sqrt{rh' + rh} \\ - \sqrt{Rh_0 + rh} + \sqrt{rh} \quad (35)$$

$$R^2 = \sqrt{Rh_2 + rh'' + Rh_0 + rh} - \sqrt{rh'' + rh} \\ - \sqrt{Rh_0 + rh} + \sqrt{rh} \quad (36)$$

As an example the formulae will be applied to the figures given for Caucasoids in table 25

$$r = \sqrt{0.135} = 36.7 \text{ per cent}$$

$$r' = \sqrt{0.010 + 0.135} - \sqrt{0.135} = 1.1 \text{ per cent}$$

$$r'' = \sqrt{0.005 + 0.135} - \sqrt{0.135} = 0.7 \text{ per cent}$$

$$R^0 = \sqrt{0.025 + 0.135} - \sqrt{0.135} = 3.3 \text{ per cent}$$

$$R^1 = \sqrt{0.530 + 0.010 + 0.025 + 0.135} - \sqrt{0.010 + 0.135} \\ - \sqrt{0.025 + 0.135} + \sqrt{0.135} = 12.3 \text{ per cent}$$

discovery of type $rh'rh''$ is one of the numerous facts which have been unearthed, supporting the theory of multiple alleles.

Family Studies Further evidence of the validity of the theory of multiple alleles was obtained by family investigations.¹²³⁻¹²⁶ In table 28 for example, are summarized the studies of Wiener and his associates during the period 1943 to 1953 on the heredity of the eight Rh types. Similar studies have been done by Race et al. and additional data have been accumulated more recently in investigations not yet published. With only two exceptions which may be ascribed to illegitimacy, the data obtained on the 2,119 children shown in table 28 support the theory of multiple alleles.

Actually a few other exceptions were encountered in certain families where one of the parents belonged to type $rh\,rh'$ or to type Rh_1Rh_2 . These apparent exceptions to the genetic theory could be explained by postulating the existence of two additional alleles in the series of allelic genes.¹²⁷⁻¹²⁹ Thus, under the theory of six allelic genes matings with parents $rh\,rh \times rh$ would have to be of genotypes $r\,r' \times r\,r$ so that half of the children would be expected to be type rh and half to be type rh'' . Actually in some of these families half of the children were type $rh'rh'$ while half were type rh . By postulating the existence of an additional allelic gene r'' , determining a corresponding agglutinin rh , having the blood factors rh' and rh'' but lacking factor Rh_2 , it was possible to account for these exceptional families. Similarly it was necessary to postulate the existence of a gene R' determining a corresponding agglutinin Rh_2 having all three factors Rh_1 , rh , and rh'' in order to account for anomalies in certain families with one or both parents of type Rh_1Rh_2 . In table 29 are shown three families illustrating the transmission of the rare genes r'' and R' .

Serology For identifying the 8 Rh types, it is necessary to have available antisera of specificities anti Rh_1 , anti rh' , and anti rh'' , respectively. Pure anti rh' serum can be obtained from type Rh_1 individuals sensitized against type Rh_1 blood, and, similarly pure anti rh'' serum is obtained from type Rh_1 individuals sensitized against type Rh_2 blood. Unfortunately, such monovalent antisera are rare and the reagents available commercially are mostly derived from type rh individuals sensitized to type Rh_1 or type Rh_2 blood respectively. Thus, the anti rh' reagents most commonly used are really anti Rh_1 sera to which anti Rh_2 blocker has been added to

TABLE 29

FAMILIES ILLUSTRATING THE TRANSMISSION OF THE RARE GENES R^s AND r^s

Mat. no.	Children				
	1	2	3	4	5
1 Rh ₁ Rh ₂ × rh $R^s r \times rr$	rh rr	Rh ₁ Rh ₂ $R^s r$			
2 rh rh × rh $r^s r \times rr$	rh rr	rh rh $r^s r$			
3 Rh ₁ Rh ₂ × rh $R^s r \times rr$	rh rr	rh rr	Rh ₁ Rh ₂ $R^s r$	rh rr	Rh ₁ Rh ₂ $R^s r$

prevent the anti Rh₀ agglutinin from giving rise to agglutination under the conditions of the test. Similarly, most anti rh'' reagents are anti Rh₀'' serums to which anti Rh₀ blocker has been added. If the directions for the tests are not followed scrupulously or when dealing with highly agglutinable red cells which are difficult to block, such as cells from homozygous type Rh₊ or type Rh₀ individuals blocking may be incomplete or may fail and the clumping resulting may be misinterpreted as indicative of the presence of the rh' and rh'' factor. Such errors have caused serious misunderstandings in medicolegal cases of disputed paternity, and have also been responsible for inaccurate reports in anthropological investigations.

Hr Factors

Levine and Javert¹²⁹ encountered an antibody in the serum of an Rh positive mother of an erythroblastotic baby which gave up approximately 30 per cent positive reactions with Caucasoid bloods. Significantly, all Rh negative bloods gave positive reactions with this serum. Because of its apparently reciprocal relation to Rh, they named the factor detected by this anti serum Hr. Independently, Race and Taylor¹³⁰ discovered an anti-serum for a blood factor present in 80 per cent of Caucasoid bloods. Wiener *et al*¹³¹ pointed out that factor Hr of Levine and Javert and the factor found by Race and Taylor were in fact the same, the difference in percentage of positive reactions being due to a gene dosage effect.

More detailed study showed that the so-called Hr factor was related to factor rh' in a manner similar to the relationship between M and N, that is, individual bloods are either rh' positive or Hr

positive, or both, and no blood was both rh' negative and Hr negative. To indicate this, Wiener introduced the designation hr' for the factor. It must be pointed out that the discovery of factor hr' made necessary no change in the genetic theory, since hr' merely defines another blood factor of agglutinogens already known and identifiable by their reactions with the three Rh antisera alone. The table of Rh allelic genes is therefore modified only by the addition of factor hr' in the "blood factor" column for agglutinogens rh , rh' , Rh_0 and Rh_2 (cf. tables 26 and 31). Thus, the number of possible genotypes remains the same but the number of possible phenotypes is increased. The additional phenotypes distinguished by the hr' antiserum have been designated as $rh'rh$ for those type rh' bloods which are positive for factor hr' , while those type rh' bloods which are negative for factor hr' are designated as $rh'rh'$. Similarly type Rh_1 bloods are subdivided by their reactions with hr' antiserum into type Rh_1rh for those bloods which react positively with hr' antiserum, and type Rh_1Rh_1 for those bloods which do not react with hr' antiserum.

Considering that factor rh' has an incidence of approximately 70 per cent in the Caucasoid population, the expected frequency of the reciprocally related factor hr' can be calculated with the aid of the formula

$$\begin{aligned}
 hr' \text{ pos} &= 1 - [1 - \sqrt{(rh' \text{ neg})}]^2 \\
 &= 1 - [1 - \sqrt{0.30}]^2 \\
 &= 1 - (1 - 0.548)^2 \\
 &= 1 - 0.204 \\
 &= 0.796 \text{ or } 79.6 \text{ per cent}
 \end{aligned}$$

This frequency, as can be seen, is very close to the value reported by Race and Taylor. If the factors rh' and hr' are considered by themselves, three phenotypes are distinguished, the expected frequencies of which would be as follows:

$$\begin{aligned}
 \text{type } rh' &= hr' \text{ negative} = 21\% \\
 \text{type } hr' &= rh' \text{ negative} = 30\% \\
 \text{type } rh'hr' &= \text{the rest} = 49\%
 \end{aligned}$$

By using an over simplification, it is possible to account for the existence of three such types. If we postulate a pair of alleles \bar{r} and \bar{h} (the bar is placed over the collective gene \bar{r} to distinguish it from gene r of the series of eight Rh alleles) there are three genotypes corresponding to the three phenotypes as follows:

Phenotype	Genotypes	Frequency %
rh	$\bar{r} \bar{r}$	21
rh hr	$\bar{r} \bar{h}$	49
hr	$\bar{h} \bar{h}$	30

It will be noted that while the incidence of the **hr** factor is 79 per cent among these 30 per cent are homozygous while 49 per cent are heterozygous. It is now known that blood from individuals homozygous for the **hr** factor reacts much more avidly with anti **hr** serum than blood from heterozygous individuals. This is known as the gene dose effect. It is probable that the original serum of Levine and Javert was detecting only homozygous **hr'** positive blood which accounts for their early observations.

Fisher¹³ noticing the reciprocal relationship between factors **rh** and **hr** postulated the existence of two additional Hr factors namely **hr''** and **Hr₀** respectively reciprocally related to factors **rh''** and **Rh₀** respectively. As will be explained later this prediction was associated with the genetic theory of linkage and a different nomenclature for the Rh Hr types. Shortly thereafter Mourant^{12a} announced the discovery of an antiserum giving the reactions expected of anti **hr''** and additional antisera of this specificity were found by Wiener and Peters^{12b} and other workers. Several announcements of the discovery of antisera supposedly of specificity anti **Hr₀** were also made but the findings could not be confirmed. Even at the present time anti **Hr₀** serum is not available although anti **hr'** and anti **hr''** sera are routinely used in many laboratories specializing in Rh Hr typing.

In table 30 are given the expected incidence of factors **hr'** and **hr''** as well as those expected of the hypothetical factor **Hr₀** among Whites, Negroes, Mongoloids and Arctic Indians estimated from the incidence of factors **rh**, **rh''** and **Rh₀** in these populations as calculated with the aid of the general formula:^{12a}

$$(\text{Hr}+) = 1 - \{1 - \sqrt{(\text{Rh}-)}\}^2 \quad (37)$$

TABLE 30
RELATIONSHIP BETWEEN Rh AND Hr ANTISERUMS

Racial Group	Observed* Percentages of Positive Reactions with Rh Antiserums			Expected Percentages of Positive Reactions with Hr antiserums		
	rh	rh'	Rh ₀	hr	hr'	Hr ₀ (Hypothetical)
Whites	70	30	85	80	97	63
Negroes	28	27	90	97.7	97.5	54
American Indians, Chinese and Japanese	85 to 95	40 to 60	99 to 100	43 to 63	86 to 95	0 to 20
Asiatic Indians	85	18	93	63	99.1	45

* A. S. Wiener, E. B. Sonn and R. B. Belkin, *Proc. Soc. Exp. Biol. and Med.*, 54: 238, 1943; A. S. Wiener, J. P. Zepeda, E. B. Sonn and H. R. Polivka, *J. Exp. Med.*, 81: 559, 1945.

Nomenclature. With the discovery of additional Rh-Hr factors, the relationships among the various agglutinogens become increasingly complicated, and this entails corresponding complexities in the notations for the various Rh-Hr types. When choosing the symbols for the Rh-Hr types, rational principles of notation were applied so that the resulting nomenclature is not only accurate and unambiguous, but also serves to translate the serological facts into symbols which can be easily manipulated both graphically and orally. Thus, no attempt is made to include everything that is known about a given agglutinin in its symbol, since this would entail repeated changes in the notation with the discovery of each new fact. Rather, the symbols chosen are in the nature of easily inscribed and spoken mnemonics, which aid the memory and at the same time stress only those characteristics needed for the identification of the agglutinin or phenotype. For example, agglutinin rh is defined and identified by its negative reactions with antisera Rh₀, rh', and rh''. With the discovery of anti-hr' and anti-hr'', no change had to be made in the symbol; instead, the definition of agglutinin rh was merely expanded to include its reactions with these two additional antisera. Thus, agglutinin rh is now more fully defined as that agglutinin which gives negative reactions with antisera Rh₀, rh', and rh'',

TABLE 31
THE EIGHT STANDARD Rh ALLELIC GENES

Gene	Corresponding Agglut. nogen	Blood Factors Present
<i>r</i>	rh	hr and hr
<i>r</i>	rh	rh and hr
<i>r</i>	rh	rh and hr
<i>r^v</i>	rh _y	rh and rh
<i>R⁰</i>	Rh ₀	Rh ₀ hr and hr
<i>R</i>	Rh ₁	Rh ₀ rh and hr
<i>R²</i>	Rh	Rh ₀ rh and hr
<i>R³</i>	Rh ₂	Rh ₀ rh and rh

and positive reactions with anti hr and anti hr^v. As was anticipated additional characteristics of agglutininogen rh have been discovered notably, hr, and more recently hr^v. It will be seen that the factor hr is not essential for the identification of agglutininogen rh, and therefore need not be included in its symbol. On the other hand since factor hr^v identifies a property of only *certain* rh agglutinogens it does require a distinctive symbol to indicate its presence.

In table 31 the situation resulting from the discovery of the additional two factors hr and hr^v is summarized for the 8 standard allelic genes. When the three Rh antisera and the two Hr antisera are used simultaneously, instead of only 8 Rh types a scheme of 18 Rh Hr phenotypes results as is shown in table 32. In order to indicate the reactions of the Hr antisera a new convention has to be introduced, as has already been indicated. To facilitate the use of this convention the names of two of the phenotypes are changed as follows: instead of rh'rh'' the symbol rh_y is used and instead of Rh₁Rh₂ the symbol Rh₂ is employed. In this way it is possible to indicate not only the results of the tests but also which antisera have been used. Thus, when tests are made using only the three Rh antisera a single symbol is used to indicate the phenotype, while a second symbol is added when tests have been carried out for the Hr factors as well. Since all type Rh₁ bloods necessarily give positive reactions with anti hr'' serum * this fact need not be indicated in the phenotype symbol. On the other hand to indicate the reaction with anti hr' the symbol Rh₁Rh₁ is used for type Rh₁ bloods which are hr' negative, while the symbol Rh₁rh is used for type Rh₁ bloods

* Rare exceptions to this rule will be described later

TABLE 32
NOMENCLATURE OF THE 18 Rh Hr BLOOD TYPES AND 36 GENOTYPES

Eight Rh Phenotypes	Reactions with Serum		Designation	
	Anti-hr	Anti-hr'	Phenotypes	Corresponding genotypes
rh	+	+	rh	<i>rr</i>
rh	+	+	rh rh	<i>r r</i>
	-	+	rh rh	<i>r r</i>
rh	+	+	rh rh	<i>r r</i>
	+	-	rh rh	<i>r r</i>
rh _y (or rh rh')	+	+	rh _y rh	<i>r r</i> and <i>r^yr</i>
	-	+	rh _y rh	<i>r^yr</i>
	+	-	rh _y rh	<i>r^yr</i>
	-	-	rh _y rh _y	<i>r^yr^y</i>
Rh ₀	+	+	Rh ₀	<i>R⁰R⁰</i> and <i>R⁰r</i>
Rh	+	+	Rh ₁ rh	<i>R¹r</i> <i>R¹R⁰</i> and <i>R⁰r</i>
	-	+	Rh ₁ Rh ₁	<i>R¹R¹</i> and <i>R¹r</i>
Rh ₂	+	+	Rh rh	<i>R²r</i> <i>R²R⁰</i> and <i>R⁰r</i>
	+	-	Rh Rh	<i>R²R²</i> and <i>R²r</i>
Rh ₂ (or Rh ₁ Rh ₂)	+	+	Rh ₂ Rh ₀	<i>R¹R²</i> <i>R r</i> <i>R²r</i> <i>R¹r</i> <i>R¹R⁰</i> and <i>R⁰r^y</i>
	-	+	Rh ₂ Rh ₁	<i>R²R¹</i> <i>R²r</i> and <i>R r^y</i>
	+	-	Rh ₂ Rh	<i>R²R²</i> <i>R²r</i> and <i>R¹r^y</i>
	-	-	Rh ₂ Rh ₂	<i>R²R²</i> and <i>R¹r^y</i>

which are hr positive. Thus the first half of the phenotype symbol represents the reactions obtained with the Rh anti-serums, while the second half of the symbol represents the reaction with the Hr anti-serums. As shown below there is generally a choice of two symbols to represent the Hr reactions.

rh or Rh₀ represents presence of both factors hr' and hr''

rh' or Rh₁ represents absence of hr' and presence of hr''

rh'' or Rh₂ represents presence of hr' and absence of hr''

rh_y or Rh₂ represents absence of both factors hr' and hr''

As shown in table 33 the symbol actually assigned to represent the Hr reactions, in most but not all cases makes the complete phenotype symbol correspond to the most frequent of the possible genotypes. It must be stressed that the adoption of this convention does not commit one to a specific genotype, since the symbol for phenotypes and genotypes always remain distinctive and unambiguous.

Family Studies Studies on the heredity of the Rh Hr blood types (136-142), support the hypothesis that hr is related to rh' , and hr'' is related to rh'' , as factor M is related to N. Many studies have been carried out with anti hr' serum but only a few studies have been made using anti hr'' serum, because the latter is less readily available. The rules of heredity can be summarized as follows:

1. Factor hr or hr'' cannot appear in the blood of a child unless it is present in one or both parents.

TABLE 33

STATISTICAL ANALYSIS OF R. R. RACE'S DATA ON THE DISTRIBUTION OF THE Rh Hr TYPES IN LONDON

Rh Hr Phenotypes	Number	Per Cent
rh	170	15.84
rh, rh	0	0
rh, rh	10	0.93
rh, rh	0	0
rh, rh	7	0.65
rh, rh	0	0
rh, rh		
rh, rh		
rh, rh		
Rh_0	19	1.77
Rh, Rh	130	17.71
Rh, rh	363	33.83
Rh, Rh	29	2.70
Rh, rh	137	12.77
$Rh_2 Rh_0$	141	13.42
$Rh_2 Rh_1$	4	0.37
$Rh_2 Rh_2$	0	0
$Rh_2 Rh_2$		

$r = 39.80$ $r = 1.15$ $r = 0.81$ $r = 2.19$ $r = 40.57$, $R = 15.98$ $R = 0.41$

TABLE 33 $\Sigma R = 100.21$

$\sqrt{(rh -)} + \sqrt{(hr -)} = 100.59$ per cent $\sqrt{(rh -)} + \sqrt{(hr -)} = 100.11$ per cent

- 2 Parents who are rh' negative cannot have hr' negative children, and hr' negative parents cannot have rh' negative children
- 3 Similarly, rh'' negative parents cannot have hr'' negative children and hr'' negative parents cannot have rh'' negative children

Only rare exceptions to these rules have been encountered, all presumably accounted for by illegitimacy. Significantly, when mother-child combinations are studied, as in material from medicolegal cases of disputed paternity, no contradictions to the second law are found. One recent exception to this statement was encountered by Levine,¹⁴ and evidence has been obtained that this baby had been interchanged with another at a foundling home.

Population Studies. Population studies in which Hr tests as well as Rh tests have been carried out yield data from which gene frequencies can be readily calculated. Formulas for these gene frequencies are derived which complement those based on data for the 8 Rh types (cf formula (31)–(36)). For example phenotype Rh_1Rh_1 includes genotypes R^1R^1 and R^1r' while phenotype $rh'rh'$ corresponds to genotype $r'r$.

$$\text{Therefore, } R^1 + r' = \sqrt{Rh_1Rh_1 + rh'rh'}$$

$$\text{Similarly, } R^2 + r'' = \sqrt{Rh_2Rh_2 + rh''rh''}$$

Since phenotypes $rh'rh'$ and $rh''rh''$ are both extremely rare the following approximations may be used

$$R^1 = \sqrt{Rh_1Rh_1} - r' = \sqrt{Rh_1Rh_1} - \sqrt{rh' + rh} + \sqrt{rh} \quad (38)$$

$$\text{similarly, } R^2 = \sqrt{Rh_2Rh_2} - \sqrt{rh'' + rh} + \sqrt{rh} \quad (39)$$

The frequency of gene R^2 can be estimated from that of phenotype Rh_2Rh_1 which includes genotypes R^2R^1 , R^2r' and R^1r'' . If we disregard the latter two genotypes which are rare in comparison to the first it can be seen that the frequency of gene R^2 is derived by dividing the frequency of phenotype Rh_2Rh_1 by twice the frequency of gene R^1 , as given by the formula (38) above. As far as the frequency of the gene r'' is concerned this is so low that a similar method of calculation cannot be used. However, the frequency of gene r'' can be estimated from the fact that in family studies, among Caucasoids, approximately half the parents of phenotype $rh'rh''$ prove to be genotype $r'r''$ and about half are genotype $r''r$.

The rarest genotype is $r''r''$, corresponding to phenotype rh_1rh_1 ,

with an estimated frequency of about 1 in 100 million among Caucasoids McGee et al¹⁴ have recently encountered such an individual and, as might be expected, he was derived from a consanguineous marriage

When the gene frequencies are estimated with the aid of the formulae, their sum is found to be close to 100 per cent, as expected

In view of the reciprocal relationship between rh' and hr' , and between rh'' and hr'' the following should hold

$$\sqrt{rh' \text{ neg}} + \sqrt{hr' \text{ neg}} = 100 \text{ per cent} \quad (40)$$

$$\text{and } \sqrt{rh'' \text{ neg}} + \sqrt{hr'' \text{ neg}} = 100 \text{ per cent} \quad (41)$$

In table 33, Race's data on the distribution of the Rh Hr types in London have been subjected to statistical analysis and it can be seen that they satisfy the requirements of the genetic theory

Serology Anti hr' serums are generally obtained from type Rh_1Rh_1 individuals who have been sensitized by pregnancy or by transfusion, and similarly, anti hr'' serums are obtained from type Rh_2Rh_2 or sensitized patients Attempts to produce anti hr' and anti hr'' serums in volunteer donors by deliberate injections of type rh blood have only occasionally been successful, which corresponds with the clinical observation that cases of hr' and hr'' sensitization are far less common than cases of Rh_0 sensitization A type Rh_1Rh_1 individual sensitized to type Rh_1rh (or type rh) blood will produce pure anti hr' , but if he is sensitized to type Rh_1Rh_2 blood he may produce the two antibodies anti hr' and anti rh'' (When an individual of the rare type Rh_2Rh_1 is sensitized his serum will necessarily contain only anti hr')¹⁵ Polyvalent anti $hr'rh''$ serum is difficult to differentiate from pure anti hr' , unless the two antibodies in the former have unequal titers or react by different methods of testing With the aid of blood of the rare type Rh_2Rh_1 however, the distinction can easily be made since such blood is hr' negative but rh'' positive Obviously failure to recognize the presence of anti rh'' in an anti hr' reagent can lead to the incorrect typing of type Rh_2Rh_1 blood as Rh_2Rh_0 Similarly, many anti hr'' reagents also contain anti rh' , and here blood of the rare type Rh_2Rh_1 is valuable for eliminating this possible source of error

An important peculiarity of anti hr' and anti hr'' serums is that they give a much more pronounced gene dosage effect than Rh anti

serums. As has already been pointed out failure to take this into account has given rise to errors in Rh Hr typing.

Additional Rh-Hr Factors

Factor rh^w Callender and Race¹⁴⁶ detected the antibody for this blood factor in serum of a woman who had lupus erythematosus and who had been immunized by numerous transfusions. She belonged to type Rh₁Rh₁ and became sensitized to factors hr and N, and to three previously undescribed blood factors of which rh^w was one. The superscript w was chosen because the name of the patient was Willis. The incidence of the new factor is relatively low—only 3 to 7 per cent among Caucasoids. A striking feature of the factor is that all persons possessing the factor also possess the factor rh. (Generally however blood having factor rh^w reacts more weakly with anti rh serum than blood lacking factor rh^w.) The existence of this factor therefore subdivides type Rh₁ for example into two types—one with and the other without the rh^w factor. These findings make it necessary to increase the number of Rh allelic genes since in place of gene R¹ there must be substituted two genes R^{1w} and gene R proper. Similarly evidence has been obtained for the existence of two kinds of r gene namely r^w and r proper.¹⁴⁷ It is possible that genes R^{1w} and r^w do exist but if they do they must be extremely rare. Family studies on the heredity of the rh^w factor support the genetic theory.^{148, 149}

Factor hr Roenfield et al.^{150, 151} encountered an antiserum whose reactions at first seemed to correspond to those expected of anti Hr. Family and population studies showed that the new serum designated anti f by the discoverers but to which the designation anti hr has been assigned by Wiener,¹⁵¹ reacted with blood cells from all individuals carrying genes r and R⁰ and did not react with blood from persons lacking these genes.

In order to crystallize the relationships among the agglutinogens determined by the blood factors described up to this point—namely Rh₁, rh', rh^w, rh'', hr, hr'', and hr, the Rh-Hr allelic genes and their corresponding agglutinogens identifiable with the aid of antisera—for these seven blood factors are summarized in table 34. As can be seen it is necessary to postulate a minimum of ten Rh allelic genes. As has already been pointed out, the discovery of the hr factor requires no extension of the series of allelic genes since it merely defines

TABLE 34

THE Rh Hr ALLELIC GENES IDENTIFIABLE WITH ANTISERUMS FOR FACTORS
Rh₀, rh, rh', rh'', hr, hr' and hr

Genes	Frequency among Caucasoids (per cent)	Corresponding Agglutinogens	Blood Factors Present
<i>r</i>	38	rh	hr hr hr
<i>r</i>	0.6	rh	rh hr
<i>r'</i>	0.003	rh'	rh rh' hr
<i>r</i>	0.5	rh	rh hr
<i>r''</i>	0.01	rh''	rh rh
<i>R⁰</i>	2.7	Rh ₀	Rh ₀ hr hr hr
<i>R¹</i>	41	Rh	Rh ₀ rh hr
<i>R'</i>	2	Rh'	Rh ₀ rh rh' hr
<i>I¹</i>	15	Rh	Rh ₀ rh hr
<i>R²</i>	0.2	Rh ₂	Rh ₀ rh rh

Omitted from the table are the genes *r''* and *R''* both presumably very rare and of which only a single family with *r''* has been found to date.

an additional serological attribute which is shared by the two agglutinogens rh and Rh₀. These two agglutinogens could formerly be identified without the aid of serum anti hr. However the anti-serum does help in distinguishing between individuals of genotypes *r'r''* and *r'r* by serological means alone because the former are hr negative while the latter are hr positive. Similarly, in persons of type Rh₁Rh₂ the use of this anti serum helps in making the distinction between those who do and those who do not carry the gene *R'* and this not infrequently is of value in cases of disputed parentage.

The number of genotypes possible with *n* alleles can be shown to be equal to $\frac{1}{2}n(n+1)$ so that the 10 Rh Hr alleles give rise to 55 possible genotypes. When all of the seven anti-serums that have been described thus far are available as many as 28 phenotypes can be distinguished corresponding to the 55 genotypes. For medicolegal work the use of anti-serums for factors Rh₀, rh', rh'', and hr is standard practice while the less readily available anti-serums anti hr' anti rh' and anti hr are used only in a limited number of laboratories. Table 35 is a systematized tabulation of the phenotypes which can be identified according to the reagents used, together with their corresponding genotypes.

Factor hr' In 1936 DeNittile et al.¹³² encountered an anti-serum for still another factor of the Rh Hr system. This factor proved to

TABLE 35

THE Rh Hr PHENOTYPES AND GENOTYPES
International Nomenclature

2 Rh Phenotypes			12 Rh Phenotypes				28 Rh Hr Phenotypes				55 Genotypes*
Des g nation	Approx imate frequencies in whites (%)	Re act on with anti Rha (or anti rhesus)	Des g nation	Approx imate fre- quencies in whites (%)	Reaction with		Designation	Approx imate fre quencies in whites (%)	Reaction with		
					Anti rh	Anti rh* Anti rh*			Anti hr Anti hr*	Anti hr	
Rh nega tive	15	-	rh	14.4	-	-	rh	14.4	+	+	rr
			rh'	0.46†	+	-	rh'rh rh rh	0.46 0036	+	+	r'r' r r'
			rh''	0.01	+	-	rh''rh rh''rh'	0.01 00006	+	+	r''r'' r''r'' or r''r''''
			rh'''	0.38	-	+	rh'''rh rh'''rh''	0.38 0025	+	+	r'''r''' r'''r'''
			rh''''	0.1	+	+	rh''''rh rh''''rh rh''''rh rh''''rh'' rh''''rh	0.06 0.08 0.001 0.001 0.00001	+	+	r''''r'''' r''''r'''' r''''r'''' r''''r'''' r''''r''''
			rh'''''	0.00035	+	+	rh'''''rh rh'''''rh'	0.00035 0.00001	+	+	r''''r''''' r''''r'''''

have an incidence of approximately 40 per cent among Africans 27 per cent in New York City Negroids, and only about 0.5 per cent among Caucasoid. It was noted that all individuals possessing the factor were also carriers of the genes r or R^0 or both, so that whenever the new factor named I is present factor hr was also present. Thus this factor bears the same relationship to factor hr that factor rh^+ bears to factor rh . For this reason it is proposed to call this factor hr^+ . Evidently in Negroids there are at least two varieties of gene r and gene R^0 namely gene r^+ and gene r proper, and gene R^{0+} and gene R^0 proper respectively. The use of anti hr^+ serum increases the number of phenotypes that can be distinguished and appropriate name for the e can readily be devised.

Complexities of the Rh_0 Factor and Associated Factors

Old blood suspensions generally give weaker agglutination reactions than fresh blood suspensions. Even when fresh blood exclusively is tested variations in intensity of the reactions are encountered dependent on the individual's blood type. For example as already pointed out blood cells from homozygous individuals not infrequently give more intense reactions than blood from heterozygous individuals—the so-called gene dose effect. In addition group A.B cells react more weakly than A_2 cells with anti A serums and type Rh_1 cells react more weakly than type rh' cells with anti rh serum. This suggests that the A.B.O. and Rh.Hr agglutinogens are each formed from a separate substrate and the presence of other structures in the agglutinin molecule may interfere with the agglutinability of the cell e.g. the presence of factor Rh_0 in type Rh_1 cells interferes with the rh /anti rh reaction and the presence of rh^+ still further reduces the agglutinability of the cell by anti rh . Aside from such phenomena agglutinability may be affected by minor variations in the structure of the agglutinin molecule itself causing corresponding minor changes in the blood factors such as in agglutinogen A_1 , A_2 and A_3 agglutinogens N and N_2 etc. Similar variations in the Rh.Hr agglutinogens especially those involving factor Rh_0 will now be discussed.

In 1914 Wiener¹³¹ showed that there were certain blood specimens which give weak but distinct reactions with certain of the Rh anti serums. These blood specimens appear to occupy a position inter

mediate between negatively reacting bloods and those which gave sharp positive reactions. Family studies indicated that this characteristic is hereditary and because of this and their serological behavior these types were designated as intermediate Rh types and the genes called 'intermediate genes'. Of the various intermediate Rh Hr variants the Rh_0 variants designated \mathfrak{Rh}_0 are the most important. As will be seen, recent investigations confirm the impression that the Rh_0 factors hold a special central position in the Rh Hr system.

Factor \mathfrak{Rh}_0 . The so called factor \mathfrak{Rh}_0 actually comprises a group of factors giving reactions of intermediate intensity with anti Rh_0 serum and the symbol \mathfrak{Rh}_0 has been used for the entire group of variant blood specimens which react typically with most anti Rh_0 serum but give slightly weaker reactions with a number are known as high grade variants while those on the other end of the scale reacting consistently weakly with all anti Rh_0 serums are low grade variants. Most of the variants give similar reactions which are characterized as follows. In saline media, they fail to react with anti Rh_0 serums even with those containing potent anti Rh_0 agglutinins. They frequently but not always react with univalent Rh_0 antisera by the conglutination method. They almost always react with such serums by the firm technique and characteristically react with such serums by the anti globulin technique although the avidity and titer of the reactions are generally lower than for typical Rh_0 positive bloods. Interestingly, the bloods of chimpanzees give reactions resembling those of high grade Rh_0 variants.

In 1915 Wiener, Under and Sonn¹⁶⁵ pointed out that Rh_0 variants are relatively rare among Caucasoids and occur more frequently among Negroids. Moreover it has been found that the factor \mathfrak{Rh}_0 like the factor Rh_0 may occur alone or together with rh or rh' , or both. To account for these findings the existence of at least three

and \mathfrak{Rh}_0
ively
had

been designated as type rh or type rh' were actually type \mathfrak{Rh}_0 or \mathfrak{Rh}' respectively. An extensive study on this question was carried out by Roenfield et al.¹⁶⁶ on a series of white blood donors in New York City. The blood specimens were first tested with saline agglutinating anti Rh , anti rh' , and anti rh'' serums and then all apparently Rh negative bloods were tested for Rh variants using the anti

globulin technique. It was found that of 161 seeming type $rh'rh$ as many as 72 or 44.7 per cent were actually type Rh_1rh , of 92 seeming type $rh''rh$ as many as 19 or 20.7 per cent proved to be type \mathfrak{Rh}_2rh while of 628 seeming type rh only 3 or 0.5 per cent were actually type \mathfrak{Rh}_0 . These results indicate that among Caucasoids of the three postulated additional rare alleles R^1 is the most frequent R^2 is next in frequency while R^0 is by far the rarest.

These findings argue against the idea that the existence of the Rh_0 variants is due to one or more modifying genes for unless such genes were completely linked to the Rh Hr genes the distribution of the Rh_0 variants would be expected to be largely independent of the Rh Hr types. A disturbing finding is the not infrequent occurrence of individuals of type Rh_1Rh_1 whose blood cells give weak reactions with anti Rh_0 serums resembling those of high grade variants. By family studies Ceppellini¹²⁷ has shown that these persons are always of genotype $R'r$ and that in other type Rh_1 individuals in the same family such as those of genotype $R'r$ the bloods give reactions of typical Rh_0 positives. Ceppellini therefore believes that the r gene in these families has a special modifying action which is responsible for the weak reactions obtained with Rh_0 antisera with the red cells of such individuals. Levine et al.¹²⁸ have recently reported a large pedigree confirming Ceppellini's hypothesis and similar observations have been made by Wiener.

The practical clinical importance of the Rh_0 variants is that transfusion of such blood to an Rh_0 negative individual may give rise to sensitization to the Rh_0 factor. As will be discussed later on in greater detail a few Rh_0 positive individuals have become sensitized to a factor related to the Rh_0 factor.

Nomenclature. The difference between factors Rh_0 and \mathfrak{Rh}_0 is not on the same plane as the difference among Rh_0 , rh and rh'' , although both are of a qualitative nature. Thus although the reactions of factor Rh_0 include all of the reactions of factor \mathfrak{Rh}_0 the reverse is not true since \mathfrak{Rh}_0 shares only some of the reactions of Rh_0 . Moreover the difference between Rh_0 and \mathfrak{Rh}_0 is not on the same plane as that between rh' and rh'' , because while serum for rh'' does not react with rh' , serum for \mathfrak{Rh}_0 always reacts with Rh_0 . Type rh individuals immunized with type \mathfrak{Rh}_0 blood produce antisera which like ordinary anti Rh_0 serums react more strongly with blood having the Rh_0 factor than those containing the \mathfrak{Rh}_0 factor. Thus in the

case of factors Rh_0 and $\bar{R}h_0$, the distinctions made are not those of sharply differentiated blood factors, but among members of a graded series of related factors. To indicate this it seems more appropriate to change the type face of the capital R to Germanic capital \mathfrak{R} than to coin a new symbol. As will be pointed out, there are also variants of the Rh_0 factor which react more strongly with anti Rh_0 serum than does ordinary Rh_0 positive blood. To indicate this the symbol of these Rh_0 variants is changed by using a capital R with a bar over it \bar{R} .

Variants of the factors rh' and rh'' also have been encountered and in the past the custom has been to enclose the prime sign and the double prime sign in parentheses to indicate such variants. Like bloods with variants of factor Rh_0 , bloods with variants of factors rh' and rh'' are rare.

Factor $\bar{R}h_0$. In 1950 Race, Sanger and Selwyn¹⁵⁹ found in the serum of the mother of a baby who died of erythroblastosis fetalis an antibody which clumped 1,400 consecutive blood specimens, but failed to clump her own cells. The mother's red cells proved to be unique in that they reacted with anti Rh_0 serum but appeared to lack both factors of the pairs $rh' hr'$ and $rh'' hr''$. To account for this observation, the investigators postulated that a deletion had occurred in the Rh 'chromosomes', and asserted that the genotype of the mother was $-D / -D-$ (cf. page 97).

The following interpretation appears more consistent with established serological and genetical principles. As Race et al. noticed in addition to the peculiarities mentioned above, the red cells were agglutinated in saline media with anti Rh_0 blocking serum which failed to clump ordinary Rh_0 positive cells under identical conditions. Thus, this individual had an agglutininogen similar to agglutininogen Rh_0 , but differing in its higher reactivity with anti Rh_0 serum.¹⁶⁰ It is this property which we consider the fundamental characteristic of this agglutininogen, due to a modification in its chemical structure which increases the reactivity with anti Rh_0 serums, and simultaneously crowds out the site of reactivity with all anti Hr serums. This new type of antigen we therefore propose to designate 'super' Rh_0 and assign the symbol 'bar' Rh_0 (or $\bar{R}h$) to its blood factor. Corresponding to agglutininogen $\bar{R}h_0$ the existence of another allelic gene \bar{h}^0 is postulated.

The existence of gene \bar{h}^0 can give rise to contradictions to the rule

that an rh negative parent cannot have an hr' negative child and an rh'' negative parent cannot have an hr negative child, and this must be taken into account in the medicolegal applications of the tests. For example, a mother whose genotype is $R\bar{R}^0$ and who is therefore hr negative could have a child who is rh negative. Fortunately the gene R^0 betrays its presence, even in its heterozygous form by the agglutination of the individual's red cells with univalent anti Rh_0 serums in saline media. In practice such cases will rarely occur because of the extreme rarity of gene R^0 but the medicolegal expert must be aware of this possibility if serious mistakes are to be avoided.

The gene R and its corresponding agglutminogen $\bar{R}h$ will probably prove to be the prototype of a whole series of such Rh Hr genes and corresponding agglutminogens. Thus corresponding to the variants Rh_0 , Rh_1 , Rh_2 etc. one may anticipate the existence in addition to $\bar{R}h_0$ of $\bar{R}h_1$, and $\bar{R}h_2$ the latter being characterized by their "super Rh_0 " factors. In fact Gunson and Donahue¹⁶ have found an individual of type $\bar{R}h^*$ who by family studies was shown to be homozygous for a new allelic gene which may be designated as R^* . His red cells lacked the four blood factors rh , hr , rh'' and hr'' .

The feature of both blood specimens $\bar{R}h_0$ and $\bar{R}h^*$ which attracted the attention of the discoverers was the simultaneous absence of both factors of the pairs rh hr and/or rh'' hr'' . According to the interpretation proposed here this is to be expected in any individual homozygous for a "super R^0 " gene. Such individuals are extremely rare and it is therefore not surprising that in both cases they were the products of consanguineous marriages. To detect the more common heterozygotes advantage should be taken of what we regard as the more fundamental characteristic of these agglutminogens, namely their high agglutinability with anti Rh_0 serums. Ordinary Rh Hr typing would not detect these carriers of a super R^0 gene since the other gene of the pair determining the genotype would provide reactions for each of the factor pairs rh' hr' and rh'' hr'' .

Theoretical Serological Considerations. Up to the present it has been impossible to demonstrate conclusively the existence of anti Hr_0 , although theoretically there seems to be no reason why a factor bearing the same reciprocal relationship to factor Rh that factor hr' has to factor rh' and factor hr'' has to factor rh'' should not exist. A possible explanation for this may be found in the following

analysis, based on the known facts concerning factor Rh and its variants Rh_0 and Rh_1 .

It has already been pointed out how the presence of one structure of factor in the Rh Hr agglutinogen molecule may interfere with the development of other Rh Hr factors, as judged from the serological reactions. Similarly, it might be anticipated that the more prominent the structure responsible for the reactions which are obtained with anti Rh_0 serums, the less prominent would be the hypothetical Hr_0 factors. On this basis, the following chart may be drawn up

Agglutinogens	Known Rh_0 factors	Hypothetical Hr_0 factors
rh rh rh etc	none	Hr_0
λh_0 ϑh_1 ϑh etc	ϑh_0	Hr_0
Rh_0 Rh Rh_1^* Rh_2 etc	Rh_0	$\$r_0$
Rh_0 Rh_1^*	Rh_0	none

It will be seen that the variations of the hypothetical Hr_0 factors are indicated by changing the type face of the capital H in the symbol in conformity with the symbols used for the various Rh_1 factors. As is well known Rh_0 antibodies are readily produced by persons lacking the Rh_0 factor. Until recently no case was known of Rh_0 antibodies being produced by individuals possessing the Rh_0 factor. However, instances of the production of anti Rh_0 by persons having the rare λh_0 variants are now well documented though such cases are quite rare. It is postulated that the difficulty encountered in finding anti Hr_0 up until now has been due to the fact that it has been searched for in individuals homozygous for the ordinary Rh_0 factor, such as persons of type Rh_1Rh_0 . As shown by the chart, however, if the concept proposed is correct such persons have a variant of the Hr_0 factor and therefore would not be apt to respond readily to injections of Hr_0 positive blood. The only individuals who might be expected to form this antibody readily would be those homozygous for a gene determining an agglutinogen having the Rh_0 factor since only such individuals would be truly Hr_0 negative according to the hypothesis. In fact, Buchanan and McIntyre²² found in the serum of his subject an antibody which agglutinated all bloods tested except his own. A similar antibody was found by Gunson and Donahue in their R^*R^* individual and significantly the bloods of the R^*R^* person and the R^*K^* person proved to be compatible. Thus it is likely according to this hypothesis that the antibodies in both instances were of the

specificity anti Hr_0 . Since the anti Hr_0 described here reacts with the blood of all persons except those homozygous for the rare super R^0 gene, it is quite different from the anti d postulated by Fisher and the hypothetical anti Hr_0 of table 30

Factors Rh^A , Rh^B , Rh^C , Wiener and Geiger^{163 164} studied a case of erythroblastosis fetalis in which the mother was Rh_0 positive, and yet her serum contained an antibody giving all of the reactions expected of anti Rh_0 except that it failed to clump her own cells. The reactions of the mother's cells indicated that they did not merely contain an Rh_0 variant, as in the case reported by Argall et al.¹⁶⁵ Moreover, serum from Argall's case clumped the cells of Wiener and Geiger's patient and conversely Rosenfield et al.¹⁶⁶ investigated another case, similar to that of Wiener and Geiger, but again cross matching of the blood specimens proved the antibodies involved to be different. To explain these observations Wiener suggested that the blood of Rh_0 positive individuals have associated with the Rh_0 factor, a series of factors Rh^A , Rh^B , etc. In certain rare individuals, one or more of the associated factors may be missing and such individuals can be sensitized to the missing blood factor. Since "normal" Rh_0 positive blood has all of the factors Rh^A , Rh^B , Rh^C , etc., anti Rh^A serum, for example, is indistinguishable from anti Rh_0 serum except that it fails to agglutinate blood cells of rare individuals lacking the factor Rh^A . If this hypothesis is correct, it would be expected that ordinary anti- Rh_0 serum could contain a mixture of the antibodies anti Rh^A , anti Rh^B , anti Rh^C etc. Indeed it was found that when anti Rh_0 serum was absorbed with cells of type Rh_0 a fraction of the antibodies remained which gave reactions corresponding to anti Rh^A .

To designate those rare Rh positive bloods which lack one of the associated factors Rh^A , Rh^B , an appropriate symbol indicating which factor is absent is used. For example, a type Rh_1^+ individual, like the mother of the erythroblastotic baby studied by Wiener and Geiger, has factor Rh_0 and all its associated factors except Rh^A . Studies¹⁶⁷ on a large series of Caucasoids showed that Rh positive bloods lacking factor Rh^A are extremely rare, but among Negroids as many as one out of 200 Rh-positive persons were found to lack factor Rh^A . If, as seems reasonable, the atypical blood types described are prototypes of a series of blood types yet to be discovered, it be

comes necessary to postulate the existence of additional rare alleles in the Rh Hr series of allelic genes such as R^{0a} , R^{1a} , R^{0b} , R^{1b} etc. Fortunately, these genes, if they exist at all, are extremely rare so that these new discoveries will have little effect on routine Rh Hr typing.

Additional Complexities of the Rh-Hr Types

Just as the Rh_0 factor has its variants, so is there evidence of the existence of rare variants of the other Rh Hr factors. Suitable terminology for designating variants of rh' and rh'' has already been indicated, and similar symbols can be devised for the variants of the other Rh Hr factors. However, knowledge concerning these rare variants is limited, and it is best not to let the symbols get ahead of the facts, or changes in terminology may become necessary later on and lead to confusion.

An antibody has been encountered for the rare blood factor, which like rh'' , occurs only in association with rh' . This antibody¹⁶⁷ may be designated anti rh^* so that at least three sharply defined types of Rh_1 agglutinin exist namely, Rh_1^* , Rh_1^x , and Rh_1 proper the latter being of course the most common. Moreover, a rare factor has been discovered which occurs only in association with factor rh'' . This antibody¹⁶⁸ may be designated anti rh^{*2} to distinguish it from the anti rh^* serum for the factor associated with rh' .

A final complexity may be mentioned and that is capacity of Rh_0 negative individuals sensitized with Rh_0 positive blood lacking the rh' factor, to produce antibodies which react with type rh' blood which lacks the factor Rh_0 . Waller et al¹⁶⁹ obtained one such serum from an immunized donor which reacted with almost equal titer on type Rh_0 and type rh' blood and could be absorbed completely with either. Thus factors rh' and Rh_0 appear to have at least one factor in common, and this has been designated rh^0 by Allen. By testing anti rh^0 serum against the red cells of individuals of type rh Allen encountered one rare blood which reacted with the serum, and this was therefore designated as type rh^0 . Wiener has pointed out the analogy between type rh^0 blood and group A_0 blood, since the latter in ordinary tests reacts like group O blood, but is agglutinated by serum from most group O individuals which contain anti C.

In conclusion it is evident that in addition to the 10 standard

Rh genes r , r' , r'^w , r'' , r^v , R^0 , R^1 , R^{1w} , R^2 , and R^3 , it is now necessary to postulate additional rare alleles r^v , r^o , R^{0v} , R^{0a} , R^{0b} , R^0 , R^0 , R^{1a} , R^{1b} , R^{1c} , R^{2a} , R^{2b} , etc

C-D-E Notations for the Rh-Iir Types

Historical Background. While in this review, Wiener's Rh Iir nomenclature has been used exclusively, those who consult the literature will find that the majority of the articles use the Fisher Race C-D-E notations, either exclusively or in combination with the Rh Iir nomenclature. Therefore, it is necessary to explain the C-D-E notations and the concepts on which they are based.

Fisher's hypothesis as presented by Race¹⁷¹ in 1946 is summarized in table 36. This table was explained by Race as follows:

"Eighteen months ago we reached this stage in agreement with, but independently of Wiener. Wiener lacked the very great help of *St* serum (*Ld* Note: *St* serum is the anti *rh'* serum of Race and Taylor described on page 71). We brought our notation into line with Wiener's. In January 1944, Fisher examined table 37 and noticed that when a gene was positive with *St* it was negative with *rh'*. He supposed that the antigens disclosed by these two sera were due to alleles, and these he called *C* and *c*. To the antibodies he gave corresponding Greek letters, *St* becoming γ , and *rh'*, Γ . He considered that the antigen recognized by anti-*Rh₀* which he called *D* had an allele *d*, and the anti-serum for which was yet to be found. The antigen recognized by anti *rh''* he called *E* and the serum he called *H*. The anti-serum for the hypothetical allele *c* was also to be found.

"In other words, there were three tightly linked loci on the chromosome responsible for *Rh*, making eight possible combinations in a chromosome (table 37). Briefly, Mourant found η , and the anti *Iir* described by Waller and Levine* has the reaction predicted by Fisher for δ ."

Shortly after the original report¹⁷² describing Fisher's hypothesis, Cippell¹⁷³ suggested that instead of using Greek letters for the Rh Iir antibodies the designations anti *C*, anti-*D*, anti-*E*, anti-*c*, anti-*d*, and anti *e* be used. This modification has been adopted by Fisher and Race.

Summarizing, Fisher postulated three pairs of contrasting genes,

* In a footnote, Race adds: "This apparent agreement has since been claimed by Levine to be due to a typographical error in his own paper."

TABLE 36
RACE'S SUMMARY OF FISHER'S HYPOTHESIS

Ant sera	Genes	Antisera
Γ (Anti <i>Rh</i>)	<i>C</i> or <i>c</i>	γ (SI)(Hr)
Δ (Anti <i>Ph</i>)	<i>D</i> or <i>d</i>	δ } then to be found
Π (Anti <i>I</i> h)	<i>F</i> or <i>f</i>	η }

TABLE 37
FISHER'S SYNTHESIS
(Modified after R. R. Race)

Name of serum (Original name)			Ant <i>Rh</i> Ant rh	Ant <i>Hr</i> Ant hr	Ant <i>Ph</i> Ant Rho	Ant <i>Ph</i> Ant rh	Not yet found Ant Hrs	found Ant hrs
Ant body present (Fisher's Cappe)			I Ant C	γ Ant c	Δ Ant D	Π Ant F	δ Ant d	η Ant f
Genes								
Original name	Present name	Fisher						
<i>Rh</i>	<i>R⁺</i>	<i>C D F</i>	+	(-)	(+)	(+)	(-)	(-)
<i>Rh</i>	<i>R</i>	<i>C D c</i>	+	-	+	-	(-)	(+)
<i>Rh₀</i>	<i>r⁺</i>	<i>c D F</i>	+	-	-	+	(+)	(-)
<i>Rh</i>	<i>r</i>	<i>c D c</i>	+	-	-	-	(+)	(+)
<i>Rh₁</i>	<i>R⁺</i>	<i>D F</i>	-	+	+	+	(-)	(-)
<i>Rh₂</i>	<i>R⁺</i>	<i>c D c</i>	-	+	+	-	(-)	(+)
<i>Rh</i>	<i>r</i>	<i>D F</i>	-	+	-	+	(+)	(-)
<i>rh</i>	<i>r</i>	<i>c D c</i>	-	+	-	-	(+)	(+)

which determine corresponding pairs of agglutinogens *C-c*, *D-d* and *I-i* respectively. The correspondence between these symbols and those used by Wiener is shown in table 38. It will be seen that Fisher treats as a gene or agglutininogen what Wiener considers a blood factor or only one of the multiple serological attributes of a given agglutininogen. No clear distinction among gene, blood factor, and agglutininogen is made in the Race-Fisher theory, and the three terms are used interchangeably. According to Fisher, the three gene pairs *C-c*, *D-d* and *I-i* are assumed to be closely linked within the same chromosome. Thus, eight 'chromosomes' are possible corresponding to Wiener's eight standard allelic genes, as shown in table 39.

As has already been pointed out, the concept of linked genes had earlier been considered by Wiener and disproved and discarded (cf. page 63).

TABLE 39

CORRESPONDENCE BETWEEN FISHER'S AND WIENER'S SYMBOLS FOR
ELEMENTARY ANTIGENS

Fisher's Genes or Agglutinogens	Wiener's Blood Factors
<i>C</i>	rh
<i>c</i>	hr
<i>D</i>	Rh ₀
<i>d</i>	(Hr ₀)*
<i>F</i>	rh
<i>e</i>	hr

* Hypothetical. Wiener now uses this symbol in a different sense (cf page 89)

TABLE 3J

COMPARISON OF THE FISHER RACE LINKED GENE HYPOTHESIS AND WIENER'S
MULTIPLE ALLELE THEORY

Fisher Race			Wiener		
Chromosomes		Corresponding antigens	Genes	Corresponding agglutinogens	Blood factors
Fisher's symbols	Short hand symbols*				
<i>c l e</i>	<i>r</i>	<i>c l</i> and <i>e</i>	<i>r</i>	rh	hr and hr
<i>C l e</i>	<i>R</i>	<i>C d</i> and <i>e</i>	<i>r</i>	rh	rh and hr
<i>c d E</i>	<i>R</i>	<i>d</i> and <i>I</i>	<i>r</i>	rh	rh and hr
<i>C d E</i>	<i>R*</i>	<i>C d</i> and <i>E</i>	<i>r*</i>	rh	rh and rh
<i>c D e</i>	<i>R₀</i>	<i>c D</i> and <i>e</i>	<i>R⁰</i>	Rh ₀	Rh, hr and hr
<i>C D e</i>	<i>I</i>	<i>C D</i> and <i>e</i>	<i>R</i>	Rh	Rh rh and hr
<i>c D f</i>	<i>R₂</i>	<i>D</i> and <i>f</i>	<i>R²</i>	Rh ₂	Rh ₀ rh and hr
<i>C D f</i>	<i>R₂</i>	<i>C D</i> and <i>f</i>	<i>R₂</i>	Rh ₂	Rh ₀ rh and rh

* For compactness use is made by Fisher and Race of symbols which they designate shorthand. As Race points out, these are based on Wiener's original symbols. However the same Rh shorthand symbols are used by Fisher and Race for both genotypes and phenotypes.

Actually, there were three predictions implicit in the Fisher hypothesis, namely, that antibodies of specificity anti-*d* and anti-*e* would be found, and that crossing over would be encountered among the components of the tightly linked gene complexes. Of these three predictions only one has been realized, namely, the discovery of anti-*e* (anti hr⁰), as has been described elsewhere (cf page 73). The various claims regarding the discovery of anti-*d* (anti Hr₀) could not

be confirmed, and as Race and Sanger state these may have been examples of anti *f* (anti *hr*). In any event anti-*d* is not available. The anti-*Hr₀* serum described on page 89, is quite different from the anti-*d* serum conceived by Fisher.

Crossing Over. If crossing over could be shown to occur among the postulated pairs of linked genes, as proposed by Fisher, this would be important evidence in favor of his concept. In actual practice, however, this has never been observed.* But, if the genes are assumed to be closely linked, crossing over might occur too rarely to be detected by family studies. Fisher^{175, 176} suggested, in support of his theory, that the rarity of crossing over actually accounted for the low incidence of the rarer genetic combinations in the population. For example, in Caucasoids the less common "chromosomes" *cDe* (R^0), *Cde* (r'), *cdE* (r''), and *CDE* (R^2) were postulated to have arisen from the more common "chromosomes" *CDe* (R^1), *cDE* (R^2), and *cde* (r) by crossing over. For example, by rare crossing over, persons of genotype *CDe/cde* (R^1r) could produce the gametes *Cde* (r') and *cDe* (R^0), similarly, *cdE/cde* (R^2r) individuals could rarely produce gametes *cDe* (R^0) and *cdE* (r''), while persons of genotype *R^1R^2* could produce gametes R^0 and R^2 (cf. fig. 3). Since every time r' , r'' or R^2 could arise by crossing over, R^0 would also be formed, the relationship $R^0 = r' + r'' + R^2$ would be expected to hold. Moreover, "chromosome" *CdE* (r^*) could arise by crossing over mainly from the rare genotypes *Cde/cdE* ($r'r''$) and *CDE/cde* (R^2r) so that the frequency of gene r^* could be expected to be extremely rare according to Fisher's hypothesis. In Mongoloids, where "chromosome" *cde* (r) is virtually absent, the original chromosomes were postulated to have been *CDe* (R^1) and *cDE* (R^2), so that from individuals of genotype R^1R^2 , by rare crossing-over, gametes R and R^0 could be expected to result in equal numbers.

In the original studies of Race et al. on 2,000 Englishmen (cf. table 40), the prediction seemed to be fulfilled, since the estimated frequency of "chromosome" R^0 was 2.57, while the sum of the frequencies of "chromosomes" r' , r'' , and R^2 was $1.24 + 0.73 + 0.22 =$

* In 1950 Lawler,¹⁷⁴ in a study on the heredity of the Rh Hr blood types commented: "No example of crossing over has been found in families tested by this Unit, nor do we know of any certain examples found elsewhere. The work done by Dr. Alexander Wiener's laboratory and this Unit has established with certainty the manner of inheritance of the allelomorphs of Rh."




FIG. 3 Fisher's Cross over Hypothesis

2.19 which is in reasonable agreement with the prediction. In addition the report by Wiener et al. on Mexican Indians (cf. table 40) in which the frequency of the R^0 chromosome was 5.8 per cent and of the R^1 chromosome 3.3 per cent was also considered to be in reasonable agreement with the expectations. Subsequently, the results of additional anthropological investigations became available and a number of these are summarized in table 40. As can be seen the initial satisfactory agreement with the hypothesis has not been maintained in other populations so that as Race and Sanger state in their book in 1954, 'it is possible to pick results strikingly in favour of the crossing over idea—and strikingly against it'.

Fisher and Race have carried the idea further by estimating the distances between the hypothetical linked genes C, D, E , from the ratio of the frequency of each rare gene to that of the genotype from which it was postulated to have arisen. Based on the calculations for

when the same calculations were applied to other populations the relationship no longer held. One of the serious weaknesses of all the calculations is that no allowance was made for agglutinogens characterized by Rh_0 variants or variants of other Rh Hr factors.

Additional evidence considered to be in support of the theory of linked genes and of the arrangement $DCLE$ was the discovery of blood specimens having factor Rh_0 (D) but lacking both members of the pairs Cc ($rh' hr'$) and Ee ($rh'' hr''$). These bloods were designated as $D-/D-$ to indicate that a deletion of the genes Cc and Ee from the chromosome had occurred. The similar example of Gunson and Donahue (cf. page 88) in which the members of the pair $rh'' hr''$ (Ie) were missing has been designated DC^*-DC^*- . Our own explanation for this situation, namely the existence of additional alleles determining agglutinogens with super Rh_0 factors which crowd out the reactivity of other blood factors is more in line with observations in other blood group systems such as the MNS types where rare blood specimens have been encountered which lack the blood factors S s as well as U. In this latter situation postulating the existence of an additional allele has proved a satisfactory solution to the problem without the need for hypothesizing the occurrence of a deletion in a chromosome. Furthermore in *Drosophila* for example large deletions are lethal while small deletions are viable only in heterozygous form. In general deletions produce marked phenotypic modifications yet all individuals encountered to date who are homozygous for the supposed deletions $D-$ and DC^*- have been entirely normal.

The discovery of factor hr (f) made the six factor six gene hypothesis completely untenable. This factor, as previously explained, occurs only in persons bearing the genes R^0 and/or r . The original explanation, namely that this factor was present when c and e were present on the same chromosome but was not found when they were present together but on different chromosomes, the so called position effect, was later changed to the postulation that a fourth locus existed closely related to the CDE loci and occupied by a new gene pair Ff with anti F yet to be found. [It may be pointed out that the postulated serum anti F would be most difficult to recognize since it should give reactions corresponding to the polyvalent serum anti $rh'+rh''$ (anti $C+F$)] As a result, the symbols for the various chromosomes had to be changed, so that for gene r the correspond-

TABLE 40

STATISTICAL TEST OF FISHER RACE CROSS OVER HYPOTHESIS
(After Wiener & S. Trans. A. J. Acad. Sci. 13 199 1951)

Population	Investigator	Number of persons tested	Calculated frequency as (r^2) of genes (chromosomes)							Calculated frequency of genotypes				Chromosome segments		
			r	r^2	R^2	R	R^2	R^2	R^2	R^2	R^2	R^2	R^2	$\frac{CE}{R^2}$	$\frac{DC}{R^2}$	$\frac{DF}{R^2}$
Caucasoids																
U S A	Wiener <i>et al</i>	2 300	36 65	1 23	0 52	3 73	42 70	15 06	05	12 86	31 30	11 04		004	039	045
U S A	Unger <i>et al</i>	7 317	38 3	1 4	0 8	2 8	42 1	15 1	—	12 71	32 25	11 57		0	043	069
England	Race <i>et al</i>	2 000	28 86	0 98	1 19	2 57	42 05	14 11	0 24	11 87	32 68	10 97		020	030	108
Canada	Chown <i>et al</i>	3 100	39 55	1 24	0 73	1 91	43 48	12 87	0 22	11 19	34 39	10 18		020	036	072
Spain	Rue <i>et al</i>	223	36 65	0	0 61	0 61	50 11	12 16	0 45	12 19	3 03	8 99		037	0	051
Czechoslovakia	Rasky <i>et al</i>	181	40 02	0 69	0 69	1 36	39 59	16 00	0 75	13 38	31 68	13 52		056	022	051
Basques	Chalmers <i>et al</i>	383	53 16	1 47	0 25	0 50	37 56	7 07	0	5 31	39 93	7 51		0	037	033
Negroes	Wiener <i>et al</i>	223	28 4	2 7	0	42 1	11 7	14 4	0	3 37	6 65	8 18		0	41	0
N Y C	Wiener <i>et al</i>	200	27 39	0 89	0 89	43 32	17 30	9 02	0	3 10	9 54	4 89		0	003	182
Pigmies, Belgian Congo	Hubinout and Speckh	94	10 5	0	0	63 2	6 2	19 5	0	2 4	1 3	4 1		0	0	0
Sinamese	Phansuboom <i>et al</i>	213	0	0	0	11 13	75 54	11 13	2 16	16 82	0	0		128	—	—
Papuan	Simmons and Graydon	100	0	0	0	2 1	94 3	2 0	1 6	3 8	0	0		42	—	—

Australian abori- gines	234	0	12 57	0	8 54	56 42	20 09	2 08	22 67	0	0	091	—
New Caledonians	325	0	0	0	5 48	33 32	10 77	0 43	17 95	0	0	025	—
Mexican Indians	98	0	0	0	5 8	64 1	26 8	3 3	34 36	0	0	096	—
Navaho Indians	305	0	17 31	2 02	8 37	31 05	35 33	5 73	21 9	0	0	262	—
Simmons and Graydon													
Simmons <i>et al</i>													
Wiener <i>et al</i>													
Boyd and Boyd													

ing chromosome becomes $dcef$ $r = dCcF$ $r'' = dcEF$ $R^0 = Dccf$ etc

To date no position within the four sets of linked genes has been assigned to the gene corresponding to the factor Hr^v or λ discussed on page 81 and similarly no provision has been made in the scheme for factors Rh^A Rh^B Rh^C

With the lack of evidence for crossing over between the various postulated gene pairs Dd Cc Ee and Ff Race has invoked the concept of complete linkage. This latest alteration in the theory makes quite meaningless the concept of linear arrangement since operationally the linked gene theory thus becomes identical with Wiener's theory of multiple alleles making the substitutions $cdef = r$ $Cdef = r'$ etc. The populations in which the relationship $R^0 = R + r + r''$ appears to hold so well strongly suggest that Fisher's suggestion that in Caucasoids the more common genes R R^2 and r may have been the original genes may be correct. However the rarer genes r' r'' R and r probably are not by crossing over but by mutation. If mutation occurred in steps as seems likely this could account for the intermediate genes corresponding to the various Rh Hr variants which are not taken into account by Fisher's original hypothesis. Similarly in Mongoloids the original genes appear to have been R and R^2 in Negroids the original genes appear to have been principally R^0 with some R^1 and R^2 accounting for the relatively high incidence of Rh_0 variants in such populations.

The C-D-F Notations. From the foregoing it is clear that the C-D-F notations differ fundamentally from the Rh Hr nomenclature.¹⁷⁷

1) Implicit in the C-D-F notations is the concept that each Rh Hr agglutinin has but a single blood factor with a single corresponding antibody making it unnecessary to make any distinction between blood factor and agglutinin. For example instead of the agglutinin Rh_1 a set of four agglutinins C D c F is utilized. As has been pointed out however a fundamental principle in immunology is that antigens have multiple corresponding antibodies and thus are characterized by multiple serological factors. This applied not only to the agglutinins of the human blood group but also to the blood groups of cattle fowl mice and even to the antigens of bacteria and viruses and soluble proteins. The Rh Hr nomenclature with its distinctive type faces for agglutinin and blood factors automatically allows for this fundamental serological reality.

2) The C-D E F notations are based on the concept of number of the agglutinogens in uniform sets of three or four if the Ff pair is included. Actually, the number of serological characters which characterize an agglutininogen is almost unlimited because of the complex character of the reacting sites, and thus varies from agglutinin to agglutininogen, as studies on the Rh Hr agglutinogens have shown (cf. table 34). In this respect, the Rh Hr nomenclature is more realistic.

3) The Rh Hr terminology takes into account the existence not only of sharply differentiated blood factors but also of graded series of related blood factors, the latter being indicated by varying the type face, as in the case of the Rh₃ variants. The C-D E notation treats the two cases in the same manner. As a result while the blood factors C*, C' have their corresponding specific antisera anti C and anti C', certain variants of Rh₃ are indicated by the analogous symbol D* even though there is no corresponding anti D* serum.

4) A logical sequel of the concept of a one-to-one correspondence between antigen and antibody is the interpretation of certain serums as containing 'linked' antibodies. For example in the case of the A B O system the anti C antibody had to be designated anti 1+B even though the more reasonable interpretation of the reaction of such anti-serum is that it identifies an additional serological factor of the red cell, namely blood factor C. Similarly in the Rh Hr system linked antibodies anti C+C*, had to be postulated to explain the reactions of anti rh' serum with bloods Rh₁* (DC*el) as well as Rh₁ (DCeF).

5) The C-D E notations pretend to indicate in the symbols all of the reactions of the red cell and are therefore presumed to be self-explanatory. The Rh Hr nomenclature makes no attempt to include all the serological attributes of the agglutininogen in the symbol which instead are in the nature of simple mnemonics as has already been explained. Of interest, are the contrasting effects on the symbol of the discovery of anti f (anti hr) and anti I (anti hr*) serums. While the Rh Hr symbol of gene r remained essentially unchanged the corresponding C-D E symbol *ede* had to be expanded first to *edef* and then to *edefv*. Moreover it has proved impossible to expand the C-D E notations to keep abreast of the newer discoveries such as Rh^A, Rh^B, Rh^C etc. In fact ~~it is impossible to expand the C-D E notations to keep abreast of the newer discoveries~~ where the discoveries have not

certain factors have been included in the notations even though the factors have not yet been proved to exist, as in the case of *d* and *F*¹³

6) A serious difficulty with the C D E F notations is the lack of agreement in the use of these symbols by different workers. This is particularly true of the symbols for the various phenotypes. For example,¹³ the phenotype Rh₁rh has been variously designated as CcDee, CDe cde (+++-), (++-+), C+ c+ D+ E- e+, DCe/-ce DCeF/dcef, C(A+ B+), D(A- B+) E(A- B+) F(A+), etc etc

7) Instead of reporting phenotypes many workers who use the C D E symbols report 'most likely genotypes'. In fact, the phrase 'most likely' is frequently omitted and 'genotypes' are reported based on serological tests alone. This procedure introduces significant statistical errors, and in a number of instances has resulted in confusion. As has already been pointed out the Rh Hr nomenclature has distinctive symbols for phenotypes and genotypes which automatically avoid this pitfall.

The Committee on Medicolegal Problems of the American Medical Association has prepared a report¹⁴ on the medicolegal application of blood grouping tests in which one of the most important problems discussed was that of Rh Hr nomenclature. In the report, the recommendation was made that "unless and until some other convention can be agreed upon the original Rh Hr notations be retained as the standard and sole nomenclature for the preparation of approved medicolegal reports on Rh types." In view of the facts which have been presented here it would appear that this recommendation should now be extended to include the applications of the Rh Hr blood types in clinical medicine and anthropology.

In conclusion it is necessary to reiterate that the essential difference between the two concepts is on a serological and not a genetic level. Blood grouping is a subspecialty of immunology and while a knowledge of genetics is helpful it is only of secondary importance for mastery of the field. The linkage theory completely evades the fundamental immunological concept of the difference between an agglutino-gen and its serological attributes: the blood factors and therein lie the pitfalls of the C-D E notations. Thus, the issue at stake is not merely a choice between two nomenclatures, but the accuracy in reporting scientific observation.

CHAPTER VI

The Kell Blood Groups

In 1946, Coombs et al.¹⁸² encountered an antibody in the serum of the mother of an erythroblastotic baby which gave reactions unrelated to any previously described blood group system. The patient's serum clumped the red blood cells of approximately 7 per cent of all persons tested. The blood factor detected by this antibody was named **Kell** after the patient Wiener and Gordon¹⁸³ independently discovered in the serum of a patient who had had a hemolytic transfusion reaction an antibody which reacted with blood samples of 12.9 per cent of individuals tested. The factor was named **S₁** after the patient and in family studies was found to be inherited as a simple Mendelian dominant, apparently independent of the A B O, M N S P, and Rh Hr blood types. Comparison by Mourant¹⁸⁴ of the two antisera anti **S₁** and anti **Kell** showed them to be identical. For the sake of uniformity, therefore, the symbol **S₁** was discarded by Wiener, and the symbol **K** is now used to designate the **Kell** or **S₁** factor.

Subsequently, other investigators found further examples of this antibody, and the distribution of the **K** factor has been studied in many populations. Among Caucasoids the incidence has been found to vary from a low of about 7 per cent to as high as 13 per cent. Among Negroids the incidence is considerably lower while among Mongoloids the **Kell** factor has not been found.

The most extensive studies have been done among Caucasoids. Of these, one of the largest series is that of Lewis, Chown et al.¹⁸⁵ who observed among 2872 Canadians 198 or 6.89 per cent positive reactors. If we assume that the **Kell** factor is transmitted as a simple Mendelian dominant the gene frequencies for this series can be readily estimated as follows:

$$k = \sqrt{\text{Kell neg}} = \sqrt{0.931} = 0.965$$

$$\text{and } K = 1 - 0.965 = 0.035$$

where **K** is the dominant gene and **k** its allele.

The calculated frequencies of the three possible genotypes for this

population then are

$$\begin{array}{lcl} \text{Genotype } KK & = & (0.035)^2 \approx 0.001225 = 0.12\% \\ \text{Genotype } Kk & = & 2(0.035)(0.965) = 0.675 \\ & = & 6.75\% \end{array} \left. \begin{array}{l} \\ \\ \end{array} \right\} \begin{array}{l} \text{Kell} \\ \text{positives} \\ 6.87\% \end{array}$$

and

$$\text{Genotype } kk = (0.965)^2 \approx 0.9312 = 93.12\%$$

Additional family studies were subsequently carried out particularly by British investigators and Race and Sanger¹⁶ have summarized studies on 336 families with 743 children. Among 507 children resulting from the mating $k- \times k-$ none were observed who were k positive.

Factor k (Cellano)

In 1949 Levine *et al.*¹⁵⁶ found in the serum of a mother of a mildly affected erythroblastotic baby an antibody which when tested against 2,500 different blood samples clumped all but five. This high frequency of positive reactors suggested that the factor it detected might be k determined by the allele of K especially since all negative reactors were **Kell** positive (cf. table 41). The expected incidence of negative reactions in New York and London was about 1 in 500 in close agreement with the frequencies obtained by Levine *et al.* with their antiserum. Therefore the antibody in question originally designated **Cellano** after the patient, is now called anti k . Additional evidence in favor of the theory was obtained by family studies. A particularly striking example is the family described by Levine *et al.* (table 42). Similar families have been described by Shapiro¹⁵⁷ and by Wiener.¹⁵⁸

The series of 2,872 Canadians described above was tested by Lewis *et al.* for factor k as well as factor K . The distribution of the three Kk types observed by them was: type KK , 0.24%, type Kk , 6.63% and kk , 93.11%. From these values the gene frequencies are readily calculated by direct count, as follows:

$$\text{gene } K = KK + \frac{Kk}{2} = 0.24 + \frac{6.63}{2} = 3.57\%$$

$$\text{gene } k = kk + \frac{Kk}{2} = 92.99 + \frac{6.63}{2} = 96.13\%$$

With these gene frequencies, the expected distribution of the three

TABLE 41
THE K & BLOOD TYPES

Tests Done Only for F_a for K		Tests Done for Both Factors K and k			
Phenotypes	React on w th Anti K	Phenotypes	Reactions w th		Genotypes
			Anti K	Anti k	
K (or $K+$)	+	KK	+	-	AA
k (or $K-$)	-	Kk	+	+	AJ
		kk	-	+	kk

TABLE 42
FAMILY SHOWING GENETIC RELATIONSHIP
BETWEEN FACTORS K AND k
(After Levine *et al* ⁴⁶)

Blood of	Reactions w th Serum		Phenotype
	Anti K	Anti k	
Mother	+	+	Kk
Father	+	+	Kk
Children 1	+	+	Kk
2	+	-	KK
3	-	+	kk
4	+	+	Kk
5	+	-	KK
6	+	-	KK
7	+	+	Kk
8	-	+	kk

Kk is type Kk 0.13% type Kk = 6.88% and type kk = 92.99%. Therefore $\chi^2 = 3.28$ and $0.1 < P < 0.5$. The large value of chi square is due mainly to the excess of individuals of the rare type Kk , namely, 7 instead of the expected 3.5 among the 2,674 persons tested. While this may well be accidental, another possibility is that some of the individuals classified as type Kk were actually type Kk but had red cells reacting weakly with anti k serum. The latter interpretation seems to be supported by some recent observations of Allen *et al* now to be described.

Additional $k-k$ Factors

Allen and Lewis⁴⁹ encountered an antibody for a previously unknown blood factor in the serum of an individual who as far as was known had never been exposed to the antigen either through preg-

nancy or transfusion. Subsequently two more examples of serums of identical specificity were found again in donor who had not been sensitized by either transfusion or pregnancy. The new factor was named **Penney** after the first donor and was assigned the symbol **kp***. Among 2 363 individuals in the Boston area 51 (2.16 per cent) were found to have the antigen in approximate gene frequency of 1 per cent.

The **Penney** factor was shown by Allen and Lewis to be related to the **Kk** system by the following observations: (1) Among 73 **Penney** positive individuals only 2 were type **K** while the expected incidence of **K+** individuals was 9.8 per cent. (2) In certain families mother who were apparently of type **kk** had children apparently of type **KK** while some mothers who were typed as **KK** had children apparently of type **kk***. In all such instances the mothers and the children proved to be **Penney** positive so that it became necessary to postulate the presence in the *c* families of a third gene allelic to **K** and **I** which determined the **Penney** antigen. That is, the mother and child in the exceptional instances actually belonged to genotypes **KI^p** and **II^p** respectively or vice versa†. (3) In matings in which one parent had both factors **K** and **kp*** and the other parent lacked both the *c* factors, no child had both factors or lacked both factors; instead half had factor **K** while half had factor **kp***. By using an exceptionally potent anti **k** serum or by the use of the absorption technique it was subsequently found that the **Penney** positive individuals also had the factor **k** or a variant thereof which was being masked by the anti **k** serum used up to that time.

In 1936 Fudenberg¹⁹ encountered in unusual anti serum in a man who had received several transfusions. This serum agglutinated the red blood cells of all but 1 of 2 363 unrelated persons. This single negatively reacting blood was **Kell** negative and **Penney** positive and gave weak reaction with anti **k**. The serum also failed to agglutinate the red cell of one of the patient's two brothers. Like the cases of the negatively reacting blood found in the random series, this blood also gave weak reactions with anti **k**. Exceptionally strong reactions were obtained, however, with both the patient's blood cells and those of his brother when tested with anti **kp*** serum. A one

* In this study only **K+** individuals were tested with anti **k** serum; all **K-** persons were assumed to be type **kk**.

† The new allelic gene is assigned the symbol **k^p** in stead of **K^p** for reasons which will become apparent later.

might expect if these individuals were of genotype $I^P k^P$, and for this reason the symbol Kp^b was assigned to the new factor (also called Rautenberg after the patient) Allen and Lewis suggest that with the four antisera anti K , anti k , anti Kp^a and anti Kp^b , probably 10 genotypes and 9 phenotypes can be distinguished (There is convincing evidence, however, for only 3 alleles, K , k , and k^P , which can yield only 6 genotypes)

No final genetic interpretation is possible at the present time because of the limited facts available. It appears that the agglutinin K determined by gene K has, in addition to factor K factor Kp^b . The gene formerly designated k seems to include at least two varieties. The more common, making up more than 98 per cent, for which the designation gene k may still be retained, determines an agglutinin characterized not only by the factor k but also factor Kp^b . The remainder, which may be assigned the symbol gene k^P , determines a different rare agglutinin having factor Kp^a , and in this case the factor k is poorly defined judging from the weaker reactions with anti k serum. According to a report by Chown,¹²¹ there may be, in addition, a fourth rare allele in this system tentatively designated K^s , which determines an agglutinin that has none of the blood factors K , k , Kp^a or Kp^b .

Serology

The factors K and k are far less antigenic than the Rh_0 factor judging from immunization experiments and clinical observations.¹²²⁻¹²⁴ As a cause of hemolytic transfusion reactions and erythroblastosis fetalis factor K ranks with factor Hr' in importance. Unfortunately, however, the tests for the K k factors are much more delicate than the tests for the Rh Hr factors. Anti K and anti- k are not reactive by the proteolytic enzyme technique, they may react by the saline agglutination or conglutination techniques, but the most satisfactory reactions are obtained with such reagents by the anti globulin method. The clumps in positive reactions are more fragile than the agglutinates of Rh Hr tests, apparently due to the smaller number of reacting sites on the red cell envelope,¹²⁴ and this makes the reactions reproducible with greater difficulty than other serological tests. Reliable results therefore, are possible only by painstaking work carried out by experienced investigators, and these tests have not been sufficiently perfected to be used routinely in medicolegal cases of disputed parentage.

CHAPTER VII

The Lutheran Blood Groups

The first anti serum for a blood factor of the Lutheran system was described in 1946 by Callender and Race¹⁹⁵. The antibody was produced by a patient with lupus erythematosus whose serum contained a number of unusual antibodies as the result of a series of blood transfusions. The factor was named **Lutheran** after the donor whose blood sensitized the patient, and was assigned the symbol Iu^a while the corresponding antibody was designated anti Iu^a . A number of additional anti-serums of specificity anti Iu^a were later encountered by other workers but the reagent is still rare, so that the number of investigations carried out with it is limited.

According to Greenwalt and Sasaki¹⁹⁶ the incidence of the $Iu(a+)$ type in pooled studies of 2 539 Englishmen was calculated by Mourant to be 6.93 per cent. If heredity by a pair of allelic genes Iu^a and Iu^b is postulated the genotype frequencies calculated from these data are Iu^aIu^a 0.12%, Iu^aIu^b 6.81% and Iu^bIu^b 93.07%.

Family Studies. Callender and Race tested the bloods of relatives of 17 $Iu(a+)$ persons and the results indicated that the Iu^a factor was transmitted as a Mendelian dominant. Moreover the Iu^a factor appeared to be inherited independently of the ABO, MN, Rh, Hr, Pp and Kk systems.

Lawler¹⁹⁷ studied the heredity of the Lutheran groups in 47 families with 97 children while Race, Singer and Thompson¹⁹⁸ tested 237 families with 594 children. The results conform with the genetic theory, in particular in 234 families where both parents lacked the blood factor all of 601 children were likewise $Iu(a-)$.

Munwaring and Pickles¹⁹⁹ reported that their anti Iu^a serum distinguished strongly and weakly reacting $Iu(a+)$ cells, analogous to the subgroups of A and postulated three corresponding allelic genes.

Mohr et al.²⁰⁰ tested many large Danish families for the Lewis and Lutheran blood groups and interpreted their results as evidence for linkage between the genes Ic and Iu . However considering the difficulties in carrying out the Lewis tests as described on page 32

it would seem premature to accept the conclusions without confirmation.

Distribution of I_u^a According to Mourant²⁰¹ among European whites different investigators have reported the frequency of the $I_u(a+)$ type to range between 3 and 10 per cent. In a limited number of studies carried out so far on natives of Asia and Australia the $I_u(a+)$ types have been observed in widely varying frequency. In Africa frequencies varying from zero for Bushman to 17 per cent for Nigerians have been reported.

Factor I_u^b Cutbush and Chinarin²⁰² found an antibody in the serum of a woman who had three normal pregnancies but no transfusions which gave the reactions expected for anti I_u . Recently Greenwalt and Sasaki¹⁹⁸ described another example of such a serum in a woman who was sensitized by a blood transfusion. To date no extended anthropological or genetic studies have been carried out with serum of this specificity.

The Lutheran blood groups have not proved to be of any considerable clinical importance up to now, and remain primarily of academic interest.

CHAPTER VIII

The Duffy Blood Groups

In 1950 Cutbush, Molh on and Parkin²⁶³ and van Loghem and Hart²⁶⁴ independently reported the discovery of a blood factor unrelated to the A B O M N-S P p Rh Hr and Kell blood type.

The antibody described by Cutbush et al. was found in the serum of a hemophilic who had sustained a mild transfusion reaction after a series of transfusions received over a period of twenty years. The blood factor was named **Duffy** after the patient. Other examples of sensitization to the **Duffy** factor were encountered by other investigators and one of us (A. S. W.) encountered such a serum with a titer of more than 500 units in a woman who had a serious hemolytic transfusion reaction caused by sensitization to the **Duffy** factor.

Terminology. In order to avoid confusion with the C D T symbols for the Rh Hr types, in stead of the letter *D* the symbol *Fy* was selected by Cutbush et al. to represent the **Duffy** factor. Wiener²⁶⁵ has suggested shortening this symbol to **F** and this latter terminology will be used in the present discussion.

Serology. All the anti **F** serums encountered up to the present time have been usable only by the anti globulin technique. Obviously, therefore, when the red cells are already coated, as in cases of auto hemolytic anemia or in erythroblastosis fetalis, the **Duffy** tests can not be carried out reliably. The clumps which characterize a positive reaction may be large, but they are quite friable, even more so than the clumps of the Kell tests. This Wiener²⁶⁶ ascribes to the smaller number of antigenic sites on the red cell envelope, which he estimates to be about one tenth as numerous as the antigenic sites of the Rh Hr agglutinogens. Because of the delicate nature of the reactions, the results are duplicated with difficulty, and it is only with painstaking work that reliable results are possible.

Distribution. Among Caucasoids the frequency of the **F** positive individuals ranges between 60 and 70 per cent, among Negroids the frequency is considerably lower, namely, between 10 and 25 per cent, while in Mongoloids the incidence is highest, in some populations approaching 100 per cent.

by
 (a) 100 - 58.72 = 41.28 type *f* From these values the gene frequencies are readily calculated as follows

$$f = \sqrt{\text{type } f} = \sqrt{0.4148} = 59.72\%$$

$$F = 100 - f = 41.28\%$$

From these values, the frequencies of the three theoretically possible genotypes can be readily calculated

$$\begin{aligned} \text{Genotype } FF &= (0.4128)^2 = 17.04\% \\ \text{Genotype } Ff &= 2(0.4128)(0.5872) = 48.48\% \\ \text{Genotype } ff &= (0.5872)^2 = 34.48\% \end{aligned} \left. \vphantom{\begin{aligned} \text{Genotype } FF \\ \text{Genotype } Ff \\ \text{Genotype } ff \end{aligned}} \right\} \text{type } F = 65.52\%$$

Heredity. The few family studies carried out to date support the hypothesis that the **Duffy factor** is transmitted as a simple Mendelian dominant. In the combined studies by Cutbush and Mollison,²⁹⁹ and Rice and Sanger³⁰⁰ there are 34 families in which both parents were type *f*, and among the 78 children of these families none were type *F*.

Factor *f* Titration

A corresponding serum specimen of genotypes *ff* and *Ff*, but not with those of genotype *FF*. In 1951, Ikem et al.³⁰¹ detected an antibody, during a routine postnatal examination, which in tests on blood specimens from 59 persons gave the reactions expected for anti *f*. Unfortunately, further studies were prevented because the serum lost its activity. In more recent studies, R. Sanger, has found that among Negroes blood lacking both factor *F* and *f* are not uncommon so that the situation is more complex than it seemed at first.

CHAPTER IX

The Kidd Blood Groups

This blood group system was first described in 1951 by Allen Diamond and Niedziela²¹⁹ who found the new antibody in the serum of the mother of an erythroblastotic baby. The blood factor detected by the antibody was named *Kidd* after the patient. It was found to be independent of the ABO MNS Pp Rh Hr Kell Duffy and Lutheran systems.

Nomenclature In order to avoid confusion with the symbol for the Kell factor the Kidd factor has been assigned the symbol *Jk* by Allen et al. Wiener²²⁰ has suggested shortening the symbol to *J* which is the symbol used in this discussion.

Distribution According to a summary compiled by Mourant²¹ among Caucasoids the incidence of factor *J* is approximately 75 per cent. Among Negroids the incidence has been found to be above 90 per cent while among Mongoloids the frequency has ranged from 30 to 100 per cent. The largest series is that of Roenfield et al.²²¹ who found 76.72 per cent type *J* and 23.28 per cent type *j* among New York whites. From these figures the theoretical gene frequencies are readily determined as follows:

$$J = \sqrt{\text{type } J} = \sqrt{0.7672} = 48.25\%$$

$$j = 100 - 48.25 = 51.75\%$$

And from these estimated gene frequencies the distribution of the three theoretically possible genotypes can be calculated:

$$\left. \begin{aligned} \text{Genotype } JJ &= (0.4825)^2 &= 23.28\% \\ \text{Genotype } Jj &= 2(0.4825)(0.5175) &= 49.91\% \\ \text{Genotype } jj &= (0.5175)^2 &= 26.78\% \end{aligned} \right\} \text{Type } J = 76.72\%$$

Factor *j* If the gene *j* gives rise to a corresponding agglutinin *j* the blood factor *j* should be present in the blood cells of all individuals of genotype *Jj* and *jj*. According to Race and Singer, at least three anti-serums have been encountered giving reactions expected for anti-*j*.

Serology The anti-*J* serums described in the literature, as well as

those encountered by one of us (A S W), have been usable only by the anti globulin technique. Therefore, these tests have the same limitations as the Duffy tests. Under ideal conditions, the reactions are clear cut and readily visible to the naked eye, but the agglutinates are friable like those obtained with Duffy antiserums. Another difficulty is the tendency of the antiserums to lose their activity upon storage. Therefore, reliable results are obtained only by carefully repeating the tests several times, using the blind technique, and including controls of known types J and j.

CHAPTER IX

The Kidd Blood Groups

This blood group system was first described in 1951 by Allen Diamond and Niedziela¹⁰ who found the new antibody in the serum of the mother of an erythroblastotic baby. The blood factor detected by the antibody was named Kidd after the patient. It was found to be independent of the ABO MN-S Pp Rh Hr Kell Duffy and Lutheran systems.

Nomenclature In order to avoid confusion with the symbol for the Kell factor the Kidd factor has been assigned the symbol *Jk* by Allen et al. Wiener¹⁰ has suggested shortening the symbol to *J* which is the symbol used in this discussion.

Distribution According to a summary compiled by Mourant²⁰ among Caucasoid the incidence of factor *J* is approximately 75 per cent. Among Negroid the incidence has been found to be above 90 per cent while among Mongoloids the frequency has ranged from 50 to 100 per cent. The largest series is that of Rosenfield et al.²¹ who found 76.72 per cent type *J* and 23.28 per cent type *j* among New York whites. From these figures the theoretical gene frequencies are readily determined as follows:

$$J = \sqrt{\text{type } J} = \sqrt{0.7672} = 48.25\%$$

$$j = 100 - 48.25 = 51.75\%$$

And from these estimated gene frequencies the distribution of the three theoretically possible genotypes can be calculated:

$$\begin{array}{lcl} \text{Genotype } JJ & = (0.4825)^2 & = 26.78\% \\ \text{Genotype } Jj & = 2(0.4825)(0.5175) & = 49.94\% \\ \text{Genotype } jj & = (0.5175)^2 & = 23.28\% \end{array} \left. \vphantom{\begin{array}{l} JJ \\ Jj \\ jj \end{array}} \right\} \text{Type } J = 76.72\%$$

Factor *j* If the gene *j* gives rise to a corresponding agglutminogen *J* the blood factor *j* should be present in the blood cells of all individual of genotype *Jj* and *jj*. According to Race and Sanger at least three anti-serums have been encountered giving reactions expected for anti *J*.

Serology. The anti *J* serums described in the literature as well as

parents were Diego negative, and in none of the 111 children did the factor appear.

High Frequency Blood Factors. As examples of blood factors with a high frequency in the general population may be mentioned the factor U of the M-N S system, factor H of the A-B O system, factor *hr*⁺ of the Rh Hr system, *T*₁ of the P system, and factors *k* and *k*^p of the Kell system. In addition to these, at least three other high frequency blood factors are known, namely, Vel, I, and Yt⁺ (cf. table I). Of these three factors, factor I has been studied most intensively and is of the greatest interest.

Anti I was found²¹³ in the serum of a group A₁B woman with acquired hemolytic anemia, whose serum also contained cold autoagglutinins in high titer. When this patient had hemolytic reactions following transfusions, her serum was also shown to contain an isohemagglutinin of high titer, anti I, reactive at room temperature with almost all other human bloods but not with the patient's own cells. No other member of the patient's family was found to be I negative (type 1). Among a series of 22,964 random donors only 5 proved to be I negative, and of these 4 were Negroes. Among the positively reacting blood specimens marked differences in titer were obtained, ranging from 25 to 5000 units, thus indicating the existence of variants of the I agglutinin, namely, I₁, I₂, I₃, etc. Recently, Shapiro²¹² has encountered a second example of anti I in South Africa, although of lower titer. In this case both the propositus and her child proved to be I negative.

TABLE 43
SOME LOW FREQUENCY BLOOD FACTORS

Name of Blood Factor	Observers	Random Series		Family of Proposita	
		+	-	+	-
Levy	Callender and Race (1916)	0	320	3	4
"	Wiener (1942)	0	ca 500	1	3
Joblin	Gillies (1947)	0	120	2	2
Becker	Libel and Prokop (1951)	0	272	3	4
Berrens	Davidsohn et al. (1953)	0	418	4	3
Cavaliere*	Wiener and Brancato (1953)	0	48	4	3

* F. H. Allen has recently shown that this blood factor is probably identical with the Wright factor (Wt⁺), described at about the same time by Holman (*Lancet* 2: 115, 1953).

Low Frequency Blood Factors As examples of the low frequency factors may be mentioned **Mi** and **Gr** of the **M N S** system **rh⁺** and **rh⁻** of the **Rh Hr** system **super Rh₀** of the **Rh Hr** system and **Kp^a** of the **Kell** system. In addition to the **e** a large number of blood factors have been reported generally detected with the serum of mothers of erythroblastotic babies. Due to the low frequency of these factors the genetic investigations have been limited to the families of the propositi. The findings in each are compatible with heredity as a simple Mendelian dominant. The more fully described of these blood factors are listed in table 43.

As an example of a low frequency factor may be mentioned the case report of Wiener and Brancato.² A woman who had babies with erythroblastosis caused by **Rh₀** sensitization then had an erythroblastotic baby who proved to be **Rh₀** negative. Tests of the maternal serum demonstrated that in addition to anti **Rh₀** there was another antibody present which was designated anti **Ca** after the patient. Among 7 **Rh₀** negative relatives the blood of 4 proved to have the factor **Ca** while 48 consecutive **Rh₀** negative donors were all **Ca** negative. Another striking example is that of Davidson *et al.*³ who found a potent antibody for a blood factor present in 4 out of 7 members of their patient's family but absent from 448 consecutive blood donors (cf. table 43).

CHAPTER XI

Medicolegal Applications

The clear cut hereditary transmission of the human blood group factors has led to their application for the solution of medicolegal problems of disputed parentage. The types of cases in which the tests have been applied are as follows:

- 1) A man accused of the paternity of a child born out of wedlock denies the charge. This is the most common type of case with which courts are confronted where blood grouping tests are used.
- 2) A man is accused of assault or rape, and tries to prove innocence by showing that he is not the father of the child born to the complainant.
- 3) A child is born in lawful wedlock, but the husband denies paternity.
- 4) It is suspected that two newborn infants have been interchanged in a nursery in a hospital.
- 5) A woman may simulate pregnancy and childbirth, and may claim that a certain child is hers to compel a man to marry her or to obtain dower in a dead husband's estate.
- 6) An individual who wished to immigrate to the U.S. asserts that his parents are American citizens living in the states, but it is suspected that the claim is fraudulent.

A number of reports^{1,2,3,4} regarding the medicolegal applications of blood grouping tests have been published by the Committee on Medicolegal Problems of the American Medical Association, and they should be consulted by those interested in this special problem. The legal aspects of the subject are discussed in detail in the excellent book by S. B. Schatkin.⁵

Blood tests can be used only to exclude and not to prove paternity. In certain jurisdictions laws have been passed empowering the courts to order blood grouping tests in cases where parentage is an issue. Naturally, the chances of disproving paternity will depend upon how complete an examination is carried out. However, the examination should be restricted to only those blood tests which have been proved to be reliable, and where an adequate number of heredity studies have been carried out. Therefore, the Committee on Medicolegal

Problems of the American Medical Association has recommended that the routine tests for court cases be limited to the A and B factors of the A B O system, the M and N factors of the M N system, and the factors Rh₀, rh', rh'', and hr' of the Rh Hr system. In private cases, or under special circumstances, qualified medicolegal experts may find it desirable to carry out more complete examinations, which will vary depending on the experience of the investigator and on the reagents available to him.

For convenience of reference, simplified tables which show the application of the A-B O, M N, and Rh-Hr tests for excluding paternity have been prepared (cf. tables 44, 45, 46).

Chances of Excluding Paternity

If a man falsely accused of paternity is found to have a blood group genetically incompatible with that of the child this establishes his

TABLE 44
INCLUSION OF PATERNITY BY THE A B O GROUPS*

Phenotype of the Putative Mother	Phenotype of Putative Father			
	O	A	B	AB
O	A B AB	B AB	A, AB	O AB
A	B AB	B AB	None	O
B	A AB	None	A AB	O
AB	O AB	O	O	O

* Find the phenotypes of the putative father and mother at the top and side columns of the tables and locate the box at which these intersect. In the box are given the groups not possible in children of the mating groups given in bold face type represent children for whom maternity is excluded.

TABLE 45
INCLUSION OF PATERNITY BY THE M N TYPES*

Phenotype of Putative Mother	Phenotype of Putative Father		
	M	N	MN
M	MN N	M N	N
N	M, N	M, MN	M
MN	N	M	None

* Find the phenotypes of the putative father and mother at the top and side columns of the table, and locate the box at which these intersect. In the box are given the types not possible in children of the mating types given in bold face type represent children for whom maternity is excluded.

innocence. If the blood groups satisfy the laws of heredity this does not necessarily prove that he is the father because of the limited number of types known and the possibility of coincidence. The chances of exonerating a falsely accused man will vary depending on the distribution of the blood groups in the population as well as on the blood group of the accused man. For example in certain tribes of American Indians where all of the members belong to group O the A B O groups would be worthless for solving such problems.

Wiener^{2, 6, 27} has derived general formulae for the chances of excluding paternity by the A B O blood groups. These are given below and those interested in their derivation should consult the original paper.

Blood group of falsely accused man	Chance of excluding paternity
Group O	$r^2(p + q) + 2pq(1 + r)$ (42)
Group A	$q(1 + r)^2$ (43)
Group B	$p(1 + r)^2$ (44)
Group AB	r^2 (45)
Group indeterminate	$p(1 + r)^2 + q(1 + r)^2 + pqr^2(p + q) + 2pqr^2$ (46)

In the above formulae p , q and r represent the frequencies of genes I^A , I^B and I respectively. The maximum chance of excluding paternity is found to be approximately 20 per cent and occurs when the gene frequencies are $p = 0.2213$, $q = 0.2213$ and $r = 0.5574$.

Similarly formulae have been derived for the chances of excluding paternity using the M N types as follows:

Blood type of falsely accused man	Chance of excluding paternity
Type M	$r(1 - r)$ (47)
Type N	$r(1 - r)$ (48)
Type MN	0 (49)
Type indeterminate	$r(1 - r)$ (50)

In the above formulae m and n represent the frequencies of gene I^M and I^N , respectively. Here the maximum chance of excluding paternity is 18.75 per cent and occurs when the gene frequencies are $m = 0.5$ and $n = 0.5$.

When the formulae are applied to populations such as that encountered in New York City table 4 could be drawn up. In making the calculations account must be taken of the independent inheritance

of the A B O and M N types. If P_1 is the chance of excluding paternity by one blood group system and P_2 is the chance of excluding paternity by a second independent system, then the chances of exclusion when both are tested for is given by the formula

$$P = 1 - (1 - P_1)(1 - P_2) \quad (51)$$

This formula can be extended to include any number of blood group systems. It is of interest that the combined chances of excluding paternity by the combined use of the A B O and M N tests in New York City is about 30 per cent, which is close to the theoretical maximum of 35 per cent.

General formulae for the chances of excluding paternity by the means of the Rh Hr types have not been derived. These would be expected to be exceedingly complex. Wiener^{22, 23} has estimated that the chances of excluding a falsely accused man of undetermined Rh Hr type is close to 25 per cent, when tests are done for the factors Rh, rh', rh'' and hr'. Thus, when tests are done for the factors mentioned of all of the three systems, the chances of excluding a falsely accused man is about 50 per cent. In practice the rate of exclusion is considerably less than 50 per cent, since most of the defendants in paternity proceedings will be the actual fathers. Mr S. B. Schitkin and Mr Sol Cooperman of the Corporation Council's Office of New York City have observed over a period of many years that approximately 17 or 18 men out of every 100 men who deny paternity are actually excluded by the tests, indicating that approximately 35 per cent of such accusations are false. In problems of disputed paternity involving married couples as in divorce actions our experience has been that the exclusion rate is considerably higher.

In selected cases it may be felt desirable to carry out additional tests, for example for the subgroups of A secretor type P, S, hr'', Kell, Duffy and other factors for which the expert has reliable anti-serums available.

The formula for the chance of excluding paternity, using only a single blood factor D, transmitted as a Mendelian dominant, is given by the formulae below.

Blood type of falsely accused man	Chance of exclusion	
D+	0	(51)
D-	dr^2	(52)
Type undetermined	dr^4	(53)

Here d represents the frequency of the dominant gene, and r the frequency of the recessive gene. The maximum chance of excluding paternity is 0.08192 or about 8.2 per cent, and occurs when the gene frequencies are $d = 0.2$ and $r = 0.8$, respectively. In practice, it turns out that the distribution of most of the blood factors inherited in this way is such that the chance of excluding paternity is less than the maximal theoretical value. Moreover, because of the overlapping among tests and the occurrence of double, triple or quadruple exclusions it becomes progressively more difficult to increase the chances of excluding paternity by the introduction of new blood tests. For example, a blood factor with an 8 per cent chance of exclusion will add only 4 per cent to the chances of exclusion since one half of the innocent men will already have been excluded by the A B O M N and/or Rh Hr tests. It has been estimated that all of the new blood tests combined add only about 12 to 15 per cent to the chances of excluding parentage, and it is questionable whether it is practicable or worth while to carry out these difficult tests in such cases, since the results may not prove acceptable to the courts.

Boyd²³⁰ has calculated in detail the chances of excluding paternity by the Rh Hr tests in a Caucasoid population. His results, including the distribution of the Rh Hr types, for the population to which his figures apply are given in table 48. For example, a falsely accused man of type rh has an almost 45 per cent chance of being excluded by tests for the Rh Hr types. By combining the figures for tables 47 and 48, it is possible to calculate the chances of excluding paternity for a falsely accused man of any A B O group, M N type, and Rh Hr type. As an example, the chances for a group AB, type M, type Rh₁Rh₂ man would be $1 - (1 - 0.607)(1 - 0.4418) = 0.782$ or 78.2 per cent.

TABLE 47
CHANCES OF PROVING NON PATERNITY WITH THE
AGGLUTINOGENS A B M AND N*
(From Wiener's *Blood Groups and Transfusion*)

Group	O			A			B			AB			Unknown
Type	M	N	MN	M	N	MN	M	N	MN	M	N	MN	
Chances (Per cent)	59.0	54.6	23.5	39.6	45.1	7.7	41.1	49.3	14.6	60.7	63.4	39.9	31.9

* Based on the frequencies of the blood types in New York City

TABLE 48
CHANCES OF EXCLUDING PATERNITY BY THE Rh Hr TYPES
(Modified from Boyd, *Amer J Human Genet* 7 229 1955)

Phenotype of Accused Man*	Frequency in Caucasoids (per cent)	Chance of Exclusion (per cent)
rh	14.36	14.95
rh rh	0.62	18.45
rh'rh	0.01	51.62
rh''	1.31	36.28
rh rh''	0.03	9.73
Rh ₀	2.41	44.14
Rh ₁ rh	35.77	10.66
Rh ₁ Rh ₁	19.73	44.48
Rh ₂	12.66	33.55
Rh ₁ Rh ₂	12.99	0
Rh ₂ Rh ₁	0.11	41.75
Unknown		25.0

* Not included in the table is the rare phenotype rh₃rh

Disputed Maternity

The blood tests may also be used to refute false claims of maternity. For example, maternity would be excluded if any of the following blood type combinations is encountered:

- 1) Mother group AB—child group O
Mother group O—child group AB
- 2) Mother type M—child type N
Mother type N—child type M
- 3) Mother type rh'rh', rh₃rh', Rh₁Rh₁, or Rh₂Rh₁—child type rh, rh'', Rh₀, or Rh₂
Mother type rh rh'', Rh₀, or Rh₁—child type rh'rh', rh₃rh', Rh₁Rh₁, or Rh₂Rh₁

The chances of excluding maternity are readily calculated as follows. For the A B O blood groups = $2(\bar{O})(\overline{AB}) = 2(0.40)(0.05)$ or approximately 4 per cent. For the M N types = $2(\bar{N})(\bar{M}) = 2(0.30)(0.20)$ or approximately 12 per cent. For the Rh Hr types = $2(\overline{rh'rh'} + \overline{Rh_1Rh_1} + \overline{Rh_2Rh_1})(\overline{rh} + \overline{rh''} + \overline{Rh_0} + \overline{Rh_2}) = 2(0.20)(0.30)$ or approximately 12 per cent. For all three systems combined = $1 - (1 - 0.04)(1 - 0.12)(1 - 0.12) = 0.258$ or approximately 26 per cent.

Situations not infrequently arise where a man falsely accused of

paternity wishes to have blood tests performed on himself and the child but the mother of the child is not available for one reason or another. In such cases the same principles apply as in cases of disputed maternity, that is the chances of excluding paternity are approximately 26 per cent or about half of the chances when all three of the parties are available.

More Complex Problems

At times a married man correctly suspects that he is not the father of more than one of his wife's children. In such cases the blood tests may exclude paternity for one or more of the children. Obviously the chances that paternity will be excluded for at least one of the children will be greater than 50 per cent. Such evidence may be and has been used as grounds for divorce. If monozygotic twins are involved the chances are not increased because such twins are invariably of identical blood groups.

In immigration cases both of the putative parents may be excluded in cases of fraudulent applications. The chances that the falsity of the claim will be disclosed by excluding at least one of the parents will necessarily be considerably greater than 50 per cent.

An unusual but interesting problem sometimes arises when it is suspected that two newborn babies have been interchanged in a hospital. In such cases the blood tests are of considerable aid because with two sets of parents and two children available the chances of obtaining conclusive results are very high. For example the chances of solving a problem of interchange of babies for a single blood factor has been shown by Wiener²¹ to be as follows:

$$1_D = 2\bar{D}\bar{R} \quad (51)$$

where 1_D is the chances of solving the problem, \bar{D} is the frequency of the blood factor in population and \bar{R} is the frequency of persons lacking the factor. The maximum chance for a single blood factor is as high as 29.03 per cent and occurs when the frequency of phenotype D is equal to 0.33 and \bar{R} equals 0.67.

Wiener has estimated the chances of detecting interchange of
 and by
 blood
 only 64

per cent. If, in addition, the Rh Hr tests are done, about 90 per cent of these problems can be solved.

Circumstantial Evidence of Paternity

As has already been pointed out, the blood tests cannot be used to prove paternity, only to exclude it, because of the possibility of coincidence of the blood types of a falsely accused man and the actual father. There are rare circumstances, however, when both the accused man and the child have a blood factor of very low incidence in the population, or both have a rare combination of blood factors. Such evidence, while of little testimonial value, may help to convince the accused man of his paternity, so that he may then be willing to accept the child as his own. As examples may be mentioned the occurrence in both the putative father and the child of the rare types rh' or rh'' . Other infrequent types sometimes encountered in both the putative father and child are Rh_i'' and K.

On the other hand, the chances of excluding paternity in cases where a false accusation has been made may be enhanced by the use of such factors as hr and rh'' . For example, a man of type $Rh_i Rh_i$ cannot have a child of type rh or Rh_i unless he is the carrier of the genes r or R^0 (genotypes R^2r , R^1R^0 or Rr'). If tests on his red cells with anti hr serum give a negative reaction, then he cannot be a carrier of either of these genes and paternity is excluded. Moreover, since factor rh'' is always associated with factor rh' in the same agglutminogen, an individual of type $Rh_i'' rh$ cannot have a child of type $Rh_i Rh_i$, nor can an individual of type $Rh_i Rh_i$ have a child of type $Rh_i'' rh$.

It should be mentioned that the existence of "super" Rh genes constitute a pitfall in the application of the Rh Hr types in disputed paternity cases. For example, a putative father belonging to type $Rh_i Rh_i$ while the child belongs to type Rh_0 , or the reverse, is not excluded if the man and child are both carriers of gene \bar{R} . Fortunately gene \bar{R} betrays its presence by the high agglutinability of the red cells by anti Rh_0 serums, as has already been explained (cf. page 87).

Case Reports

The application of blood grouping tests in actual practice is best illustrated by citing a few cases (cf. Wiener,²² Alvarez,^{23a} Unger,²⁴ Sussman and Schutkin²⁵ and Witchky and Wexler²⁶).

Case 1. This case, which was tried before the Court of Special Sessions in New York City, is unusual because a man was accused of the paternity of a

children born out of wedlock. When he denied the charge, blood grouping tests were carried out, upon order of the court, on the defendant and on the complainant and her six children with the results shown in table 49. As can be seen, the complainant belongs to group O, three of her children to subgroup A_1 , and three to subgroup A_2 . The defendant belongs to subgroup A_1 , corresponding to which there are three possible genotypes according to the genetic theory. If we assume that the defendant is of genotype $I^{A_1}I^{A_1}$, half his children with a group O mother would be expected to be subgroup A_1 and half subgroup A_2 , as actually occurred in this case.

As far as the M-N types are concerned, the complainant is type N and the defendant type M, and all six children are type MN, as would necessarily result assuming the defendant to be their father. Moreover, the complainant is type Rh_{hrh} and the defendant type Rh_{hrh} , three of the children type Rh_{hrh} , and three of type Rh_{hrh} , again in agreement with the theoretic expectations.

The excellent fit of the blood group findings of the six children with the genetic expectations strongly suggests that the defendant actually is their father, as charged. While blood tests that fail to exclude paternity are inconclusive because of the possibility of coincidence, the results here are so striking that they helped convince the defendant of his paternity, so that he contributed willingly to the children's support in conformity with the decision of the court.

The use here of tests for the subgroups of A is inconsistent with the opinion already cited that these tests are not sufficiently reliable for routine use in medicolegal cases. However, subgrouping tests are so easy to perform and the results so informative that the senior author carries them out regularly. At times the findings are illuminating, as in the present case. When tests for the subgroups of A indicate an exclusion of paternity but all other tests are inconclusive, the safest practice is to omit the subgrouping results from the official report, and to present them instead in a covering letter, explaining in detail the limitations of the test, for the guidance of the court.

Case 2. A man accused of the paternity of a child denied the charge. As shown in table 50, the blood tests indeed showed that he was not the father of the child. In fact, the results in this case were unusual in that the tests provided triple proof of nonpaternity.

Case 3. Four young Chinese adults came to the United States claiming de

TABLE 49
RESULTS OF BLOOD GROUPING TESTS IN A CASE OF DISPUTED PATERNITY INVOLVING SIX CHILDREN

Blood Group System	Putative Father	Mother	Children					
			1st	2nd	3rd	4th	5th	6th
A B O	Phenotypes Genotypes $A_1 I^A I^A$, $I^A I^O$, or $I^A I^{A_2}$	O $I^O I^O$	$A_2 I^A I^O$	$A_1 I^A I^O$	$A_2 I^A I^O$	$A_1 I^A I^O$	$A_1 I^A I^O$	$A_1 I^A I^O$
M N	Phenotypes Genotypes M $L^M L^M$	N $L^N L^N$	MN $L^M L^N$	MN $L^M L^N$	MN $L^M L^N$	MN $L^M L^N$	MN $L^M L^N$	MN $L^M L^N$
Rh Hr	Phenotypes Genotypes $Rh Rh_1$ $R^+ R^+$ or $R^{+} r'$	$Rh_1 h$ $R^+ R^{+} R^0$, or $R^{+} r'$	$Rh_1 rh$ $R^+ r$, $R^+ R^0$ or $R^{+} r'$	$Rh_1 Rh_1$ $R^+ R^+$ or $R^{+} r'$	$Rh_1 Rh_1$ $R^+ R^+$ or $R^{+} r'$	$Rh_1 h$ $R^+ r$, $R^+ R^0$ or $R^{+} r'$	$Rh_1 h$ $R^+ r$, $R^+ R^0$ or $R^{+} r'$	$Rh_1 Rh_1$ $R^+ R^+$ or $R^{+} r'$

Thus the claims of derivative citizenship of at least three of the four applicants were proved to be fraudulent.

Nomenclature

The current controversy regarding Rh Hr nomenclature has already been discussed in detail (cf. page 100). This important question has been the subject of study by a special Subcommittee on Blood Grouping Tests of the Committee on Medicolegal Problems of the American Medical Association. As is pointed out in their report²³ 'The submission to jurists by different experts of medicolegal reports with conflicting symbolisms cannot help but confuse them and shake their confidence in the blood tests. Moreover reports in two different symbolisms require translation from one into the other when comparisons are to be made, there is always danger of error in the process. Therefore the adoption of one uniform nomenclature for medicolegal reports is most desirable.

From a study of the available evidence the committee concluded in part as follows: 'Although corresponding to the two systems of designating the Rh Hr types there are two conflicting genetic theories the crux of the problem is actually on a serologic level. It is most important to distinguish clearly between agglutinogens and their serologic attributes—the blood factors. The C D I notations for the Rh Hr types make no allowance for the difference between a blood factor and an agglutininogen and incorporate the tacit incorrect assumption that every agglutininogen has but a single corresponding antibody. This has led to a number of paradoxes. Indeed much of the evidence that has accumulated regarding the serology and genetics of the Rh Hr types refutes the interpretation implicit in the C D I notations. On the other hand the original Rh Hr nomenclature presents the data objectively without committing the user to a preconceived interpretation. The committee, therefore recommends that the C D I notations for the Rh Hr types be discarded and that, in approved medicolegal reports unless and until some other convention is agreed upon the original Rh Hr nomenclature be retained as the sole nomenclature for this blood group system.'

Recently the recommendation that the Rh Hr nomenclature be used exclusively for medicolegal reports has been approved by the Section on Immunology of the American Academy of Forensic Sciences. The time has now come to extend this recommendation to

the applications of the Rh Hr types in clinical medicine and anthropology. Confusion will remain until editors of scientific journals require that all articles dealing with the Rh Hr types conform with this recommendation as has been done in the case of the analogous recommendation made in a report²³⁴ published in 1937, regarding the nomenclature of the A B O blood groups.

Qualifications of Experts

The field of immunohematology has grown in scope and complexity and there are relatively few workers who have devoted the time and effort required to become expert in the field. As a guide for lawyers and courts desiring to use blood tests as aid in cases of disputed parentage, the Law Department of the American Medical Association has available a list of qualified blood group specialists. A similar panel of qualified workers is used by the Court of Special Sessions in New York City, from which lawyers may select an expert to carry out blood tests in cases of dispute of paternity. However failure of many jurisdictions to exert equal care in the selection of experts to carry out blood grouping examinations has been responsible for a number of unfortunate mistakes, and this is a problem which still requires a satisfactory solution.

In blood transfusion practice, it is considered *a priori* evidence of negligence if a patient of group O is given group A blood, and if the patient dies, the physician, hospital and/or blood bank may be subjected to costly lawsuits. Similarly, a mistake in blood grouping

TABLE 50
TRIPLE EXCLUSION BY BLOOD TESTS OF MAN
ALLEGEDLY ACCUSED OF PATERNITY

Blood Group System		Blood of		
		Putative father	Mother	Child
A B O	Phenotypes Genotypes	$I^A I^A$ or $I^A I^O$	$I^O I^O$	$I^B I^O$
M N	Phenotypes Genotypes	$I^N L^N$	$L^N L^N$	$L^M L^N$
Rh Hr	Phenotypes Genotypes	Rh ₊ Rh ₊ $R^+ R^+$ or $R^+ r$	rh ⁺ $r^+ r$	rh rr

TABLE 51
CASE ILLUSTRATING THE APPLICATION OF BLOOD TESTS TO DISPROVE FALSE CLAIMS OF DERIVATIVE CITIZENSHIP

Blood Group system	Total	Father	Putative Mother	Child			
				1	2nd	3d	4th
ABO	Genotypes	$I^A I^A$ $I^A I^O$ or $I^A I^O$	AB $I^A I^B$	O $I^O I^O$	AB $I^A I^B$	A $I^A I^A$ $I^A I^A$ or $I^A I^O$	O $I^O I^O$
MN	Genotypes	MN $L^M L^N$	M $I^M I^M$	MN $L^M L^N$	M $L^M L^M$	MN $L^M L^N$	M $L^M L^M$
Rh	Genotypes	Rh R ₁ R R ₂ R ₃ R ₄ R ₅ R ₁ R ₂ R ₃ R ₄ R ₅ R ₁ R ₂ R ₃ R ₄ R ₅ R ₁ R ₂ R ₃ R ₄ R ₅	Rh rh R R R ₂ or R ₂ R ₃	Rh R ₁ R R or R r	Rh R ₁ R R or R r	Rh Rh R ₂ R ₂ R ₂ r or R r	Rh rh R r R R ₂ or R ₂ r

in a medicolegal case of disputed paternity erroneously excluding paternity or maternity could conceivably lead to a lawsuit for malpractice or for damages for defamation of character. An error in medicolegal cases is even less defensible than in clinical cases because the claim that the tests had to be done under the pressure of an emergency cannot be used. The medicolegal expert has plenty of time in which to carry out the tests as many times as desired and with as many controls as necessary. When in doubt he can draw new samples and repeat the tests and if he wishes he can send the blood samples to an independent expert for confirmation. Unfortunately however the least experienced workers whose reports are most in need of verification are generally the least sensitive to the possibility that their findings may be in error.

A fully qualified expert in blood grouping is not merely a careful technician but an individual thoroughly versed in the fundamentals of blood group serology and genetics. Such an individual prefers accuracy to simplicity and has no difficulty in using the recommended Rh Hr nomenclature which merely translates the results obtained into symbols. Such a worker has no difficulty in preparing a clear self-explanatory report when presenting his evidence to a lay bench and jury. Wiener²²⁵ maintains therefore that with certain outstanding exceptions those who find it necessary to resort to the incorrect C D E notations for the sake of simplicity do not understand the subject thoroughly and he suggests that this may be used as one of the criteria by which to judge the qualifications of an expert.

CHAPTER XII

Anthropological Studies on the Blood Types

Until recently the use of blood grouping tests in racial studies attracted relatively little attention among physical anthropologists. Thus most journals and text books on physical anthropology concentrate a good deal of attention on skeletal measurements and external physical characteristics but devote little space to blood grouping. This has been due principally to the limited value of the findings until the Rh Hr blood types were discovered. For example when tests for the A B O groups alone are carried out closely related ethnic groups may exhibit significant differences in distribution while totally unrelated peoples may have similar distributions. The situation was improved somewhat by the introduction of the subgroups of A and the M N types but the greatest advances resulted from the discovery of the Rh Hr types.

At first the studies were restricted to the Rh factor alone because the only serums available for the earlier studies were the animal antirhesus serums or human serums of specificity anti Rh₀. It was soon observed that while the Rh₀ factor is present among approximately 85 per cent of Caucasoids it occurs more frequently among Negroids and among practically 100 per cent of Mongoloids.²⁶ The results were quite striking and became more so when the tests were extended to include the rh and rh₁ factors. For example Negroids were characterized by a high frequency of type Rh₀ a type of blood which occurs infrequently among Caucasoids and Mongoloids.²⁷ The introduction of tests for the factors hr and hr₁ did not alter the situation to any great degree because as has already been pointed out the reactions given by anti Hr serums can largely be predicted in advance from the reactions of the anti Rh serums. The principal value of Hr serums in anthropological investigations is as an aid in recognizing individuals carrying the rare gene R⁺. In this way it was possible to demonstrate that gene R⁺ which is rare among Caucasoids and has not been demonstrated to date among Negroids occurs in as many as 3 to 6 per cent of Mongoloids.²⁸

Based on the newer findings concerning the distribution of the blood group factors, especially Rh Hr Wiener has proposed a semi-qualitative classification of human races as follows:

- I Caucasoid group—highest frequency of gene r gene I^{A2} present as well as gene I^{A1} gene L^M slightly more frequent than gene L^N
- II Negroid group—highest frequency of gene R^0 and Rh 0 variants gene I^{A2} present as well as gene I^{A1} also frequent & intermediates gene L^M slightly more frequent than gene L^N
- III Mongoloid group—highest frequency of the rare gene R genes r and I^{A2} absent or rare
 - a Asiatic subdivision—genes L^M and L^N almost equal in frequency
 - b Pacific subdivision (including Australian aborigines and Ainu)—low frequency of gene L^M and high frequency of gene L^N
 - c American subdivision (including Amerindians and Eskimos)—high frequency of gene L^M and low frequency of gene L^N

Before the use of blood tests was introduced anthropologists classified races on the basis of their external physical characteristics or skeletal measurements. This led to a number of manifest contradictions and paradoxes. One reason for this is that the external physical characteristics are subject to conscious selection while conscious selection would hardly play a role for the blood groups. Moreover whereas physical characteristic such as skin color texture of the hair height and shape of the nose have considerable adaptive value the possession or lack of any particular blood factor has no special advantage except in relation to isosensitization in pregnancy. Based on their physical characteristics Australian aborigines were considered to be intermediate between Negroids and Caucasoids while Papuans were placed in the Negroid group. Studies on the Rh Hr types, on the other hand, clearly demonstrated that these peoples belong to the Mongoloid group, a conclusion which is more reasonable judging from geographical and historical considerations.

Study of the distribution of the blood group factors, especially the Rh Hr types, is of great value in analyzing the results of crosses between different ethnic groups. For example the distribution of the Rh types among Puerto Ricans (cf. table 25) clearly indicates that this ethnic group arose from a cross between Negroids and Caucasoids. However, one should be cautious before drawing conclusions from

the results of blood tests alone. For example, in the case of Puerto Ricans the presence of shovel-shaped incisors indicates that Mongoloids—that is, American Indians—also participated in the cross from which the Puerto Ricans originated, as had been pointed out by Washburn.

That the A B O groups and Rh Hr types are not entirely without selective value has been demonstrated by the discovery of their role in isosensitization by pregnancy. For example, Rh-negative mothers (genotype rrh) who have become sensitized to the Rh_0 factor will have a certain number of Rh-positive babies (genotype Rhr) who are stillborn or die during the neonatal period from erythroblastosis. As Wiener³⁹ pointed out, if we assume that we are dealing with a population of constant size containing x Rh genes and y rh genes, then the initial distribution of the genes would be as follows:

$$Rh = \frac{x}{x+y} \quad \text{and} \quad rh = \frac{y}{x+y}$$

If the number of fetuses and newborn that die from erythroblastosis during one generation is c , then the distribution of the genes during the second generation would be

$$Rh = \frac{x-c}{x+y-2c} \quad \text{and} \quad rh = \frac{y-c}{x+y-2c}$$

If to begin with the number of Rh genes is equal to the number of rh genes, this process would have no effect on the relative distribution of the genes. If, on the other hand, the incidence of the two genes is initially unequal, the less frequent gene would be affected to a greater extent than the more common gene, so that eventually, other things being equal, over a period of thousands of generations, the incidence of the less common gene would be substantially reduced or it might be practically eliminated.

This process could explain the virtual absence of the Rh_0 -negative type among Mongoloids, but it then becomes difficult to account for the relatively high frequency of gene rh among Caucasoids. Wiener³⁹ and Haldane⁴⁰ independently pointed out that if one assumed the existence of a population in the past, and possibly still existing at the present time, consisting largely of Rh-negative persons, a hybrid population could result with a serological composition resembling that of Caucasoids. This hypothesis has received support

from the findings in Basques by Etcheverry²⁴¹ and confirmed by Mourant. Among these people who differ from other Europeans in their isolated habitat, unique language and social customs, the frequency of gene *rh* is extraordinarily high, namely about 60 per cent. It had been previously shown by the Hirszfelds²⁴² that the incidence of gene *I^B* decreases from east to west in Eurasia. According to Candela²⁴³ gene *I^B* was introduced at the time of the Mongolian invasion; the degree of crossing between European and Mongoloids being indicated by the frequency of gene *I^B*. Here again the Basques are unique in that the frequency of gene *I^B* is the lowest in Europe. For these reasons it seems probable that the Basques represent most closely the original European stock which must have had a very high frequency of gene *rh* as well as a very low frequency of gene *I^B*.

Analysis of Mixed Populations

A number of methods have been suggested for analyzing blood group frequencies of mixed populations. The principle of the calculations is the same as for computing the results of mixing solutions of different concentrations. Here we shall describe a simple method which has been used by Mourant.²⁴⁴ This method has the advantage that it can be applied to many blood factors at the same time and thus give a comprehensive view.

Let *p* represent the frequency of the gene for a given blood factor in a population A and let *q* represent the frequency of the same gene in a population B. Then suppose a new population C is formed with *x* individuals from population A and *y* individuals from population B. If *r* represents the frequency of the gene in the mixed population, this is obviously the weighted mean of *p* and *q* and is readily calculated by the following formula:

$$r = \frac{px + qy}{x + y} \quad (22)$$

The value of *r* can also be derived geometrically. The frequency *p* for population A is erected on one vertical line and the frequency *q* for population B on another vertical line. The interval between the two vertical lines A and B is divided in inverse proportion to *x* and *y* and a new vertical line C erected. A straight line is then drawn connecting the points *p* and *q* and the point at which it intersects line C represents the frequency of the gene in the mixed population.

Conversely if it is known that a certain mixed population C resulted from a cross between two other populations A and B the proportion of individuals from populations A and B which entered the cross can be determined if the frequencies of the gene for any agglutino-gen is known for all three populations. One first marks the frequency p for population A on one vertical line and the frequency q for population B on a second vertical line and a straight line is drawn between the two points. A second line is then drawn parallel with the horizontal axis at distance r from it. At the point of intersection of the two points a third line is erected and this represents the composition of the population C .

An example of the practical application of the method is given in figure 4 from the study of Chalmers et al.⁴⁵ In constructing this diagram the frequencies of gene r were used as the basis. A straight line was drawn connecting the points representing the frequencies of gene r in Basques on the one hand and Siamese on the other. Then the horizontal lines were drawn representing the frequencies of gene r in England, Latvia and India and through points of intersection obtained vertical lines were then erected for the three populations. On the five vertical lines points representing the fre-




FIG. 4.—Analysis of Blood Group Gene Frequencies of Mixed Populations (After Chalmers et al.)

quencies of genes R^1 , R^2 , R^0 , I^A and L^M were marked and the corresponding points on the five lines connected producing the results shown in the figure. If the populations in England, Latvia and India had been produced as a result of crossing between Basques and Siamese one would expect the lines representing each of the gene frequencies to be straight. As a matter of fact the lines are not all straight which is to be expected since the populations in question are not merely such simple mixtures.

The purpose of this chapter has been to present the broad principles involved in the application of blood groups in anthropology. No attempt has been made to tabulate the numerous studies that have been reported. Readers who have occasion to consult the original sources should bear in mind that some of the data in the literature suffer from errors in blood typing and this possibility should always be borne in mind especially when the findings appear to be bizarre or unique. The first compendium of the literature was prepared by Boyd³⁴⁶ and included principally the earlier studies on the A B O groups. In Wiener's book⁴⁷ a less complete compilation was given but this source also lists findings on the subgroups of A and the M N types. The most comprehensive and most valuable source is the recent book by Mourant²⁴⁸ which includes the voluminous data especially on the Rh Hr blood types and also on other blood group systems including P, Kell, Duffy, Kidd and Lutheran. One of the most important investigators in this field is R. T. Simmons^{254, 255} whose numerous valuable contributions should be studied and used as models by those interested in the subject.

REFERENCES

- 1 a LANDSTEINER, K *The Specificity of Serological Reactions*, 2nd rev ed , 1945, Harvard University Press, Cambridge, Mass
- b WIENER A S *Blood Groups and Transfusion* 3rd ed , 1943, C C Thomas, Springfield, Ill
- c WIENER, A S *Rh Hr Blood Types, Applications in Clinical and Legal Medicine and Anthropolgy* 1954, Grune & Stratton, New York
- d WIENER, A S *Rh Hr Syllabus* 1954 Grune & Stratton, New York, 1955, Georg Thieme, Stuttgart, 1955, V Idelson, Naples, 1956, Prensa Med Mexicana, Mexico, 1957, Tokyo (Japanese)
- e RACE R R , AND SANGER, R *Blood Groups in Man* 3rd ed , 1958, Blackwell Scientific Publ , Oxford
- f SCHIFF, F , AND BOYD, W C *Blood Grouping Technic* 1942, Interscience Publishers, Inc , New York, N Y
- 2 LANDSTEINER K *Zentralbl f Bakteriologie* , 27: 357 (1900), *Wien Klin Woch* , 14: 1132 (1901)
- 3 LANDSTEINER, K , AND LEVINE, P *Proc Soc Exp Biol and Med* , 24 600, 941 (1927)
- 4 LANDSTEINER, K , AND WIENER, A S *Proc Soc Exp Biol and Med* , 43: 223 (1940)
- 5 WIENER, A S , AND SONN GORDON, F B *Amer J Clin Path* 17 67 (1947)
- 6 a RENKONEN, K O *Ann Med Exp Biol, Fenn* , 26 66 (1948)
- b KRÜPE, M *Blutgruppenspezifische Pflanzliche Fixeisskörper*, 1956, Ferdinand Enke, Stuttgart
- 7 FISHER, R A *Statistical Methods for Research Workers*, 6th ed , 1936, Oliver & Boyd, Edinburgh
- 8 a WIENER, A S , AND WEXLER, I B *Bact Rev* , 16 69 (1952)
- b WIENER A S , AND WEXLER, I B L GEDDA's *Notant' Annali delle Leggi Mendeliane*, 1956, pp 147-162, Istituto Gregorio Mendel, Rome
- c WIENER, A S AND KAROWE, H *J Immunol* , 49 51 (1944)
- 9 WIENER, A S *J Immunol* , 34: 87 (1938)
- 10 FRIEDSTEIN, A A , AND OTTENBERG, R *Trans A Y Pathol Soc* , 8. 157 (1908)
- 11 VON DUNGERN, E , AND HIRSZFELD, L *Ztschr f Immunitäts* , 6 281 (1910)
- 12 BERNSTEIN, F *Ztschr f indukt Abstamm u Vererbungslehre*, 37. 237 (1925)
- 13 KIRIHARA, S , AND HAKURINSAI, K *Nagoya J Med Sci* , 2: 75 (1932)
- 14 FURUHATA, T *Japan Med World*, 7: 197 (1927)
- 15 LANDSTEINER, K , AND LEVINE, P *Proc Soc Exp Biol and Med* , 24: 911 (1927)
- 16 THOMSEN, O , FRIDFELDERICH, V , AND WORSAAE, F *Acta Path et Microbiol Scand* , 7. 157 (1930)

- 17 LANDSTEINER, K, AND LEVINE, P *J Exp Med*, 48 731 (1928)
 (1947)
 ed 18800,
 153 1444 (1953)
 22 GREENWALT, T, SASAKI, T, SANGER, R, AND RACE, R R *Proc Nat Acad Sci*, 40 1126 (1956)
 23 SHAPIRO, M *J Forensic Med (S Africa)*, 3 152 (1956)
 24 LANDSTEINER, K AND WIENER A S *J Exp Med*, 74 309 (1911)
 25 WIENER, A S *Science*, 98 182 (1943)
 26 WIENER, A S *Proc Soc Exp Biol and Med*, 54 316 (1943)
 27 MURRAY, J, RACE R R AND TAYLOR G L *Nature (Lond)* 183 360 (1911)
 28 WIENER, A S *Nature (Lond)* 2 735 (1948)
 29 WIENER, A S *Science*, 100 595 (1944)
 b KEMPTHORNE, O *An Introduction to Genetic Statistics*, 1957, John Wiley & Sons, Inc, N Y
 35 HIRSZFELD, L AND HIRSZFELD, H *Lancet* 2 675 (1919)
 36 BERNSTEIN, I *Ztschr f indukt Abstamm u Vererbungsl* 86 233 (1930)
 ing, 1932, Julius Springer, Berlin
 40 LANDSTEINER, K, AND LEVINE, P *J Immunol* 12 441 (1926)
 41
 42
 43
 44
 45
 (1910)
 46 GROVE RASMUSSEN, M, SOUTTER, L, AND LEVINE, P *Am J Clin Path* 22 1157 (1952)
 47 DUNSFORD, I, AND ASPINALL, P *Ann Eugen (Lond)* 17 30 (1952)
 48 UNGER, L J, AND WIENER, A S *J Lab & Clin Med* 44 387 (1954)
 49 SCHIFF, F, AND ADELSBERGER, L *Ztschr Immunitäts*, 40 235 (1924)
 50 FRIEDENREICH, V, AND WITZ, S *Ztschr Immunitäts*, 78 152 (1933)
 51 OWEN, R D *J Immunol*, 73 29 (1954)
 52 LEHR, H *Ztschr Immunitäts*, 66 175 (1930)
 53 PLTAONEN, T *Acta Med Fenn 'Duodecim'* 14 Part 2 (1930)
 54 WIENER, A S AND DELBEN, R R *J Immunol*, 47 467 (1943)

- 55 SCHIFF F AND SASAKI Z *Ztschr Immunitäts* 77 129 (1932) *Klin Woch*, 34 1426 (1932)
- 56 UYETAMA R *Jap J Med Sci* 3 13 (1940)
- 57 MURANT A E *Nature* (Lond) 158 237 (1946)
- 58 ANDRESEN P H *Acta Path et Microbiol Scand* 24 616 (1948)
- 59 ANDRESEN P H *Acta Path et Microbiol Scand* 25 728 (1948)
- 60 GRUBB R *Nature* (Lond) 162 933 (1948)
- 61 MORGAN W T J *Proceedings 11th International Congress of Blood Transfusion* pp 285-290 1955 ANDERSON L F and MORGAN W T J *Biochem J* 50 462 (1952)
- 62 RACE R R SANGER R LAWLER S D AND BERTINSHAW D *Brit J Exp Path* 30 73 (1949)
- 63 ANDRESEN P H ANDRESEN A JORDAL K AND HENNINGSEN L *Rev d Hematol* 5 305 (1950)
- 64 SNEATH J S AND SNEATH P H A *Nature* (Lond) 176 172 (1955)
- 65 NICHOLAS J W JENKINS W L AND MARSH W I *Brit Med J* 1 1458 (1957)
- 66 GRUBB R AND MORGAN W T J *Brit J Exp Path* 30 195 (1949)
- 67 WIENER A S *Laf Digest* (St Louis) Vol 18 No 5 November 1954
- 68 CEPELLINI R personal communication
- 69 LANDSTEINER K AND MILLER C P *J Exp Med* 43 835 (1925)
- 70 WIENER A S unpublished observations
- 71 WIENER A S CANDELA P B AND GOSS I J *J Immunol* 45 229 (1942)
- 72 CANDELA P B WIENER A S AND GOSS L J *Zoologica* 25 513 (1940)
- 73 WIENER A S AND GORDON I B *Brit J Hematol* 2 305 (1956)
- 74 WEINER W LEWIS B MOORES P SANGER R AND RACE R R *Vox Sang* 2 25 (1957)
- 75 YOKOTAMA M STACY S M AND DUNFORD I *Vox Sang* 2 348 (1957)
- 76 BHENDE Y M DESHPANDE C K BHATIA H M SANGER R RACE R R MORGAN W T J AND WATKINS W *Lancet* 1 903 (1952)
- 77 LEVINE P ROBINSON I CYRANO M BRICCS O AND TALKINBURG I *Blood* 10 1100 (1955)
- 78 LANDSTEINER K AND LEVINE P *J Exp Med* 47 757 (1928)
- 79 WIENER A S ZINSSER R AND SEIKOWE J *J Immunol* 27 431 (1934)
- 80 LANDSTEINER K AND LEVINE P *J Exp Med* 48 731 (1928)
- 81 WIENER A S *J Immunol* 21 157 (1931) *Human Biology* 7 222 (1935)
- 82 CROMB W *Deutsche Ztschr f d ges gerichtl Med* 24 167 (1935)
- 83 FRIEDENREICH V *Deutsche Ztschr f d ges gerichtl Med* 25 338 (1936)
- 84 WIENER A S *J Immunol*, 34 57 (1938)
- 85 LANDSTEINER K, AND WIENER A S *J Immunol*, 33 19 (1937)
- 86 WALSH R J AND MONTGOMERY C *Nature* (Lond) 160 504 (1947)
- 87 SANGER R RACE R R, WALSH R J AND MONTGOMERY, C *Heredity* 2 131 (1948)
- 88 WIENER A S DiDIEGO, N, AND SOKOL, S *Acta Med Genet et Gemellol*, 2 391 (1953)

- 89 WIENER A S *Amer J Human Genet* 4 463 (1952)
- 90 LE JAE P KUBSICHEL A B WIGOD M AND KOCH F *Proc Soc Exp Biol and Med* 78 218 (1951)
- 91 WIENER A S *Acta Med Genet et Gemellol* 3 314 (1954)
- 92 WIENER A S *Acta Med Genet et Gemellol* 6 95 (1957)
- 93 FISHER R A Cited after RICE R R and SINGER R *Blood Groups in Man* 2nd ed p 62 1954
- 94 RACE R R AND SANGER R *Blood Groups in Man* 2nd ed 1954
- 95 WIENER A S UNGER I J AND GORDON I B *J Amer Med Assoc* 153 1441 (1953)
- 96 WIENER A S UNGER I J AND CHEN L S *ence* 119 31 (1954)
- 97 GREENWALT T et al *Proc Natl Acad Sci* 40 1197 (1954)
- 98 LAND TEINER K STRUTTON W R AND CLARKE M W *J Immunol* 27 469 (1954)
- 99 CHALMERS J N M IAIN J W AND MOURANT A P *Brit Med J* 2 15 (1953)
- 100 SHAPIR M *J Fœtal Med (S Afr)* 3 152 (1954)
- 101 VAN DER HART M B SSIAN H AN VAN IGHET J J *Var Sang* 4 108 (1954)
- 102 IEVINE I et al *Proc Soc Exp Biol and Med* 77 402 (1951)
- 103 WALLACE J MINE G R M N J MORFIS I SANGER R AND RACE R R *Nature (London)* 179 478 (1957)
- 104 ALLEN R CORCORAN P A KENYON H B AND BRIDGES N in press
- 105 LANDSTEINER K AND LEVINE I *J Immunol* 17 1 (1950)
- 106 WIENER A S AND IEFERS H R *Ann Int Med* 13 2308 (1940) Wiener A S *Amer J Clin Path* 12 302 (1952)
- 107 FURUKATA T AND ISABURA I *Jap J Genetics (Japanese)* 12 20 (1956)
- 108 WIENER A S *J Immunol* 66 287 (1951)
- 109 CAMFORD C I AND STAFFORD J M *Nature (London)* 179 147 (1957)
- 110 IEVINE P ORBITT O WALLACE R K AND IEFERS H R *Proc Soc Exp Biol and Med* 77 43 (1951)
- 111 SANGER R *Proc 6th Cong Internat Soc of Blood Transfusion* 1 p 110 113 SINGER R N Y 1953
- 112 LANDSTEINER K AND WIENER A S *Proc Soc Exp Biol and Med* 43 23 (1940)
- 113 WIENER A S *Proc Soc Exp Biol and Med* 70 56 (1950)
- 114 WIENER A S *Proc Soc Exp Biol and Med* 74 309 (1951)
- 115 56 171 (1951)
- 116 24 122 (1951)
- 117 DAVENPORT K AND ABELSON N M *J Clin Invest* 30 672 (1951)
- 118 WIENER A S *J Lab and Clin Med* 31 821 (1945)
- 119 DAVENPORT K AND DENTON R I *J Lab and Clin Med* 31 821 (1945)
- 120 CAMFORD C I AND RACE R R *Nature (London)* 2 15 (1951)

- 121 MORTON, J A , AND PICKLES, M M *Nature* (London), 159 779 (1947)
- 122 WIENER, A S , AND LANDSTEINER, K *Proc Soc Exp Biol and Med* , 53: 167 (1943)
- 123 WIENER, A S *Science*, 98 182 (1943), WIENER, A S and SONN, E B *J. Immunol* , 47 461 (1943)
- 124 WIENER, A S *Proc Soc Exp Biol and Med* , 54: 316 (1943)
- 125 WIENER, A S , SONN, E B , AND BELKIN, R B *Proc Soc Exp Biol and Med* , 54 238 (1943)
- 126 WIENER, A S , SONN, E B , AND BELKIN, R B *J Exp Med* , 56 171 (1944)
- 127 WIENER, A S , AND HYMAN, M A *Amer J Clin Path* , 18 921 (1948)
- 128 WIENER, A S *Nature* (London), 2 735 (1948)
- 129 Cited by LEVINE, P *J Ped* , 23 656 (1943)
- 130 RACE, R R , AND TAYLOR, G L *Nature* (London), 152 300 (1943)
- 131 WIENER, A S , DAVIDSON, I , AND POTTER, E L *J Exp Med* , 81: 63 (1945)
- 132 FISHER, R A , cited after RACE, R R *Nature* (London), 153 771 (1944)
- 133 MOURANT, A E , *Nature* (London), 155 544 (1945)
- 134 WIFNER, A S , AND PETERS, H R *Amer J Clin Path* , 18: 533 (1948)
- 135 WIENER, A S *Science*, 102 479 (1945)
- 136 WIENER, A S , DAVIDSON, I , AND POTTER, E L *J Exp Med* , 81: 63 (1945)
- 137 WIENER, A S , SONN, E B , AND POLIVKA, H *Proc Soc Exp Biol and Med* , 61: 382 (1946)
- 138 WIENER, A S , AND SONN GORDON, E B *J Immunol* , 57: 203 (1947)
- 139 WIENER, A S *Proc Eighth Internat Cong Genetics, Hereditas*, Suppl Vol, pp 500-519 (1949)
- 140 RACE, R R , TAYLOR, G L , IKIN, I W , AND PRIOR, A W *Ann Eugen* (London), 12 206 (1944)
- 141 RACE, R R , TAYLOR, G L , IKIN, I W , AND DORSON, A M *Ann Eugen* (London), 12. 261 (1945)
- 142 MCFARLANE, M *Ann Eugen* (London), 13: 15 (1946)
- 143 LEVINE, P personal communication
- 144 MCGEE, R P , LEVINE, P , AND CEFANO, M *Science*, 125: 1043 (1957)
- 145 WIFNER, A S , AND BRANCATO, G J *J Lab and Clin Med* , 40: 27 (1952)
- 146 CALLENDER, S T , AND RACE, R R *Ann Eugen* (London), 13 102 (1946)
- 147 WIENER, A S , BRANCATO, G J , AND GORDON, E B *Acta Genet Med et Gemellol* , 6 195 (1957)
- 148 LAWLER, S D , BERTINSHAW, D , SANCER, R , AND RACE, R R *Ann Eugen* (London), 15: 258 (1950)
- 149 WIENER, A S , GORDON, E B , AND HANDMAN, L *Amer J Human Genet* , 1: 127 (1949)
- 150 ROSENFELD, R F , VOGLF, P , GIBBEL, N , SANGFR, R , AND RACE, R R . *Brit Med J* , 1: 975 (1953)

- 153 DeNATALE A CAHAN A JACK J A RACE R R AND SANGER R
J Amer Med Assoc 159 247 (1955)
- 154 WIENER A S *Science* 100 595 (1944)
- 155 WIENER A S UNGER L J AND SONN L B *Proc Soc Exp Biol and Med* 58 89 (1945)
- 156 ROSENFELD R L VOGL P MILLER E B AND HABER G *Blood* 6 1123 (1951)
- 157
- 158 RACE R R SANGER R AND SELWYN J G *Brit J Exp Path* 32 124 (1951)
- 159 STURGEON P *J Immunol* 68 277 (1952)
- 160 GUYSE
- 161 BUCH
- 162 WIENE
- 163 GEIGE
- 164
- 165 ARGALL C L BALL M AND TRENTLMAN F *J Lab and Clin Med* 41 895 (1953)
- 166 ROSENFELD R F HABER G GIBBEL N *Proc 8th Congress of the International Society of Blood Transfusion* pp 90-95 S Harger New York 1958
- 167 STRATTON F AND RENTON P H *Brit Med J* 1 962 (1954)
- 168 GREENWALT T SANGER R AND RACE R R cited after Race and Sanger *Blood Groups in Man* 2nd ed 1954
- 169 WALLER R K AND WALLER M *J Lab and Clin Med* 34 770 (1949)
- 170 ALLEN F H *10th Ann Meeting Amer Assoc of Blood Banks* Chicago Nov 4 1957
- 171 RACE R R *Ann N Y Acad Sci Vol XLVI Art 3 pp 388-390* (Nov 8) 1946
- 172 FISHER R A cited by Race R R *Nature (Lond)* 153 771 (1944)
- 173 CAPPELL D F *Brit Med J* 2 601 (1946)
- 174 LAWLER S D BERTINSHAW D SANGER R AND RACE R R *Ann Eugen (London)* 15 208 (1950)
- 175 FISHER R A AND RACE R R *Nature (London)* 157 48 (1946)
- 176 FISHER R A *Proc Scient* 35 95 (1947)
- 177 WIENER A S *Proc Soc Exp Biol and Med* 2 224 (1955)
- 178 DODD I
- 179 STORMO
- 180 WIENER
- 181 WIENER *J Amer Med*
- 182 COMBS R R A MOURANT A F AND RACE R R *Lancet* 1 761 (1946)
- 183 WIENER A S AND SONN G ROON L B *Lee d'Hématol* 2 3 (1957)

- 184 MOURANT A F personal communication
- 185 LEWIS M CHOWN B AND PETERSON R F *Amer J Phys Anthropol* **13** 323 (1955)
- 186 LEVINE P WIGOD W BACKER A M AND PONDER R *Blood* **7** 869 (1949)
- 187 SHAPIRO M S *Amer Med J* **26** 931 (1952)
- 188 WILNER A S BRANCATO A J AND WAITMAN R *J Lab and Clin Med* **42** 570 (1953)
- 189 ALLEN I AND LEWIS S J *For Surg* **2** 81 (1957)
- 190 FULDENBERG H cited by Allen and Lewis
- 191 CHOWN B cited by Allen I personal communication
- 192 RACE R R *Bull Central Lab & Biochem* **2** 191 (1952)
- 193 WIENER A S SAMWICK A A MORRISON H AND CHOWN I *Exp Med and Surg* **13** 347 (1955)
- 194 WILNER A S AND GORDON I B *Amer J Clin Path* **23** 429 (1958)
- 195 CATTENDER S T AND RACE R R *Ann Eugen (London)* **13** 102 (1946)
- 196 GREENWAIT T J AND SASAKI T *Blood* **12** 998 (1957)
- 197 LAWLER S D *Ann Eugen (London)* **15** 255 (1950)
- 198 RACE R R SANGER R AND THOMSON J S cited by Race & Sanger *Blood Groups in Man* 2nd ed
- 199 MAINWARING H R AND PICKLES M M *J Clin Path* **1** 202 (1948)
- 200 MOHR J *A Study of Linkage in Man* 1954 Ejnar Munksgaard Copenhagen
- 201 MOURANT A F *The Distribution of the Human Blood Groups* 1951 Blackwell Sci Publ Oxford
- 202 CUTBUSH M AND CHANARIN I *Nature (London)* **178** 855 (1956)
- 203 CUTBUSH M AND MOLLISON P I *Heredity* **4** 383 (1949)
- 204 VAN LOUHEM J J VAN DER HART M *Nederl Tijdschr & Gercoek* **11** 745 (1950)
- 205 WIENER A S *Amer J Clin Path* **23** 987 (1953)
- 206 WIENER A S AND SOHN I B *Amer J Clin Path* **23** 708 (1953)
- 207 IKIN I W MOURANT A F AND PIAUT G *Brit Med J* **1** 581 (1950)
- 208 RACE R R AND SANGER R *Heredity* **6** 111 (1952)
- 209 IKIN I W MOURANT A F PETTENKOFER H J AND BLUMENTHAL G *Nature (London)* **168** 1077 (1951)
- 210 ALLEN I H DIAMOND I K NIEDZIELA B *Nature (London)* **167** 482 (1951)
- 211 ROSENFIELD R I AND VOEL P GIBBEL N ORINO G AND HABER G *Amer J Clin Path* **23** 1222 (1953)
- 212 LEVINE P KOCH I A MCGEE R T AND HILL G H *Amer J Clin Path* **24** 292 (1954)
- 213 LEVINE P ROBINSON I A LAYRISSSE M, ARENDS T AND DOMINICUEZ SISCO R *Nature (London)* **177** 40 (1956)
- 214 LAYRISSSE M ARENDS T AND DOMINICUEZ SISCO R *Acta Med Venet* **3** 172 (1955)
- 215 LAYRISSSE M AND ARENDS T *Science* **123** 623 (1957)

- 216 LEWIS, M, KAITA, H, AND CHOW, B *Amer J Human Genet* 9 274 (1957)
- 217 LAYRISSE, M, AND ARENDS, T *Blood*, 12 115 (1957)
- 218 WIENER A S, UNGER L J, COHEN, I, AND FELDMAN, J *Ann Int Med*, 44: 221 (1956)
- 219 SHAPIRO, M personal communication
- 220 WIENER, A S, AND BRANCATO, G J *Amer J Human Genet*, 5 350 (1953)
- 221 DAVIDSOHN, I, STERN, K, STRAUSSER, F R, AND SPURRIER W *Blood* 8 747 (1953)
- 222 HEKTOEN, L, LANDSTEINER, K, AND WIENER A S *J Amer Med Assoc* 108 2138 (1937)
- 223 DAVIDSOHN, I, IRVINE P, AND WIENER A S *J Amer Med Assoc* 148 699 (1952)
- 224 WIENER, A S, WEYLER I B, STORMONT C AND OWEN, R *J Amer Med Assoc*, 161 233 (1956), 164 2036 (1957)
- 225 SCHATKIN, S B *Disputed Paternity Proceedings* 3rd edition, Matthew Bender & Co, N Y, 1952
- 226 WIFNER, A S, LEDERER, M AND POLAYES S H *J Immunol*, 19 250 (1930)
- 227 WIENER, A S *J Immunol*, 24 413 (1933)
- 228 WIFNER, A S *Amer J Human Genet* 2 177 (1950)
- 229 WIENER, A S *The 1950 Proceedings of the International Society of Hematology* pp 207 220, Grune & Stratton New York
- 230 BOYD, W C *Amer J Human Genet* 7 229 (1955)
- 231 WIENER A S *Ztschr f indukt Abstammungs u Vererbungs* 59 227 (1931)
- 232 WIFNER, A S *Amer J Hum Genet* 2 177 (1950)
- 232a ALVAREZ JOSÉ DE JS Aplicaciones Médico legales y Antropologicas de los Grupos Sanguíneos en la Republica Dominicana VI Congreso Médico Dominicano Trujillo 1951
- 233 WIFNER A S, OWEN R D, STORMONT C, AND WEYLER I B *J Amer Med Assoc* 164 2036 (1957)
- 234 HEKTOEN, L, LANDSTEINER K, AND WIFNER A S *J Amer Med Assoc* 108 2138 (1937)
- 235 WIFNER A S *J Forensic Med* 3 139 (1956)
- 236 LANDSTEINER, K, WIENER A S, AND MATSON G A *J Exp Med* 76 73 (1912)
- 237 WIFNER A S, SONN F B AND BEKIN, R B *Proc Soc Exp Biol and Med*, 64 316 (1913)
- 238 WIFNER A S, ZEPEDA J P, SONN, F B, AND POLIYKA H *J Exp Med*, 81. 579 (1945)
- 239 WIENER A S *Science*, 96 407 (1942)
- 240 .
- 241 . (1911)
- 242 ..

- 243 CANDELA P B *Hum Biol* 14 413 (1942)
- 244 MOURANT A E *Cold Spr Harb Symp Quant Biol* 15 291 (1950)
- 245 CHALMERS J N M IRIN E W AND MOURANT A E *Ann Eugen (Lond)* 17 168 (1953)
- 246 BOYD W C *Blood groups Tabul biol (Hague)* 17 113-240 (1939)
- 247 WIENER A S *Blood Groups and Transfusion* 3rd ed Thomas Springfield 1943
- 248 MOURANT A F *The Distribution of the Human Blood Group* Blackwell Oxford 1954
- 249 WIENER A S AND LEFF I L *Genetics* 25 187-196 (1940)
- 250 LINGER L J *J Amer Med Assoc* 152 1006-1010 (1953)
- 251 SUSSMAN I N AND SCHATKIN S B *J Amer Med Assoc* 164 249-250 (1957)
- 252 SUSSMAN I N AND MILLER E B *Rev d Hematol* 7 368-371 (1952)
- 253 SCHATKIN S B SUSSMAN I N AND YARBROUCH D R *Crin Lau Rev* 2 (1) 44-56 (1956)
- 254 WYELEGALA V B AND WITEBSKY F *Buffalo La Rev* 7 (2) 209-230 (1958)
- 255 WITEBSKY E KLENSHOF N C AND McNEILL C *Proc Soc Exp Biol & Med* 55 167 (1944)
- 256 WALSH R J KOOPITZ OF O LANCASTER H Q AND PRICE A V C *Oceania* 24 146-151 (1953)
- 257 OTTENSOOSER F AND PASQUALIN R *Amer J Hum Genet* 1 141-150 (1949)
- 258 SIMMONS R T GRAYDON J J SEMPLE N M AND FRY F I *Amer J Phys Anthropol* 13 667-690 (1955)
- 259 SIMMONS R T GRAYDON J J AND SEMPLE N M *Med J Austral* 2 589-596 (1953)
- 260 SIMMONS R T *Anthropos* 51 500-512 (1956)
- 261 MOLLIS N P L *Blood Transfusion in Clinical Medicine* 2nd ed Blackwell Oxford 1956
- 262 HARLEY D *Mexico Legal Blood Group Determination* W He nemann London 1943
- 263 CRADWELL R B H *Clinical Laboratory Methods and Diagnosis* 5th ed chapter V pp 994-1189 C V Mosby St Louis 1956
- 264 GONZALES T A VANCE M HELPERN M AND UMBERGER C J *Legal Medicine Pathology and Toxicology* 2nd ed chapter 27 pp 634-673 Appleton Century Crofts New York 1954
- 265 COLIN E C *Elements of Genetics* 3rd ed pp 108-117 McGraw Hill New York 1956

INDEX

- A B O groups, 16 *ff*
 - A_m blood groups, 38
 - A_m blood group, 27
 - Bernstein's theory of heredity, 12
 - "Bombay" blood group, 39
 - C factor, 26 *ff*.
 - discovery of, 1, 4
 - distribution of, 19
 - family studies, 18, 22
 - gene frequency analysis, 16
 - heredity of, 12, 16 *ff*
 - homozygous and heterozygous, 20
 - in apes and monkeys, 38
 - medicolegal applications, 118 *ff*
 - mother child combinations, 21
 - paternity, changes of excluding 120
 - secretor types, 29 *ff*
 - serology of 26 *ff*
 - subgroups of A, 21 *ff*
- Agglutination reaction, 11
- Agglutinogens, 2, 10
 - and blood factors, 10
 - mosaic nature of, 11
 - A B O, 28 29
 - M N S, 48, 55
 - P, 58
 - Rh Hr, 75, 81, 90
- Alleles, multiplicity of, 14
- A_m blood group, 38
- Anthropoid apes 38, 47
 - A B O groups in 38
 - M factors in, 47
- Anthropology, blood groups and 132 *ff*
- Antibodies, 10
 - anti A and anti B, 16
 - anti A₁, 23
 - anti C, 27
 - anti-d, 94
 - anti F, 111
 - anti H or anti O, 21, 30
 - anti hr, 80, 81, 82, 83
 - anti hr', 71, 79
 - anti hr'', 73, 79
 - anti Hr_o, 7, 73, 89
 - anti hr'', 81
 - anti K and anti k, 103 *ff*
 - anti Kp^a and anti Kp^b, 106 107
 - anti I c, 32
 - anti Ia, 108
 - anti M and anti N, 41
 - anti O or anti H, 21, 30
 - anti P, 57
 - anti Rh^A, anti Rh^B, etc., 90
 - anti Rh_o, 60 *ff*
 - anti rh', 63 *ff*
 - anti rh'', 64 *ff*
 - anti rh^G, 91
 - anti rh'', 80, 82 83
 - anti rh^s, 91
 - anti S, 57
 - linked 101
- Antigen antibody combination, 10
- Anti-serums, sources of, 2
- Basques, blood groups in, 135
- Becker blood factor 115
- Bernstein's theory, 12 *ff*
- Berrens blood factor, 115
- Bi, B_{ii}, B_{iii}, blood factors 27, 28 29
- Blood factors, 3 *ff*
 - definition of 10
 - high frequency, 117
 - list of, 4 *ff*
 - low frequency, 118
 - test for independence of, 3
 - variants of, 14
- Blood groups
 - history of, 4 *ff*
 - nomenclature of, 4 *ff*, 14
 - technique of tests 10
- Blood group chimeras, 33
- Blood group serology, principles of, 1
- Blood group systems, list of, 4 *ff*
- Body fluids, group substances in 29
- "Bombay" blood type, 39

- C blood factor 26 ff 29
 Ca blood factor 115
 C D E notations 92 ff
 Cellano blood factor 104
 Ceppellini's hypothesis 36
 Chimpanzees 38
 A B O blood groups in 38
 M blood factors in 47
 Chi square test 3
 M N types 44
 Contingency table 3
 M N S types 49
 Crossing over 9a
 Deletions " 97
 Diego blood groups 114
 Distribution in population of
 A B O groups 19
 Duffy groups 111
 Kell groups 103
 Lewis groups 35
 P groups 57
 Rh Hr groups 67 82
 Duffy blood groups 110 ff
 F and f blood factors 110 ff
 FA blood factor 28 29
 Fisher Race hypothesis 92 ff
 notations 92 ff
 Furukata's linkage hypothesis, 12
 Gene dose effect 73 79
 Genetics of blood groups 4 ff
 A B O groups 16 ff
 Duffy blood groups 111
 eight Rh blood types 64 ff
 Kell blood groups 103 ff
 Lewis blood groups 34 ff
 Lutherin blood groups 108
 M N S types 49 ff
 P blood groups 58
 principles of 12 ff
 Rh₀ blood factor 61
 Rh Hr blood types, 71 ff, 82, 83
 secretor types, 31
 subgroups of A, 21 ff
 three M N types 42 ff
 Gorillas A B O groups in 38
 Gr blood factor 56
 Group specific substances 29 ff
 Grubb and Morgan hypothesis 3
 H blood factor 4 24 29
 and Lewis blood factors 34 3
 He blood factor 55
 Heterozygotes
 in group A and group B 20
 maximum frequency of 45
 selection against 131
 Hierarchy of the blood groups 4 ff
 Homozygote 20
 Hr blood factors 71 ff
 nomenclature of 74
 relation to Rh factors 74
 Hr₀ factor 71 8)
 hr factor 80
 hr factor 72
 hr' factor 73
 hr' factor 81
 Hu blood factor 53
 I blood factor 117
 Interchange of babies 121
 International nomenclature 82 83
 J and j blood factors 112 ff
 Jobbins blood factor 115
 K and k blood factors, 103 104
 Kell blood groups 103 ff
 Cellano blood factor 106
 distribution 103
 heredity 104
 Pennes blood factor 106
 serology, 107
 Kidd blood groups 112 113
 Kp^a and Kp^b blood factors 106
 Landsteiner, Karl, *frontispiece*
 Landsteiner's A-B-O groups, 16 ff
 Landsteiner and Levine's theory
 42 ff
 Landsteiner and Wiener

- discovery of Rh factor, 6
 heredity of Rh factor 60 61
 I^e and I^e_H blood factors 32
 Lectins 21
 Le⁺ blood factor, 115
 Lewis blood groups 34 ff
 and T substance, 34
 and H blood factor 34
 family studies 32
 genetics of 34 ff
 in Bombay' type 39
 in twins 33
 Linked gene hypothesis 12
 A B O groups 12
 Rh Hr blood types 92 ff 100
 Lutheran blood groups 108 109
 Maternity exclusion by blood tests 123
 Medicolegal applications 117 ff
 illustrative cases 125 ff
 maternity exclusion of, 123
 nomenclature in 128
 paternity exclusion of 118
 qualification of experts 120
 M, M₁, M₂, blood factors 47 49
 M₁ blood factor 50
 M N types 41 ff
 chi square test 44
 family studies 43
 gene frequency analysis 43
 heredity of 42 ff
 paternity chances of excluding 120
 serology of 41 46
 M N S types 48 ff
 distribution 41
 factors Gr and Me, 54
 factors Hu and He, 55
 factor U 54
 family studies 53
 gene frequency analysis 52
 genetic theory 49
 gene *le* 50
 Monkey A B O groups in 38
 M factors in 47
 Mosaic nature of agglutinogens 11
 A B O agglutinogens 29
 Kell agglutinogens 107
 M N agglutinogens 48
 Rh agglutinogens 84 ff
 Rh Hr agglutinogens 63, 68 81
 Mother child combinations, 21 45
 Multiple allele theory 12 ff
 A B O groups 12 16 ff
 Kell blood groups 106 107
 P blood groups 58
 Rh Hr blood types 63 66 ff 82 83
 Subgroups of A 21
 N₁ N₂ blood factors 48
 N agglutininogen 46
 Nomenclature of blood groups 14
 A B O groups 27 29
 American Medical Association report 102
 C D E notations 100
 Hr factors 74
 in medicolegal practice 128
 international 82 83
 Kell types 105
 Lewis types 31
 M N S types 51
 Rh₀ variants 80
 Rh Hr types 82 83
 Non secretors 29
 and I^e(a+) type 32
 T substance in saliva 34
 O blood factor 20
 O₁ blood group 39
 Parentage exclusion of by blood tests 124
 Paternity chances of excluding by blood tests 118 ff
 P blood groups 57 ff
 Penny blood factor 10
 Position effect 97
 Principles of
 blood group genetics 12
 blood group nomenclature 14
 blood grouping tests 1 10 ff
 Q blood factor, 57

INDIX

- anti Hr serums 79
 - Duffy blood groups 110
 - eight Rh types 70
 - Kell blood groups 107
 - Kidd blood groups 117
 - P blood groups 57
 - Rh₀ blood factor 61
 - Rh₀ and Hr₀ blood factors 88 ff
 - S₁ blood factor 103
 - Subgroup of A 21 ff
 - A₂ and A₂B 76
 - family studies 24
 - gene frequency analysis 71
 - serology of 23
 - Super Rh₀ 87
- Technique of blood grouping 7 10
 - Tj₁ blood factor 58
 - Transubstance 34
 - Twin mosaics 33
- U blood factor 54
- Variants of Rh₀ factor 84 ff
 - Vel blood factor 110
 - Ven blood factor 9
 - Vw blood factor 9
- Wiener nomenclature 66 74 82 83
 - Wiener's theory of heredity of the
 - Rh Hr blood types 13 13 ff 87
 - 83 84 ff
 - Wra blood factor 115
 - Yt₁ blood factor 115
- Yt₂ blood factor 115
- Yt₃ blood factor 115
- Yt₄ blood factor 115
- Yt₅ blood factor 115
- Yt₆ blood factor 115
- Yt₇ blood factor 115
- Yt₈ blood factor 115
- Yt₉ blood factor 115
- Yt₁₀ blood factor 115
- Yt₁₁ blood factor 115
- Yt₁₂ blood factor 115
- Yt₁₃ blood factor 115
- Yt₁₄ blood factor 115
- Yt₁₅ blood factor 115
- Yt₁₆ blood factor 115
- Yt₁₇ blood factor 115
- Yt₁₈ blood factor 115
- Yt₁₉ blood factor 115
- Yt₂₀ blood factor 115
- Yt₂₁ blood factor 115
- Yt₂₂ blood factor 115
- Yt₂₃ blood factor 115
- Yt₂₄ blood factor 115
- Yt₂₅ blood factor 115
- Yt₂₆ blood factor 115
- Yt₂₇ blood factor 115
- Yt₂₈ blood factor 115
- Yt₂₉ blood factor 115
- Yt₃₀ blood factor 115
- Yt₃₁ blood factor 115
- Yt₃₂ blood factor 115
- Yt₃₃ blood factor 115
- Yt₃₄ blood factor 115
- Yt₃₅ blood factor 115
- Yt₃₆ blood factor 115
- Yt₃₇ blood factor 115
- Yt₃₈ blood factor 115
- Yt₃₉ blood factor 115
- Yt₄₀ blood factor 115
- Yt₄₁ blood factor 115
- Yt₄₂ blood factor 115
- Yt₄₃ blood factor 115
- Yt₄₄ blood factor 115
- Yt₄₅ blood factor 115
- Yt₄₆ blood factor 115
- Yt₄₇ blood factor 115
- Yt₄₈ blood factor 115
- Yt₄₉ blood factor 115
- Yt₅₀ blood factor 115
- Yt₅₁ blood factor 115
- Yt₅₂ blood factor 115
- Yt₅₃ blood factor 115
- Yt₅₄ blood factor 115
- Yt₅₅ blood factor 115
- Yt₅₆ blood factor 115
- Yt₅₇ blood factor 115
- Yt₅₈ blood factor 115
- Yt₅₉ blood factor 115
- Yt₆₀ blood factor 115
- Yt₆₁ blood factor 115
- Yt₆₂ blood factor 115
- Yt₆₃ blood factor 115
- Yt₆₄ blood factor 115
- Yt₆₅ blood factor 115
- Yt₆₆ blood factor 115
- Yt₆₇ blood factor 115
- Yt₆₈ blood factor 115
- Yt₆₉ blood factor 115
- Yt₇₀ blood factor 115
- Yt₇₁ blood factor 115
- Yt₇₂ blood factor 115
- Yt₇₃ blood factor 115
- Yt₇₄ blood factor 115
- Yt₇₅ blood factor 115
- Yt₇₆ blood factor 115
- Yt₇₇ blood factor 115
- Yt₇₈ blood factor 115
- Yt₇₉ blood factor 115
- Yt₈₀ blood factor 115
- Yt₈₁ blood factor 115
- Yt₈₂ blood factor 115
- Yt₈₃ blood factor 115
- Yt₈₄ blood factor 115
- Yt₈₅ blood factor 115
- Yt₈₆ blood factor 115
- Yt₈₇ blood factor 115
- Yt₈₈ blood factor 115
- Yt₈₉ blood factor 115
- Yt₉₀ blood factor 115
- Yt₉₁ blood factor 115
- Yt₉₂ blood factor 115
- Yt₉₃ blood factor 115
- Yt₉₄ blood factor 115
- Yt₉₅ blood factor 115
- Yt₉₆ blood factor 115
- Yt₉₇ blood factor 115
- Yt₉₈ blood factor 115
- Yt₉₉ blood factor 115
- Yt₁₀₀ blood factor 115

